

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
21 June 2001 (21.06.2001)

PCT

(10) International Publication Number
WO 01/44239 A2

(51) International Patent Classification⁷: **C07D 413/12**,
413/14, 403/14, 487/04, 261/16, 401/12, 417/12, A61K
31/42, A61P 9/12

John, E. [US/US]; 258 Old Farm Road, Flemington, NJ
08822 (US). **GU, Zhengxiang** [CN/US]; 92 York Road,
Princeton, NJ 08540 (US).

(21) International Application Number: PCT/US00/33730

(74) Agents: **ALGIERI, Aldo, A.** et al.; Bristol-Myers Squibb
Co., P.O. Box 4000, Lawrenceville-Princeton Road,
Princeton, NJ 08543 (US).

(22) International Filing Date:
13 December 2000 (13.12.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/464,037 15 December 1999 (15.12.1999) US
09/481,197 11 January 2000 (11.01.2000) US
09/513,779 25 February 2000 (25.02.2000) US
09/604,322 26 June 2000 (26.06.2000) US
09/643,640 22 August 2000 (22.08.2000) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE,
DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID,
IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT,
LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL,
PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ,
UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **BRIS-
TOL-MYERS SQUIBB CO.** [US/US]; P.O. Box 4000,
Lawrenceville-Princeton Road, Princeton, NJ 08543-4000
(US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **MURUGESAN,
Natesan** [US/US]; 42 Millbrook Drive, Princeton Junc-
tion, NJ 08550 (US). **TELLEW, John, E.** [US/US]; 28
Avalon Road, Pennington, NJ 08534 (US). **MACOR,**

Published:

— Without international search report and to be republished
upon receipt of that report.

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: BIPHENYL SULFONAMIDES AS DUAL ANGIOTENSIN ENDOTHELIN RECEPTOR ANTAGONISTS

(57) Abstract: Novel biphenyl sulfonamide compounds which are combined angiotensin and endothelin receptor antagonists are claimed along with methods of using such compounds in the treatment of conditions such as hypertension and other diseases, as well as pharmaceutical compositions containing such compounds.




WO 01/44239 A2


BIPHENYL SULFONAMIDES AS
DUAL ANGIOTENSIN ENDOTHELIN RECEPTOR ANTAGONISTS


The present invention relates to biphenyl sulfonamide compounds which are combined angiotensin and endothelin receptor antagonists, to methods of using such compounds in the treatment of conditions such as hypertension and other diseases, and to pharmaceutical compositions containing such compounds.

The present invention provides biphenyl sulfonamide compounds of the following formula I, enantiomers (including atropisomers), diastereomers, salts and metabolites thereof:



wherein:







R₂ is hydrogen, halogen, -CHO, alkyl, haloalkyl, (cycloalkyl)alkyl, alkenyl,
 5 alkylnyl, alkoxyalkyl, haloalkoxyalkyl, alkoxy, aryloxy alkoxyalkoxy,
 cyano, hydroxy, hydroxyalkyl, nitro, -CH(OR₁₃)(OR₁₄), -(CH₂)_wY;
 with the proviso that when R₁ is B, R₂ is not hydrogen, halogen,
 alkyl, haloalkyl, alkoxy, hydroxyalkyl, nitro, -(CH₂)_wNR₁₉R₂₀ or
 -NHSO₂R₂₂;


10 R₃ is heteroaryl;

R₄ and R₅ are each independently alkyl, hydroxyalkyl, cycloalkyl, hydroxy
 substituted cycloalkyl, alkoxyalkyl, or hydroxy substituted
 alkoxyalkyl, or R₄ and R₅ together form a cyclobutyl, cyclopentyl,
 cyclohexyl, tetrahydrofuranyl or tetrahydropyranyl ring which may
 15 be optionally substituted with one or more hydroxy groups;

R₆ is alkyl, hydroxyalkyl, haloalkyl, hydroxy substituted haloalkyl,
 cycloalkyl, hydroxy substituted cycloalkyl, (cycloalkyl)alkyl,
 hydroxy substituted (cycloalkyl)(alkyl), aralkyl, alkoxy, hydroxy
 substituted alkoxy, alkoxyalkyl, hydroxy substituted alkoxyalkyl, or
 20 -NR₁₆R₁₇;


- R_7 is $-(CH_2)_w-CO_2R_{15}$, $-(CH_2)_w-(C=O)NR_{16}R_{17}$, $-(CH_2)_w-NR_{15}(C=O)NR_{16}R_{17}$, $-(CH_2)_w-CH_2OH$, $-(CH_2)_w-(C=O)R_{15}$, tetrazolyl, oxadiazolyl or triazolyl wherein said tetrazolyl, oxadiazolyl or triazolyl may optionally be substituted with hydrogen, alkyl, hydroxy or halogen;
- 5 R_8 , R_9 , R_{9a} , R_{10} and R_{12} are each independently hydrogen, halogen, alkyl, hydroxyalkyl, cycloalkyl, (cycloalkyl)alkyl, aryl, heteroaryl, arylalkyl, alkylthioalkyl, alkoxy or alkoxyalkyl, or R_9 and R_{9a} together with the carbon atom to which they are bonded form a cycloalkyl ring;
- 10 R_{11} and R_{11a} are each independently hydrogen, alkoxy, or together form a carbonyl;
- R_{13} and R_{14} are alkyl or together form a five to six-membered ring;
- R_{15} , R_{16} and R_{17} are independently hydrogen, alkyl, hydroxyalkyl, cycloalkyl, (cycloalkyl)alkyl, alkoxyalkyl, aralkyl, heterocycloalkyl,
- 15 aryl, heteroaryl or $(CH_2)_wQ$, or R_{16} and R_{17} may together form a four to six-membered heterocyclic ring;
- n is 1 or 2;
- w is 0, 1, or 2;
- Y is heteroaryl, $-COOH$, $-COOR_{18}$, $-CONR_{19}R_{20}$, $-NR_{19}R_{20}$, $-NR_{19}-OR_{20}$, -
- 20 $NR_{21}(C=O)R_{22}$, $-NR_{21}(C=O)NR_{19}R_{20}$, $-N(R_{19})-(alk)-NR_{21}(C=O)R_{22}$, $-NR_{21}(C=O)OR_{18}$, $-NR_{21}SO_2R_{22}$, $-SO_2R_{22}$, Q , R or S ;





R_{18} , R_{19} , R_{20} , R_{21} and R_{22} are each independently hydrogen, alkyl, haloalkyl, alkoxyalkyl, cycloalkyl, alkenyl, alkynyl, aryl, aralkyl, heteroaryl, or R_{19} and R_{20} may together form a four to seven-membered heterocyclic ring;

R_{23} and R_{24} are each independently hydrogen, alkyl or cycloalkyl, or may together form a three to seven membered cycloalkyl ring;



x is 2, 3 or 4;


R_{25} , R_{26} and R_{27} are each independently hydrogen, alkyl or cycloalkyl, or R_{26} and R_{27} may together form a three to seven-membered cycloalkyl ring;

R_{101} , R_{102} , R_{103} , and R_{104} are each independently hydrogen, halogen, -CHO, alkyl, haloalkyl, (cycloalkyl)alkyl, alkenyl, alkynyl, alkoxyalkyl, haloalkoxyalkyl, alkoxy, alkoxyalkoxy, cyano, hydroxy, hydroxyalkyl, nitro, -CH(OR₁₃)(OR₁₄), or -(CH₂)_wY; wherein said rings; aryl alone or as part of another group; or heteroaryl alone or as part of another group may each optionally be substituted by one or more hydrogen, halogen, cyano, alkyl, hydroxyalkyl, alkoxy, nitro or trifluoromethyl groups.

The compounds of the formula I and salts thereof may be used as combined endothelin and angiotensin receptor antagonists.


Compounds of the formula I and salts thereof wherein one or more, and especially all, of R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , R_8 , R_9 , R_{10} , R_{11} , R_{11a} , R_{12} , R_{15} , R_{16} , R_{17} , R_{18} , R_{19} , R_{20} , R_{21} , R_{22} , R_{23} , R_{24} , R_{25} , R_{26} , R_{27} , R_{101} , R_{102} , R_{103} , R_{104} , n , w , Y ,

Q, Z, and x are selected from the following definitions, are preferred compounds of the present invention:




- 5 R_2 is alkyl, haloalkyl, (cycloalkyl)alkyl, alkoxyalkyl, haloalkoxyalkyl, alkoxy, alkoxyalkoxy, hydroxyalkyl, or $-(CH_2)_wY$, or when R_1 is D, R_2 is hydrogen, alkyl, haloalkyl, (cycloalkyl)alkyl, alkoxyalkyl, haloalkoxyalkyl, alkoxy, alkoxyalkoxy, hydroxyalkyl, or $-(CH_2)_wY$;
- 10 R_3 is isoxazolyl, pyridizynyl, pyrazynyl or pyrimidinyl, each optionally substituted with one to three of the following substituents:
hydrogen, halogen, cyano, alkyl, alkoxy, trifluoromethyl or nitro;
 R_4 and R_5 are each independently alkyl, cycloalkyl, or R_4 and R_5 together form a cyclobutyl, cyclopentyl or cyclohexyl ring;
- 15 R_6 is alkyl, haloalkyl, cycloalkyl or alkoxy;
 R_7 is $-CO_2R_{15}$, $-(C=O)NR_{16}R_{17}$ or $-CH_2OH$;
 R_8 , R_9 , R_{10} and R_{12} are each independently hydrogen, halogen, alkyl, cycloalkyl, alkoxy or alkoxyalkyl;
 R_{11} and R_{11a} are each independently hydrogen, alkoxy, or together form a carbonyl;
- 20 R_{15} , R_{16} and R_{17} are independently hydrogen, alkyl or cycloalkyl or R_{16} and R_{17} may together form a four to six-membered heterocyclic ring;
n is 1 or 2;
w is 0, 1, or 2;

Y is $-\text{COOR}_{18}$, $-\text{NR}_{21}(\text{C}=\text{O})\text{R}_{22}$, $-\text{NR}_{21}(\text{C}=\text{O})\text{NR}_{19}\text{R}_{20}$, $-\text{NR}_{21}(\text{C}=\text{O})\text{OR}_{18}$,
 $-\text{NR}_{21}\text{SO}_2\text{R}_{22}$, $-\text{SO}_2\text{R}_{22}$ or $\underline{\text{Q}}$;



R_{18} , R_{19} , R_{20} , R_{21} and R_{22} are each independently hydrogen, alkyl, cycloalkyl,
 5 or R_{19} and R_{20} may together form a four to seven-membered
 heterocyclic ring;

R_{23} and R_{24} are each independently hydrogen, alkyl or cycloalkyl, or may
 together form a three to seven membered cycloalkyl ring;




10 x is 2, 3 or 4;

R_{25} , R_{26} and R_{27} are each independently hydrogen, alkyl or cycloalkyl, or R_{26}
 and R_{27} may together form a three to seven-membered cycloalkyl
 ring;

R_{101} , R_{102} , R_{103} , and R_{104} are each independently hydrogen, halogen, alkoxy or
 15 alkyl.

Compounds of the formula I and salts thereof wherein one or more,
 and especially all, of R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , R_8 , R_9 , R_{10} , R_{11} , R_{11a} , R_{15} , R_{16} ,
 R_{17} , R_{18} , R_{19} , R_{20} , R_{21} , R_{22} , R_{23} , R_{24} , R_{25} , R_{26} , R_{27} , R_{101} , R_{102} , R_{103} , R_{104} , n, w, Y, $\underline{\text{Q}}$,
 Z, and x are selected from the following definitions, are more preferred
 20 compounds of the present invention:



R_2 is alkyl, haloalkyl, (cycloalkyl)alkyl, alkoxyalkyl, haloalkoxyalkyl, alkoxy, hydroxyalkyl, or $-(CH_2)_wY$; or when R_1 is D, R_2 is hydrogen, alkyl, haloalkyl, (cycloalkyl)alkyl, alkoxyalkyl, haloalkoxyalkyl, alkoxy, alkoxyalkoxy, hydroxyalkyl, or $-(CH_2)_wY$;

R_3 is isoxazolyl, optionally substituted with one or two of the following substituents: hydrogen, halogen, cyano, alkyl, alkoxy, trifluoromethyl or nitro;

R_4 and R_5 are each independently alkyl, cycloalkyl, or R_4 and R_5 together form a cyclobutyl, cyclopentyl or cyclohexyl ring;

R_6 is alkyl, haloalkyl, cycloalkyl or alkoxy;

R_7 is $-CO_2R_{15}$ or $-(C=O)NR_{16}R_{17}$;

R_8 , R_9 and R_{10} are each independently hydrogen, halogen, alkyl, cycloalkyl alkoxy or alkoxyalkyl;


R_{11} and R_{11a} together form a carbonyl;

R_{15} , R_{16} and R_{17} are independently hydrogen, alkyl, or cycloalkyl or R_{16} and R_{17} may together form a four to six-membered heterocyclic ring;

n is 2;


w is 0, 1, or 2;

Y is $-NR_{21}(C=O)R_{22}$, $-NR_{21}(C=O)NR_{19}R_{20}$, $-NR_{21}(C=O)OR_{18}$, $-NR_{21}SO_2R_{22}$, $-SO_2R_{22}$ or Q;



R_{18} , R_{19} , R_{20} , R_{21} and R_{22} are each independently hydrogen, alkyl, cycloalkyl, or R_{19} and R_{20} may together form a four to seven-membered heterocyclic ring;

- 5 R_{23} and R_{24} are each independently hydrogen, alkyl or cycloalkyl, or may together form a three to seven membered cycloalkyl ring;




x is 2, 3 or 4;

- R_{25} , R_{26} and R_{27} are each independently hydrogen, alkyl or cycloalkyl, or R_{26} and R_{27} may together form a three to seven-membered cycloalkyl ring;

R_{101} , R_{102} , R_{103} , and R_{104} are each independently hydrogen, halogen, or alkyl.

- Compounds of the formula I and salts thereof wherein one or more, and especially all, of R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_8 , R_{10} , R_{11} , R_{11a} , R_{18} , R_{19} , R_{20} , R_{21} , R_{22} , R_{23} , R_{24} , R_{25} , R_{26} , R_{27} , R_{101} , R_{102} , R_{103} , R_{104} , w, Y, Q, Z, and x are selected from the following definitions, are most preferred compounds of the present invention:



R_2 is alkyl, haloalkyl, (cycloalkyl)alkyl, alkoxyalkyl, haloalkoxyalkyl, alkoxy, alkoxyalkoxy, hydroxyalkyl, or $-(CH_2)_wY$ or when R_1 is D, R_2 is hydrogen, alkyl, haloalkyl, (cycloalkyl)alkyl, alkoxyalkyl, haloalkoxyalkyl, alkoxy, alkoxyalkoxy, hydroxyalkyl, or $-(CH_2)_wY$;

5

R_3 is isoxazol-5-yl or isoxazol-3-yl independently substituted with two of following substituents: alkyl or halogen;

R_4 and R_5 are each independently alkyl, cycloalkyl, or R_4 and R_5 together form a cyclobutyl, cyclopentyl or cyclohexyl ring;

10 R_6 is alkyl, halo alkyl, cycloalkyl or alkoxy;


R_7 is $-CO_2R_{15}$ or $-(C=O)NR_{16}R_{17}$;

R_8 , R_9 , and R_{10} are independently H, alkyl, cycloalkyl, alkoxy or alkoxyalkyl;

n is 2;

15 w is 0, 1, or 2;


Y is $-NR_{21}(C=O)R_{22}$, $-NR_{21}(C=O)NR_{19}R_{20}$, $-NR_{21}(C=O)OR_{18}$, $-NR_{21}SO_2R_{22}$ or Q;



R_{18} , R_{19} , R_{20} , R_{21} and R_{22} are each independently hydrogen, alkyl, cycloalkyl, or R_{19} and R_{20} may together form a four to seven-membered

20 heterocyclic ring;

R_{23} and R_{24} are each independently hydrogen, alkyl or cycloalkyl, or may together form a three to seven membered cycloalkyl ring;



x is 2;

R_{25} , R_{26} and R_{27} are each independently hydrogen, alkyl or cycloalkyl, or R_{26} and R_{27} may together form a three to seven-membered cycloalkyl ring;

R_{101} , R_{102} , R_{103} , and R_{104} are each independently hydrogen, halogen, or alkyl.

5 Especially preferred are compounds where

R_1 is selected from A, D, or E;

R_2 is selected from alkyl, alkoxyalkyl, and haloalkoxyalkyl, and further selected from hydrogen when R_1 is D;

R_3 is isoxazol-3-yl independently substituted with two of following

10 substituents: alkyl or halogen;

R_4 and R_5 together form a cyclobutyl, cyclopentyl or cyclohexyl ring;

R_6 is alkyl;

R_7 is $-(C=O)NR_{16}R_{17}$;

R_8 , R_9 , and R_{10} are independently alkyl or alkoxy; and

15 R_{101} , R_{102} , R_{103} , and R_{104} are each independently hydrogen, halogen, or alkyl.

20 The following are definitions of terms used in this specification.

The initial definition provided for a group or term herein applies to that group or term throughout the present specification, individually or as part of another group, unless otherwise indicated.

25 The terms "alk" or "alkyl" refer to straight or branched chain hydrocarbon groups having 1 to 12 carbon atoms, preferably 1 to 8 carbon atoms. Lower alkyl groups, that is, alkyl groups of 1 to 4 carbon atoms, are most preferred.

The term "alkenyl" refers to straight or branched chain hydrocarbon groups having 2 to 12 carbon atoms, preferably 2 to 4 carbon atoms, and
30 at least one double carbon to carbon bond, such as ethenyl.

The term "alkynyl" refers to straight or branched chain hydrocarbon groups having 2 to 12 carbon atoms, preferably 2 to 4 carbon atoms, and at least one triple carbon to carbon bond, such as ethynyl.

5 The term "alkoxy" refers to an alkyl group bonded through an oxygen (-O-).

The term "aryloxy" refers to an aryl group bonded through an oxygen (-O-).

The term "thioalkyl" refers to an alkyl group bonded through a sulfur (-S-).

10 The term "carbonyl" refers to the group $-(C=O)-$.

The terms "ar" or "aryl" refer to phenyl, naphthyl and biphenyl. Phenyl is a preferred aryl group. Aryl groups may be optionally substituted with one or more (such as one to three) of the following substituents: hydrogen, halogen, cyano, alkyl, alkoxy, nitro or
15 trifluoromethyl groups.

The term "cycloalkyl" refers to fully saturated cyclic hydrocarbon groups having 3 to 8 ring carbon atoms.

The terms "halogen" and "halo" refer to fluorine, chlorine, bromine and iodine. Haloalkyl refers to an alkyl chain substituted with from one
20 to three halogens.

The term "heteroaryl" refers to furyl, thienyl, pyrrolyl, pyridyl, pyrimidyl, pyrazinyl, pyridazinyl, triazinyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, oxadiazolyl, thiadiazolyl, imidazolyl, triazolyl, and tetrazolyl, each of which may optionally be substituted where appropriate by one or
25 more (such as one to three) of the following: hydrogen, halogen, cyano, alkyl, hydroxyalkyl, alkoxy, nitro or trifluoromethyl.

The terms "heterocyclic" or "heterocyclo" refer to optionally substituted, non-aromatic cyclic groups, for example, 4 to 7 membered monocyclic, 7 to 11 membered bicyclic, or 10 to 15 membered tricyclic ring
30 systems, which have at least one heteroatom in at least one carbon atom-containing ring. Each ring of the heterocyclic group containing a

heteroatom may have 1, 2, 3 or 4 heteroatoms selected from nitrogen atoms, oxygen atoms and/or sulfur atoms, where the nitrogen and sulfur heteroatoms may optionally be oxidized and the nitrogen heteroatoms may optionally be quaternized. The heterocyclic group may be attached at any
5 heteroatom or carbon atom of the ring or ring system.

Exemplary monocyclic heterocyclic groups include azetidiny, pyrrolidiny, oxetany, imidazolinyl, oxazolidiny, isoxazolinyl, thiazolidiny, isothiazolidiny, tetrahydrofurany, piperidiny, piperazinyl, 2-oxopiperazinyl, 2-oxopiperidiny, 2-oxopyrrolodiny, 2-oxoazepiny,
10 azepiny, 4-piperidony, tetrahydropyrany, morpholinyl, thiamorpholinyl, thiamorpholinyl sulfoxide, thiamorpholinyl sulfone, 1,3-dioxolane and tetrahydro-1,1-dioxothienyl, and the like.

The term "ring" encompasses homocyclic (i.e., as used herein, all the ring atoms are carbon) or "heterocyclic" (i.e., as used herein, the ring
15 atoms include carbon and one to four heteroatoms selected from N, O and /or S, also referred to as heterocyclo), where, as used herein, each of which (homocyclic or heterocyclic) may be saturated or partially or completely unsaturated (such as heteroaryl), and each of which (homocyclic or heterocyclic) may optionally be substituted by one or more (such as one to
20 three) hydrogen, halogen, cyano, alkyl, alkoxy, nitro or trifluoromethyl groups.

Throughout the specification, groups and substituents thereof may be chosen to provide stable moieties and compounds.

The compounds of formula I form salts which are also within the
25 scope of this invention. Reference to a compound of the formula I herein is understood to include reference to salts thereof, unless otherwise indicated. The term "salt(s)", as employed herein, denotes acidic and/or basic salts formed with inorganic and/or organic acids and bases. In addition, when a compound of formula I contains both a basic moiety and
30 an acidic moiety, zwitterions ("inner salts") may be formed and are included within the term "salt(s)" as used herein. Pharmaceutically

acceptable (i.e., non-toxic, physiologically acceptable) salts are preferred, although other salts are also useful, e.g., in isolation or purification steps which may be employed during preparation. Salts of the compounds of the formula I may be formed, for example, by reacting a compound I with an amount of acid or base, such as an equivalent amount, in a medium such as one in which the salt precipitates or in an aqueous medium followed by lyophilization.

The compounds of formula I which contain a basic moiety may form salts with a variety of organic and inorganic acids. Exemplary acid addition salts include acetates (such as those formed with acetic acid or trihaloacetic acid, for example, trifluoroacetic acid), adipates, alginates, ascorbates, aspartates, benzoates, benzenesulfonates, bisulfates, borates, butyrates, citrates, camphorates, camphorsulfonates, cyclopentanepropionates, digluconates, dodecylsulfates, ethanesulfonates, fumarates, glucoheptanoates, glycerophosphates, hemisulfates, heptanoates, hexanoates, hydrochlorides (formed with hydrochloric acid), hydrobromides (formed with hydrogen bromide), hydroiodides, 2-hydroxyethanesulfonates, lactates, maleates (formed with maleic acid), methanesulfonates (formed with methanesulfonic acid), 2-naphthalenesulfonates, nicotines, nitrates, oxalates, pectinates, persulfates, 3-phenylpropionates, phosphates, picrates, pivalates, propionates, salicylates, succinates, sulfates (such as those formed with sulfuric acid), sulfonates (such as those mentioned herein), tartrates, thiocyanates, toluenesulfonates such as tosylates, undecanoates, and the like.

The compounds of formula I which contain an acidic moiety may form salts with a variety of organic and inorganic bases. Exemplary basic salts include ammonium salts, alkali metal salts such as sodium, lithium, and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salts with organic bases (for example, organic amines) such as benzathines, dicyclohexylamines, hydrabamines (formed with

N,N-bis(dehydroabietyl)ethylenediamine), N-methyl-D-glucamines, N-methyl-D-glucamides, t-butyl amines, and salts with amino acids such as arginine, lysine and the like. Basic nitrogen-containing groups may be quaternized with agents such as lower alkyl halides (e.g. methyl, ethyl, 5 propyl, and butyl chlorides, bromides and iodides), dialkyl sulfates (e.g. dimethyl, diethyl, dibutyl, and diamyl sulfates), long chain halides (e.g. decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides), aralkyl halides (e.g. benzyl and phenethyl bromides), and others.

Prodrugs and solvates of the compounds of the invention are also 10 contemplated herein. The term "prodrug", as employed herein, denotes a compound which, upon administration to a subject, undergoes chemical conversion by metabolic or chemical processes to yield a compound of the formula I, or a salt and/or solvate thereof. Solvates of the compounds of formula I are preferably hydrates. Any tautomers which may exist are 15 also contemplated herein as part of the present invention.

All stereoisomers of the present compounds, such as those which may exist due to asymmetric carbons on the R substituents, including enantiomeric forms (which may exist even in the absence of asymmetric carbons, e.g., atropisomers) and diastereomeric forms, are contemplated 20 within the scope of this invention. Individual stereoisomers of the compounds of the invention may, for example, be substantially free of other isomers, or may be admixed, for example, as racemates or with all other, or other selected, stereoisomers. The chiral centers of the present invention can have the S or R configuration as defined by the IUPAC 1974 25 Recommendations.

The present invention can be applied to the extensive prior art in the field of angiotensin antagonists to obtain additional novel compounds possessing potent antagonist activity at both endothelin and angiotensin 30 receptors. In particular, a large number of groups (known to be useful

within the field of angiotensin receptor antagonists) can be substituted at the R₁ position of Formula I without departing from the scope of the present invention. The table below outlines examples of additional suitable R₁ groups.

5

Patent No.	Description of R₁
JP 09291078	Substituted imidazoles
JP 09301956	Substituted imidazoles
JP 09323991	carboxy methylidene cyclohepta imidazoles
WO 97/40040	Pyrimidin-4-ones
US 5674879	tetrahydro-imidazopyridines
WO 97/30036	N-acylaminoacid derivatives
JP 09110691	Imidazotetrahydropyridines
JP 08208640	Benzimidazoles
JP 08165292	purine derivatives
JP 08143552	Cyclohepta imidazoles
JP 08113572	Imidazopyridines
WO 96/10559	Ureas
EP 708103	Imidazolones
WO 96/08476	Pyrimidinones
JP 08041053	Pyrimidines
JP 08034780	Cycloheptimidazoles
WO 96/05195	naphtho-fused lactams
WO 96/04273	Pyrazoles
EP 696583	Benzimidazoles
JP 07316055	Pyrimidines
WO 95/34564	Pyridylimidazoles
JP 07309871	Hydrazotriazoles
WO 95/32198	Pyridylimidazoles
AU 95/16257	Imidazopyridines
WO 95/24902	Imidazoles
DE 4407488	Pyridones
WO 95/22543	Imidazoles
WO 95/21838	Pyridylimidazoles
JP 07157485	fused pyrimidinones
US 5411980	N-arylsubstituted-1,2,4-triazolinones
WO 95/16677	pyrimidine/pyrimidinones

WO 95/16675	Benzazepin-3-yl-ureas
WO 95/16692	Benzazepin-3-yl-ureas
US 5426105	Dihydroimidazopyridines
US 5424450	Imidazolinones
DE 4341453	4-oxo-imidazo-pyridines
JP 07112975	Amino-azoles
DE 4339868	4-oxo-imidazo-pyridazines
WO 95/12598	benzazepinyl urea macrocycles
EP 648763	Imidazoles
WO 95/09632	benzazepin-3-yl-ureas
EP 647627	pyridinones
JP 07048357	aminoacids
WO 95/03290	benzofused lactams
DE 4342724	2-oxo-1,2-dihydro-pyridines
WO 93/17023	pyrazolopyrimidines
US 5385894	6-aminoquinazolines
JP 07002776	N-arylmethylpyridines
JP 06340668	pyrazolotriazoles
US 5380719	quinoxalines and its N-oxides
GB 2280438	carboxymethylidene cycloheptimidazoles
WO/95/00517	imidazoles
US 5378704	benzo- and pyrido-1,2,4-thiadiazines
JP 06287182	alkylglycines
EP 623610	pyridine and pyridones
US 5358947	pyrazolotriazinones
WO 94/22838	pyrazoles
EP 621276	2,3,6-substituted quinazolinone derivatives
WO 94/21629	1-phenyl-imidazol-2-ones
JP 06211814	thioureas
EP 618207	5,8-dihydro-6H-pyrido-pyrimidin-7-ones
DE 4305279	triazolopyrimidines
WO 94/17069	pyrazoles
WO 94/17067	pyrimidones
US 5338736	2,3,6-substituted quinazolinone derivatives
JP 06184086	thioureas
DE 4300912	1-benzyl-1,2,3,4-tetrahydroquinazolin-2-ones or related analogs
US 5330987	pyridopyrimidinones
EP 607077	4-pyrimidinones
WO 94/13675	pyrazolotriazoles
US 5326776	5-membered heterocyclic rings comprising 1-4 N or

	2 N and 1 O and comprising 0-2 double bonds
EP 602521	imidazopyridines
WO 94/11379	pyridooxazinones
WO 94/11012	dipeptide analogs
EP 591891	benzimidazoles
US 5315013	pyrazoles
JP 06100541	pyrazolinones
WO 94/04516	5-membered ring heterocycles containing 1-4 heteroatoms
JP 06087833	vinyl imidazoles
EP 594022	2-pyridones
EP 594019	2-pyridones
JP 06073008	alkoxy-pyridines
JP 06072985	ureas and thioureas
DE 4233590	benzimidazoles
JP 06065236	dihydrobenzimidazolones
JP 06056826	quinoxalines
EP 589665	4,5,6,7-tetrahydro-1H-imidazo[4,5-c]pyridine-6-carboxamides
JP 06041122	quinoxalines
DE 4230464	imidazolylbenzimidazoles
US 5296480	2-alkyl-5,6,7,8-tetrahydro-pyrido (4,3-d)-pyrimidine-4-ones
US 5294617	quinazolinones
US 5294611	quinazolinones
US 5292734	quinazolinones
JP 06032782	pyrazoles
US 5290780	quinazolinones
JP 06025250	thieno[3,4-d]imidazoles
JP 06025229	imidazoles
WO 94/03453	pyrazolyl[4,3-c]pyridines
WO 94/03449	imidazoles
WO 94/03435	imidazole-5-carboxylic acids
US 5288720	quinazolinones
JP 06016661	imidazoles
US 5284853	quinazolinones
US 5284852	quinazolinones
JP 06009638	pyrazolopyrimidines
WO 94/02467	imidazoles
US 5281604	quinazolinones
US 5281603	quinazolinones

US 5281602	pyridopyrimidines
DE 4224133	benzimidazoles
US 5276048	imidazoles
GB 2268743	pyridines/pyrazines/pyrimidines
JP 05310696	pyridines
EP 574846	imidazopyridines
DE 4237656	benzimidazoles
FR 2690442	pyrazolopyrimidines
GB 2267089	cycloheptimidazoles
EP 573218	imidazoles
WO 93/23391	indoles, 7-azaindoles
US 5260325	N-linked amides or thioamides
DE 4212748	benzimidazoles
EP 566020	benzimidazoles
DE 4212250	benzimidazoles
US 5250548	aminopyridines
GB 2265900	pyrazoles
EP 562936	imidazolines
WO 93/18035	aza-phthalimides
WO 93/17682	bicyclic heterocycles
WO 93/17681	5-membered ring heterocycles
EP 561252	2-oxoquinolines
JP 05201994	pyridazinones
GB 2264709	imidazopyridines
JP 05194418	2-aminopyrimidines
WO 93/16049	4-oxo-dihydropyridines
JP 05178836	pyridines
EP 556080	pyrazolo[1,5-a]pyridines or imidazo[1.5-a]pyridines
US 5231094	triazolopyrimidines
EP 554107	1,2,4-triazole
EP 554098	imidazo[4,5b]pyridines
WO 93/14086	1-isoquinolones
EP 552765	benzimidazoles
JP 05155884	benzopyrazines or pyridopyrazines
WO 93/13077	hydantoins connected to imidazoles
US 5225408	oxadiazinone
JP 05140152	quinazolindione
EP 550313	pyrazolone and pyrimidinones
US 5219856	imidazoles
EP 547514	imidazopyridines
US 5218125	imidazoles

GB 2262096	4-aminopyrimidines
US 5214153	imidazoles
US 5212195	indoles, azaindoles
EP 543263	benzimidazoles
DE 4221583	pyridones
US 5208234	imidazolephosphonic acids
WO 93/08193	imidazo-tetrahydropyridazines
WO 93/08171	pyrimidocycloalkanes
WO 93/08169	aminopyrimidines
EP 539086	pyrimidinolactams
EP 537937	pyrazinopyrimidinones
DE 4132632	imidazolylpropenoic acids
EP 535465	imidazolylpropenoic acids
EP 535463	imidazolylpropenoic acids
EP 535420	imidazoles
EP 534706	quinazolinone or pyridopyrimidinones
WO 93/05044	pyrazolotriazines
WO 93/05025	pyrazoles
WO 93/04059	imidazoles
WO 93/04045	imidazolinones
JP 05032661	imidazolinones
EP 532410	imidazolinones and larger ring analogs
EP 531876	imidazoindolizines
EP 531874	4,5,6,7-tetrahydroimidazo[4,5-c]pyridine-4-carboxylic acids
EP 530702	1,2-dihydro-2-oxopyridines
WO 93/03040	thienopyrimidin-4-ones
WO 93/03033	imidazo[4,5-d]pyridazines
WO 93/03018	pyrimidines
US 5187168	quinazoline derivatives
JP 05017480	thienoimidazoles
US 5185340	oxypyrimidines
US 5182288	lactams
FR 2677016	acyl amino acid derivatives
WO 93/00341	imidazoles
WO 92/22533	4-aminoquinolines
US 5124335	fused pyrroles
EP 490820	acylamino derivatives
EP 490587	pyrazolo-pyrimidine and imidazopyridazines
EP 495626	oxypyridines, oxyquinolines, and imidazoles
EP 497150	3-quinazolin-4-ones

EP 497121	imidazoles
EP 498721	1,4-dihydroquinolin-4-ones
US 5132216	imidazopyridines
EP 499416	oxypyridines
EP 499415	aminopyridines
EP 499414	Fused oxypyridines
EP 500409	4-pyrimidones
EP 500297	2-pyridones and 2-pyrimidones
EP 502725	fused pyrimidones
EP 502575	1-(2H)-isoquinolinones
EP 502314	benzimidazoles
EP 505098	imidazoles
EP 503785	imidazoles
JP 04230683	7-aza and 4- azabenzimidazoles (i.e., imidazopyridines)
EP 507594	quinolines
JP 04235988	imidazopyridazinediones
EP 505893	imidazopyridines
FR 2672891	pyrazolones
US 5145699	pyridopyrimidines
EP 515265	pyrimidine derivatives
EP 511791	pyrrolopyridines
JP 04257564	benzimidazoles, imidazopyridines, imidazopyrazines
WO 92/16524	benzo-fused lactams
EP 516392	naphyridones and pyrido[c,b]pyrrolidones
WO 92/19211	imidazobenzoquinones
JP 04295478	imidazo[4,5-b]pridines
EP 5212768	triazolo pyrimidines
WO 92/21666	thiazole derivatives
EP 518033	imidazo[4,5-c]pyridine-4-carboxylates and other heterocycles
US 5087702	3H-imidazo-[4,5-b] pyridines
US 5087634	N-substituted imidazol-2-ones
WO 92/00977	imidazoles
EP 475206	various 6-membered heterocycles
WO 92/02508	oxyquinolines
WO 92/04343	tetrahydrobenzazoles
WO 92/04335	1H-1,2,4-triazoles
EP 475898	Azacyclic compounds including imidazolinones
EP 481448	dihydropyrimidines

QO 92/05161	1,3,5-trisubstituted 1,2,4-triazoles
WO 92/07852	xanthine derivatives
JP 04120072	pyrimidine derivatives
EP 487252	quinolines and 1,5-naphthyridine derivatives
EP 483683	thienoimidazoles
EP 470543	fused imidazoles
EP 468470	fused imidazoles
EP 467207	purines
WO 91/19715	imidazo[4,5-d]pyridazines
WO 91/19697	pyridines
EP 465368	imidazoles
EP 465323	pyrimidines
WO 91/18888	triazolones
EP 461039	benzimidazoles
US 5066586	pyridoimidazoles
EP 459136	benzimidazoles
WO 91/17148	triazoles
EP 456510	pyridoimidazoles
EP 456442	quinolines
WO 91/15479	imidazoles, oxazoles, thiazoles
WO 91/15209	pyrimidines
EP 453210	pyridines
WO 91/14679	imidazolinones or pyrimidinones
US 5053329	imidazopyridines
US 5049565	conjugated imidazopyridines
EP 449699	pyrazoles
EP 446062	pyrazoles
EP 445811	pyridin-4-ones and pyrimidin-4-ones
EP 443983	amides, sulfonamides, carbamates
EP 443568	thienopyridin-4-ones, thienopyrimidin-2,4-diones
EP 442473	pyrimidin-2,4-diones
EP 435827	pyrimidinones
EP 434038	fused imidazoles
EP 432737	cycloheptimidazolones
WO 91/07404	azaquinolines
EP 430300	xanthines
EP 426021	fused imidazoles
EP 425921	benzimidazoles
EP 424317	pyrimidines
EP 420237	fused imidazoles
EP 419048	pyrimidinones

EP 415886	fused imidazoles
EP 412848	oxyquinolines
EP 412594	triazolinones
EP 411766	quinazolinones
EP 411507	pyrazole-3-carboxylates
WO 91/00281	imidazoles
WO 91/00277	imidazoles
EP 409332	triazoles
EP 407342	pyrimidinones
EP 407102	fused imidazoles
EP 401030	fused imidazoles
EP 400974	fused imidazoles
EP 400835	benzimidazoles
EP 399732	benzimidazoles
EP 399731	fused imidazoles
EP 392317	benzimidazoles
EP 890719	imidazoles
EP 323841	pyrroles, pyrazoles, and triazoles
EP 291969	biphenyltetrazoles
EP 253310	imidazoles
WO 95/28419	indazoles
EP 638572	fused imidazoles
DE 4320432	aminopyridyl, imidazoles, fused imidazoles
JP 06279437	imidazoles
EP 624583	pyridones
EP 623611	pyridines or 2-pyridones
US 5348955	diacyl piperazines
WO 94/07486	benzo-fused lactams
JP 06128256	imidazopyridines
US 5298517	imidazoles
US 5286729	quinazolinones
DE 4203872	imidazo[1,2-a]pyridines
GB 2263637	fused imidazoles
US 5177097	imidazolones
EP 519831	imidazoles and pyrimidines
US 5153347	phosphonates or phosphinates
DE 4034728	thienoimidazoles
DE 4032522	thienoimidazoles
EP 461040	fused imidazoles
DE 4006693	benzimidazoles
US 4916129	imidazoles

US 4880804	benzimidazoles
WO 97/15556	3-spiroindolin-2-ones
US 5266583	imidazoles
EP 573218	imidazoles
US 5264447	imidazoles
EP 569794	benzopyridones or pyridopyridones
US 5091390	fused imidazoles
US 5256658	morpholines, piperidines, piperazines, thiomorpholines
EP 546358	benzimidazoles

Methods of Preparation

The compounds of the present invention may be prepared by
5 methods such as those illustrated in the following Schemes I to XIII.
Solvents, temperatures, pressures, and other reaction conditions may be
selected by one of ordinary skill in the art. Starting materials are
commercially available or readily prepared by one of ordinary skill in the
art.

10 The following are the definitions of symbols used throughout
Schemes I to XIII:


- AA hydrogen, halogen (chloro, bromo, iodo) or $-\text{OSO}_2\text{CF}_3$;
- BB suitable nitrogen protecting group, exemplified by
methoxymethyl- [MOM], benzyloxymethyl- [BOM], 2-
15 (trimethylsilyl)ethoxymethyl- [SEM], methoxyethoxymethyl-
[MEM], or *t*-butyl groups;
- DD $\text{S}_{\text{n}}2$ or $\text{S}_{\text{n}}1$ leaving group exemplified by halogen (Cl, Br, I) and
sulfonates ($-\text{OSO}_2$ -aryl (e.g., $-\text{OSO}_2\text{Ph}$ or $-\text{OSO}_2\text{PhCH}_3$), or
 $-\text{OSO}_2$ -alkyl (e.g., $-\text{OSO}_2\text{CH}_3$ or $-\text{OSO}_2\text{CF}_3$));
- 20 EE halogen (chloro, bromo, iodo) or $-\text{OSO}_2\text{CF}_3$;
- GG boronate ester or boronic acid, or trialkylstannane;
- HH metal atom such as tin, zinc, magnesium or lithium as part of an
organometallic compound used as an intermediate for transition
metal mediated aryl-aryl coupling reactions;

JJ -CN, -CHO, or -CO₂R₂₀ wherein R₂₀ is hydrogen or C₁ to C₃ alkyl.

Exemplary conditions for forming and removing suitable nitrogen protecting groups may be found in T. W. Greene and P. G. M. Wuts, 5 Protecting Groups in Organic Synthesis, John Wiley & Sons, Inc, New York, 1991, pp. 309-405. One skilled in the art can also recognize that the heteroaryl sulfonamide-NH in compounds of the invention will also have carboxylic acid character, and accordingly, methods used to protect carboxylic acids may be applicable to protecting the nitrogen NH of the 10 sulfonamides in the invention, including intermediates to compounds of formula I. Exemplary conditions for forming and removing suitable carboxylic acid protecting groups may be found in T. W. Greene and P. G. M. Wuts, Protecting Groups in Organic Synthesis, John Wiley & Sons, Inc, New York, 1991, pp. 175-276.

15

SCHEME I



Compounds of formula I may be prepared from the deprotection of a
 5 compound of formula II wherein BB is a suitable nitrogen protecting
 group. Exemplary conditions for deprotection, and nitrogen protecting
 groups, may be found in T. W. Greene and P. G. M. Wuts, Protecting
Groups in Organic Synthesis, John Wiley & Sons, Inc, New York, 1991,
 pp. 309-405. Preferred nitrogen protecting groups are the methoxymethyl
 10 (MOM), methoxyethoxymethyl (MEM), and 2-(trimethylsilyl)ethoxymethyl
 (SEM) groups.

Compounds of formula II may be prepared from a palladium
 catalyzed coupling of a compound of formula III with a compound of
 formula VIII, in the presence of a suitable base in an inert solvent.
 15 Exemplary palladium catalysts include tetrakis(triphenylphosphine)
 palladium(0), palladium(II) chloride or palladium(II) acetate. The

preferred palladium catalyst is tetrakis(triphenylphosphine) palladium(0). Exemplary bases include tertiary amines, such as, but not limited to, triethylamine, or aqueous potassium, sodium, or cesium carbonate. The preferred base is aqueous sodium carbonate. Exemplary solvents include
5 tetrahydrofuran, 1,4-dioxane, acetonitrile, toluene, benzene, or straight chain alcohols, or a combination thereof. The preferred solvent is a mixture of toluene and ethanol. Exemplary reaction temperatures are between about 25°C to 125°C, preferably between about 65°C and 110°C.

Compounds of formula III may be prepared from a compound of
10 formula IV via displacement of the leaving group (DD) by the conjugate base of a compound R_1-H , wherein R_1 is as previously defined, using a base in an inert solvent. Exemplary bases include sodium carbonate, potassium carbonate, cesium carbonate, sodium hydride, potassium hydride, or alkyl lithiums. The preferred base is sodium hydride.
15 Exemplary inert solvents include ethers (tetrahydrofuran, 1,4-dioxane, diethyl ether), or N,N-dimethylformamide. The preferred solvent is N,N-dimethylformamide. Exemplary reaction temperatures are between about 0°C to 154°C, preferably between about 65°C and 110°C.

Compounds of formula III may also be prepared via a Mitsunobu
20 reaction between a compound of formula VI and the conjugate acid R_1-H , preferably using a phosphine and oxidizing agent, in an inert solvent. Exemplary phosphines include trialkylphosphines, triarylphosphines and polymer supported triarylphosphines. The preferred phosphine is triphenylphosphine. Exemplary oxidizing reagents include diethyl
25 azodicarboxylate, diisopropyl azodicarboxylate, or carbon tetrabromide. The preferred oxidizing reagent is diethyl azodicarboxylate. Exemplary inert solvents include ethers (tetrahydrofuran, 1,4-dioxane, diethyl ether), acetonitrile or N,N-dimethylformamide. The preferred solvent is N,N-dimethylformamide. Exemplary reaction temperatures are between about
30 0°C to 154°C, preferably between about 20°C and 65°C.

Compounds of formula IV (especially, where DD is -OSO₂Ph, -OSO₂PhCH₃, -OSO₂CH₃, -OSO₂CF₃) may be prepared from the reaction of a compound of formula VI with ClSO₂Ph, ClSO₂PhCH₃, ClSO₂CH₃ or (CF₃SO₂)₂O in the presence of a base in an inert solvent.

- 5 Compounds of formula VI may be prepared from reduction of a compound of formula VII using a suitable reducing agent in an inert solvent.

Compounds of formula VII are either commercially available or available by means known to one skilled in the art.


- 10 Compounds of formula VIII may be prepared via lithiation of a compound of formula IX wherein AA is hydrogen or a halogen (chloro, bromo, iodo), and reacting the resulting aryl lithium with an appropriate borate derivative.

- 15 Compounds of formula IX may be prepared via the protection of the nitrogen in a compound of formula XI. Exemplary nitrogen protecting groups and methods of protecting the nitrogen are similar to those for protecting amines, such as those described in T. W. Greene and P. G. M. Wuts, Protecting Groups in Organic Synthesis, John Wiley & Sons, Inc, New York, 1991.

- 20 Compounds of formula XI may be prepared from the reaction of a compound of formula XII with a compound R₃-NH₂.

Compounds of the formula XII are either commercially available or available by means known to one skilled in the art.

SCHEME II



Compounds of formula I may be prepared from the deprotection of a
5 compound of formula II as described in Scheme I.

Compounds of formula II may be prepared from a compound of
formula XIV via displacement of the leaving group (DD) by the conjugate
base of a compound R_1-H , wherein R_1 is as previously defined, using a base
in an inert solvent. Exemplary bases include sodium carbonate,
10 potassium carbonate, cesium carbonate, sodium hydride, potassium
hydride, or alkyl lithiums. The preferred base is sodium hydride.
Exemplary inert solvents include ethers (tetrahydrofuran, 1,4-dioxane,
diethyl ether), or N,N-dimethylformamide. The preferred solvent is N,N-
dimethylformamide. Exemplary reaction temperatures are between about
15 0°C to 154°C , preferably between about 25°C and 110°C .

Compounds of formula II may also be prepared via a Mitsunobu
reaction between a compound of formula XV and the conjugate acid R_1-H
using a phosphine and oxidizing agent in an inert solvent. Exemplary

phosphines include trialkylphosphines, triarylphosphines and polymer supported triarylphosphines. The preferred phosphine is triphenylphosphine. Exemplary oxidizing reagents include diethyl azodicarboxylate, diisopropyl azodicarboxylate, or carbon tetrabromide.

- 5 The preferred oxidizing reagent is diethyl azodicarboxylate. Exemplary inert solvents include ethers (tetrahydrofuran, 1,4-dioxane, diethyl ether), acetonitrile or N,N-dimethylformamide. The preferred solvent is N,N-dimethylformamide. Exemplary reaction temperatures are between about 0°C to 154°C, preferably between about 20°C and 65°C.

- 10 Compounds of formula XIV may be prepared from compounds of formula XV using methods well known in the art. For example, compounds of formula XIV (DD = Br) may be prepared by the treatment of compound XV with carbon tetrabromide and triphenylphosphine in a suitable solvent such as toluene or tetrahydrofuran.


- 15 Compounds of formula XV may be prepared from reduction of a compound of formula XVI using a suitable reducing agent in an inert solvent. R₂ is preferably not an amide, an ester, a carboxylic acid or an aldehyde during this operation.

- 20 Compounds of formula XVI may be prepared from a palladium catalyzed coupling of a compound of formula VII with a compound of formula IX in the presence of a suitable base and an inert solvent as described in Scheme I.

Compounds of formula VII are available by means known to one skilled in the art.

25

SCHEME III



Compounds of formula XVII (which are compounds of formula I

5 wherein R_2 is R_{2a} where R_{2a} is $-\text{CH}_2\text{N}(R_{21})(\text{C}=\text{O})\text{N}(R_{19})R_{20}$, $-\text{CH}_2\text{N}(R_{21})(\text{C}=\text{O})\text{OR}_{18}$ or $-\text{CH}_2\text{N}(R_{21})(\text{C}=\text{O})R_{22}$), may be prepared from compounds of formula XVIII (also compounds of formula I, where R_2 is $-\text{CH}_2\text{NHR}_{21}$) by reaction of a compound of formula XVIII with an active ester (i.e., from a carboxylic acid such as $R_{22}\text{COOH}$ in the presence of a

10 suitable coupling agent such as dicyclohexylcarbodiimide (DCC)), or an acid chloride (i.e., $R_{22}(\text{C}=\text{O})\text{Cl}$), or an isocyanate (i.e., $R_{19}\text{N}=\text{C}=\text{O}$), or a chloroformate (i.e., $R_{18}\text{O}(\text{C}=\text{O})\text{Cl}$) in the presence of a suitable base such as triethylamine and catalyst such as 4-dimethylaminopyridine in an inert solvent. This step may, for example, be conducted combinatorially and a

15 library of such compounds created.

Compounds of formula XVIII may be prepared from reductive amination of compounds of formula XIX (compounds of formula I where R_2 is

-CHO) using a primary amine such as $R_{21}NH_2$ in the presence of a suitable reducing agent such as sodium triacetoxyborohydride in an inert solvent.

Compounds of formula XIX may be prepared by deprotection of compounds of formula XX wherein BB is a suitable nitrogen protecting group as described in Scheme I.

Compounds of formula XX may be prepared from a palladium catalyzed coupling of a compound of formula XXI with a compound of formula VIII in the presence of a suitable base and an inert solvent as described in Scheme I.

Compounds of formula XXI may be prepared in two steps from a compound of formula XXIII, first by displacement of the leaving group (DD) by the conjugate base of a compound R_1-H , wherein R_1 is as previously defined using a suitable base in an inert solvent as described in Scheme I to provide a compound of formula XXII. Subsequent deprotection of the acetal in a compound of formula XXII using methods known in the art is employed to provide a compound of formula XXI.

Compounds of formula XXII may also be prepared via a Mitsunobu reaction between a compound of formula XXIV and the conjugate acid R_1-H using a phosphine and oxidizing agent in an inert solvent.

Exemplary phosphines include trialkylphosphines, triarylphosphines and polymer supported triarylphosphines. Exemplary oxidizing reagents include diethyl azodicarboxylate, diisopropyl azodicarboxylate, or carbon tetrabromide. Exemplary inert solvents include ethers (tetrahydrofuran, 1,4-dioxane, diethyl ether), acetonitrile or N,N-dimethylformamide.


Compounds of formula XXIII may be prepared from compounds of formula XXIV using methods well known in the art. For example, compounds of formula XXIII (DD = Br) may be prepared by the treatment of compound XXIV with carbon tetrabromide and triphenylphosphine in a suitable solvent such as toluene or tetrahydrofuran.

Compounds of formula XXIV may be prepared in two steps from compound XXV via a partial reduction of the nitrile group to the aldehyde

using a suitable reducing agent such as diisobutylaluminum hydride, with subsequent reduction of the aldehyde to hydroxymethyl using an agent such as sodium borohydride.

Methods for the preparation of compounds XXV and XXVI are known in the art [H.-Y. Zhang, et al., *Tetrahedron*, 50, 11339-11362 (1994)].

SCHEME IV




Compounds of formula XXIX (which are compounds of formula I where R₂ is -CH₂Y) may be prepared from the deprotection of a compound of formula XXX such as is described in Scheme I.

Compounds of formula XXX may be prepared from a compound of formula XXXI via displacement of the leaving group (DD) by the conjugate base of a compound Y-H, wherein Y is as previously defined using a base in an inert solvent. Exemplary bases include sodium carbonate, potassium carbonate, cesium carbonate, sodium hydride, potassium hydride, or alkyl lithiums. The preferred base is sodium hydride.


Compounds of formula XXXI may be prepared from compounds of formula XX using methods well known in the art. For example, compounds of formula XXXI (DD = Br) may be prepared from compound XX in two steps: first by reducing the aldehyde to a hydroxymethyl group using a suitable reducing agent such as sodium borohydride, and second, conversion of the hydroxymethyl group to the bromomethyl function using carbon tetrabromide and triphenylphosphine in a suitable solvent such as toluene or tetrahydrofuran.

SCHEME V



5 Compounds of formula XXXII (which may be employed, for example, in the methods of the preceding Schemes) may be prepared by cyclization of compounds of formula XXXIII in the presence of orthoester XXXIV using a catalytic amount of a weak acid such as acetic acid. Exemplary reaction temperatures are between about 25°C to 154°C, preferably
10 between about 60°C and 110°C.

Compounds of formula XXXIII (e.g., where R_{13} and R_{14} , together with the atoms to which they are bonded, form the five-membered ring



) may be prepared from compounds of formula XXXVI in two steps:

- (1) acylation of compound XXXVI with an N-protected amino acid in the presence of a suitable coupling agent such as dicyclohexylcarbodiimide (DCC) or (benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate (PyBOP) in a suitable solvent such as N,N-dimethylformamide, and (2) removal of the protecting group. Suitable conditions and suitable nitrogen protecting groups and the corresponding
15 deprotection conditions may be found in T. W. Greene and P. G. M. Wuts,
20


Protecting Groups in Organic Synthesis, John Wiley & Sons, Inc, New York, 1991, pp. 309-405.

Compounds of formula XXXVI may be prepared via reduction of a compound of formula XXXVII using an appropriate reducing agent such as diborane or lithium aluminum hydride in an appropriate solvent such as tetrahydrofuran.

Compounds of formula XXXVII may be prepared from compounds of formula XXXVIII as described in Scheme III.

Compounds of formula XXXVIII may be prepared by methods known in the art.

SCHEME VI



Compounds of formula I may be prepared from the deprotection of a compound of formula II as described in Scheme I.

Compounds of formula II may be prepared by a palladium catalyzed coupling of a compound of formula IX (as described in Scheme I) wherein AA here is $-\text{OSO}_2\text{CF}_3$ or a halogen (chlorine, bromine, or iodine; preferably bromine or iodine) with a compound of formula XXXIXa, wherein GG is a boronic acid or ester, in the presence of a base and an inert solvent as described in Scheme I.

Compounds of formula II may also be prepared by a palladium or nickel catalyzed coupling of a compound of formula IX (as described in

Scheme I) wherein AA is a halogen (chlorine, bromine, or iodine; preferably bromine or iodine) with a compound of formula XXXIXb wherein HH is a suitable metal atom bearing appropriate ligands. Exemplary metal atoms include tin, zinc, magnesium, and lithium.


- 5 Exemplary catalysts include tetrakis(triphenylphosphine)palladium(0) and dichlorobis(triphenylphosphine)nickel(II).

Compounds of formula XXXIXa or XXXIXb may be prepared via lithiation of a compound of formula III wherein EE is a halogen (chlorine, bromine, or iodine; preferably bromine or iodine), then reacting the
 10 resulting aryl lithium with an appropriate borate derivative or with an appropriate zinc, tin, or magnesium reagent.

Compounds of formula III may be prepared by the methods described in Scheme I.

SCHEME VII

15



Compounds of formula I may be prepared from the thermal reaction of a compound of formula XL with a heterocyclic compound of formula

R_3 -X, wherein X is a halogen (fluorine, chlorine, bromine, or iodine), in the presence of a base and an inert solvent. Exemplary bases include sodium hydride, potassium carbonate, potassium hydride, and potassium bis(trimethylsilyl)amide, preferably sodium hydride. Exemplary solvents
5 include N,N-dimethylformamide and N,N-dimethylacetamide. Exemplary reaction temperatures are between about 80°C and 150°C, preferably between 110°C and 130°C.

Compounds of formula I may also be prepared from the reaction of a compound of formula XL with a heterocyclic compound of formula R_3 -X,
10 wherein X is a halogen (chlorine, bromine, or iodine), in the presence of a palladium catalyst, a phosphine ligand, a base, and an inert solvent. Exemplary palladium catalysts include palladium acetate and tris(dibenzylideneacetone)palladium(0), and the preferred palladium catalyst is palladium acetate. The preferred phosphine ligand is 2,2'-
15 bis(diphenylphosphino)-1,1'-binaphthyl. Exemplary bases include sodium hydride and sodium t-butoxide. The preferred base is sodium hydride. Exemplary reaction temperatures are between about 20°C and 110°C, preferably between 85°C and 110°C.

Compounds of formula XL may be prepared by deprotection of a
20 compound of formula XLI, wherein BB is a suitable nitrogen protecting group. Exemplary conditions for protection and deprotection of nitrogen functionalities may be found in T. W. Greene and P. G. M. Wuts, Protecting Groups in Organic Synthesis, John Wiley & Sons, Inc., New York, 1991, pp. 309-405. The preferred protecting group BB for Scheme
25 VII is *tertiary*-butyl. Exemplary deprotection conditions include the use of acids, such as trifluoroacetic acid.


Compounds of formula XLI may be prepared by a palladium catalyzed coupling of a compound of formula III (as described in Scheme I) wherein EE is a halogen (chlorine, bromine, or iodine; preferably
30 bromine or iodine) with a compound of formula XLII, wherein GG is a

boronic acid or ester, in the presence of a base and an inert solvent as described in Scheme I.

Compounds of formula XLII may be prepared via the lithiation of a compound of formula XLIII in an inert solvent, followed by reacting the resulting aryl lithium with an appropriate borate derivative. Exemplary reagents for the lithiation reaction include *n*-butyllithium and *t*-butyllithium. Exemplary solvents include ethers such as tetrahydrofuran, either alone or in combination with hydrocarbon solvents such as hexane. The preferred solvent is a mixture of tetrahydrofuran and hexane.

Compounds of formula XLIII are either commercially available or available by means known to one skilled in the art.

SCHEME VIII



Compounds of formula XXIX (which are certain compounds of formula I where R₂ is -CH₂Y, as described in Scheme IV) may be prepared from a compound of formula XIX (which is a compound of formula I where R₂ is -CHO, as described in Scheme III) via a two step process: 1) reductive amination of XIX in the presence of a primary amine R₃₀NH₂,


wherein R_{30} is carboxyalkyl, alkoxycarbonylalkyl, hydroxyalkyl, or aminoalkyl, using a suitable reducing agent such as sodium triacetoxyborohydride, yields an intermediate amine; 2) subsequent cyclization using an appropriate cyclization reagent yields a compound of formula XXIX. When R_{30} is carboxyalkyl, appropriate cyclization reagents include carbodiimides such as diisopropylcarbodiimide. When R_{30} is hydroxyalkyl or aminoalkyl, appropriate cyclization reagents include phosgene and 1,1'-carbonyldiimidazole. When R_{30} is alkoxycarbonylalkyl, appropriate cyclization reagents include tertiary amine bases such as triethylamine and N,N-diisopropylethylamine.

Compounds of formula XIX may be prepared via deprotection of a compound of formula XX as described in Scheme III.

Compounds of formula XXIX may also be prepared by deprotection of a compound of formula XXX as described in Scheme I.

Compounds of formula XXX may be prepared from a compound of formula XX, using the two step process described above for the formation of compounds of formula XXIX from a compound of formula XIX.

SCHEME IX



5 Compounds of formula XLIV (which are compounds of formula I wherein R_1 is D as defined for formula I) may be prepared from the deprotection of a compound of formula XLV as described in Scheme I.

10 Compounds of formula XLV may be prepared via the acylation of a compound of formula XLVI using either a carboxylic acid such as R_6COOH in the presence of a suitable coupling agent such as dicyclohexylcarbodiimide, or the corresponding acid chloride or acid anhydride in the presence of a suitable base such as triethylamine.


15 Compounds of formula XLVI may be prepared from reduction of a compound of formula XLVII in the presence of a primary amine such as $H_2NCHR_7R_9$ in the presence of a suitable reducing agent such as sodium triacetoxyborohydride.

Compounds of formula XLVII may be prepared via reduction of a compound of formula XVI, as described in Scheme II, wherein JJ is -CN, or -CO₂R₂₀ wherein R₂₀ is hydrogen or C₁ to C₃ alkyl, using means known to one skilled in the art.

- 5 Compounds of formula XLVII may also be prepared via oxidation of a compound of formula XV, as defined in Scheme II, using means known to one skilled in the art.

SCHEME X

10




- Compounds of formula XLIV (which are compounds of formula I wherein R₁ is D as defined for formula I) may be prepared by the acylation of a compound of formula LXI using either a carboxylic acid such as R₆COOH in the presence of a suitable coupling agent such as dicyclohexylcarbodiimide, or the corresponding acid chloride or acid anhydride in the presence of a suitable base such as triethylamine.
- 15

- Compounds of formula LXI may be prepared by the reduction of a compound of formula LXII in the presence of a primary amine such as
- 20

$H_2NCHR_7R_9$ in the presence of a suitable reducing agent such as sodium triacetoxyborohydride.

Compounds of formula LXII may be prepared via deprotection of a compound of formula LXIII (which is a compound of formula XVI wherein
5 JJ is CHO), as described in Scheme I.

SCHEME XI



10 Compounds of formula LXIV (which are compounds of formula I wherein R_2 is $-N(R_{19})R_{20}$) may be prepared via reduction of a compound of formula LXV in the presence of an aliphatic, aromatic, or heteroaromatic aldehyde using a suitable reducing agent such as sodium triacetoxyborohydride.

15 Compounds of formula LXV (which are compounds of formula I wherein R_2 is $-NHR_{19}$) may be similarly prepared via reduction of a compound of formula LXVI in the presence of an aliphatic, aromatic, or heteroaromatic aldehyde using a suitable reducing agent such as sodium triacetoxyborohydride.


20 Compounds of formula LXVI (which are compounds of formula I wherein R_2 is $-NH_2$) may be prepared by reduction of a compound of

formula LXVII using a suitable reducing agent such as tin (II) chloride dihydrate in a suitable solvent such as ethyl acetate.

Compounds of formula LXVII (which are compounds of formula I wherein R_2 is $-\text{NO}_2$) may be prepared by deprotection of a compound of formula LXVIII as described in Scheme I.

Compounds of formula LXVIII (which are compounds of formula II wherein R_2 is $-\text{NO}_2$) may be prepared by the methods described for the preparation of compounds of formula II in Schemes I and II.

SCHEME XII




Compounds of formula LXIX (which are compounds of formula I wherein R_2 is $-\text{N}(\text{R}_{21})(\text{C}=\text{O})\text{R}_{22}$) may be prepared via acylation of a compound of formula LXVa (prepared as described in Scheme XI for a compound of formula LXV) using either a carboxylic acid such as R_{22}COOH in the presence of a suitable coupling agent such as dicyclohexylcarbodiimide, or the corresponding acid chloride or acid anhydride in the presence of a suitable base such as triethylamine.

Compounds of formula LXX (which are compounds of formula I wherein R_2 is $-\text{N}(\text{R}_{21})\text{SO}_2\text{R}_{22}$) may be prepared via sulfonylation of a compound of formula LXVa using either a sulfonyl chloride such as


$R_{22}SO_2Cl$ or the corresponding sulfonic anhydride, in the presence of a suitable base such as triethylamine.

5


SCHEME XIII

Compounds of formula LXXI (which are compounds of formula I wherein R_2 is $-CH_2N(R_{21})(SO_2R_{22})$) may be prepared via sulfonylation of a compound of formula XVIII (prepared as described in Scheme III) using either a sulfonyl chloride such as $R_{22}SO_2Cl$ or the corresponding sulfonic anhydride in the presence of a suitable base such as triethylamine.


The present invention further provides the following novel compounds, which may be employed as intermediates in the preparation of compounds of the formula I and salts thereof:



wherein R_2 , R_3 , R_{101} , R_{102} , R_{103} , and R_{104} are as defined for a compound of formula I, R_{50} is hydroxy, chloro, bromo, iodo, $-OSO_2$ -alkyl, or $-OSO_2$ -aryl, and R_{51} is hydrogen, $-CH_2OCH_2CH_2OCH_3$, $-CH_2OCH_2CH_2Si(CH_3)_3$, $-CH_2OCH_3$, $-CH_2OCH_2$ -aryl, or other suitable nitrogen protecting group;




- wherein R_1 , R_2 , R_3 , R_{101} , R_{102} , R_{103} , and R_{104} are as defined for a compound of
 5 formula I and BB is $-\text{CH}_2\text{OCH}_2\text{CH}_2\text{OCH}_3$, $-\text{CH}_2\text{OCH}_2\text{CH}_2\text{Si}(\text{CH}_3)_3$, $-\text{CH}_2\text{OCH}_3$, $-\text{CH}_2\text{OCH}_2$ -aryl, or other suitable nitrogen protecting group; and



- 10 wherein R_1 , R_2 , R_{101} , and R_{102} are as defined for a compound of formula I, and R_{52} is chloro, bromo, iodo, or $-\text{OSO}_2\text{CF}_3$.


- The present invention also provides the following novel method for the preparation of a compound of the formula I or salt thereof, wherein
 15 said method comprises at least one of the following steps:

- a) displacement of a leaving group R_{50} via the anion of a compound R_1 -H from a compound of the formula




wherein R_1 , R_2 , R_3 , R_{101} , R_{102} , R_{103} , and R_{104} are as defined for a compound of formula I, R_{50} is hydroxy, chloro, bromo, iodo, $-\text{OSO}_2$ -alkyl or $-\text{OSO}_2$ -aryl, and R_{51} is hydrogen or a suitable nitrogen protecting group, using a Mitsunobu reaction or S_N1 or S_N2 displacement reaction, with removal of
 5 said nitrogen protecting group as appropriate;

b) removal of the nitrogen protecting group R_{51} from a compound of formula




10 wherein R_1 , R_2 , R_3 , R_{101} , R_{102} , R_{103} , and R_{104} are as defined for a compound of formula I, and R_{51} is a suitable nitrogen protecting group;

c) organometallic coupling of a compound of formula




15 with a compound of formula



wherein R_1 , R_2 , R_3 , R_{101} , R_{102} , R_{103} , and R_{104} are as defined for a compound of formula I and R_{51} is a suitable nitrogen protecting group. R_{52} is chloro, bromo, iodo or $-\text{OSO}_2\text{CF}_3$ and R_{54} is a boronic acid, boronic ester or stannane derivative. Or R_{52} is a boronic acid, boronic ester or stannane derivative, and R_{54} is chloro, bromo, iodo or $-\text{OSO}_2\text{CF}_3$;
 20

d) acylation of a compound of the formula




wherein R_1 , R_3 , R_{101} , R_{102} , R_{103} , and R_{104} are as defined for a compound of formula I, and R_{51} is hydrogen or a suitable nitrogen protecting group,

5 with an acylating agent of the formula $R_{55}-(C=O)R_{22}$, $R_{19}N=C=O$, $R_{55}-CO_2R_{18}$, $R_{55}SO_2R_{22}$, wherein R_{18} , R_{19} and R_{22} are as defined for a compound of formula I and R_{55} is an activating group for an acid, or made using an acid activating agent, with removal of said nitrogen protecting group as appropriate; or


10

e) reductive amination of a compound of formula



wherein R_1 , R_3 , R_{101} , R_{102} , R_{103} , and R_{104} are as defined for a compound of formula I, and R_{51} is hydrogen or a suitable nitrogen protecting group,

15 with an amine of the formula



wherein R_{23} , R_{24} , and x are as defined for a compound of formula I, with removal of said nitrogen protecting group as appropriate.

5

Utility

The compounds of formula I and salts thereof are antagonists of both endothelin (especially, ET-1) and angiotensin II (especially, subtype AT_1) receptors ("dual angiotensin endothelin receptor antagonists") and are useful in treatment of conditions associated with increased ET levels and/or increased angiotensin II levels and of all endothelin-dependent or angiotensin II-dependent disorders. They are thus useful as antihypertensive agents. By the administration of a composition having one (or a combination) of the compounds of this invention, the blood pressure of a hypertensive mammalian (e.g., human) host is reduced.

10 They are also useful in portal hypertension, hypertension secondary to treatment with erythropoietin and low renin hypertension.

The compounds of the present invention are also useful in the treatment of disorders related to renal, glomerular and mesangial cell function, including acute (such as ischemic, nephrotoxic, or glomerulonephritis) and chronic (such as diabetic, hypertensive or immune-mediated) renal failure, diabetic nephropathy, glomerular injury, renal damage secondary to old age or related to dialysis, nephrosclerosis (especially hypertensive nephrosclerosis), nephrotoxicity (including nephrotoxicity related to imaging and contrast agents and to cyclosporine),

20 renal ischemia, primary vesicoureteral reflux, glomerulosclerosis and the like. The compounds of this invention are also useful in the treatment of disorders related to paracrine and endocrine function. The compounds of this invention are also useful in the treatment of diabetic nephropathy, hypertension-induced nephropathy, and IGA-induced nephropathy.

30 The compounds of the present invention are also useful in the treatment of endotoxemia or endotoxin shock as well as hemorrhagic

shock. The compounds of the present invention are also useful in alleviation of pain associated cancer, such as the pain associated with prostate cancer, and bone pain associated with bone cancer. The compounds of the present invention are further useful in the prevention
5 and/or reduction of end-organ damage associated the cell-poliferative effects of endothelin.

The compounds of the present invention are also useful in hypoxic and ischemic disease and as anti-ischemic agents for the treatment of, for example, cardiac, renal and cerebral ischemia and reperfusion (such as
10 that occurring following cardiopulmonary bypass surgery), coronary and cerebral vasospasm, and the like.

In addition, the compounds of this invention are also useful as anti-arrhythmic agents; anti-anginal agents; anti-fibrillatory agents; anti-asthmatic agents; anti-atherosclerotic and anti-arteriosclerotic agents
15 (including anti-transplantation arteriosclerotic agents); additives to cardioplegic solutions for cardiopulmonary bypasses; adjuncts to thrombolytic therapy; and anti-diarrheal agents. The compounds of this invention may be useful in therapy for myocardial infarction; therapy for peripheral vascular disease (e.g., Raynaud's disease, intermittent
20 claudication and Takayashu's disease); treatment of cardiac hypertrophy (e.g., hypertrophic cardiomyopathy); treatment of primary pulmonary hypertension (e.g., plexogenic, embolic) in adults and in the newborn and pulmonary hypertension secondary to heart failure, radiation and chemotherapeutic injury, or other trauma; treatment of central nervous
25 system vascular disorders, such as stroke, migraine and subarachnoid hemorrhage; treatment of central nervous system behavioral disorders; treatment of gastrointestinal diseases such as ulcerative colitis, Crohn's disease, gastric mucosal damage, ulcer, inflammatory bowel disease and ischemic bowel disease; treatment of gall bladder or bile duct-based
30 diseases such as cholangitis; treatment of pancreatitis; regulation of cell growth; treatment of benign prostatic hypertrophy; restenosis following

angioplasty or following any procedure including transplantation and stenting; therapy for congestive heart failure including inhibition of fibrosis; inhibition of left ventricular dilatation, remodeling and dysfunction; and treatment of hepatotoxicity and sudden death. The compounds of this invention are useful in the treatment of sickle cell disease including the initiation and/or evolution of the pain crises of this disease; treatment of the deleterious consequences of ET-producing tumors such as hypertension resulting from hemangiopericytoma; treatment of early and advanced liver disease and injury including attendant complications (e.g., hepatotoxicity, fibrosis and cirrhosis); treatment of spastic diseases of the urinary tract and/or bladder; treatment of hepatorenal syndrome; treatment of immunological diseases involving vasculitis such as lupus, systemic sclerosis, mixed cryoglobulinemia; and treatment of fibrosis associated with renal dysfunction and hepatotoxicity. The compounds of this invention are useful in therapy for metabolic and neurological disorders; cancer; insulin-dependent and non insulin-dependent diabetes mellitus; neuropathy; retinopathy; epilepsy; hemorrhagic and ischemic stroke; bone remodeling; psoriasis; and chronic inflammatory diseases such as arthritis, rheumatoid arthritis, osteoarthritis, sarcoidosis and eczematous dermatitis (all types of dermatitis).

The compounds of this invention are additionally useful in the treatment of disorders involving bronchoconstriction and disorders of chronic or acute pulmonary inflammation such as chronic obstructive pulmonary disease (COPD) and adult respiratory distress syndrome (ARDS).

The compounds of this invention are also useful in the treatment of sexual dysfunction in both men (erectile dysfunction, for example, due to diabetes mellitus, spinal cord injury, radical prostatectomy, psychogenic etiology or any other cause) and women by improving blood flow to the genitalia, especially, the corpus cavernosum.

The compounds of this invention are also useful in the treatment of dementia, including Alzheimer's dementia, senile dementia and vascular dementia.

5 Additionally the compounds of the present invention are further useful in the reduction of general morbidity and/or mortality as a result of the above utilities.

The present invention thus provides methods for the treatment of all endothelin-dependent or angiotensin II-dependent disorders, comprising the step of administering to a subject in need thereof at least
10 one compound of the formula I in an amount effective therefor. Other therapeutic agents such as those described below may be employed with the inventive compounds in the present methods. In the methods of the present invention, such other therapeutic agent(s) may be administered prior to, simultaneously with or following the administration of the
15 compound(s) of the present invention.

The effective amount of a compound of the present invention may be determined by one of ordinary skill in the art, and includes exemplary dosage amounts for a human of from about 0.1 to about 100 mg/kg, preferably about 0.2 to about 50 mg/kg and more preferably from about 0.5
20 to about 25 mg/kg of body weight (or from about 1 to about 2500 mg, preferably from about 5 to about 500 mg) of active compound per day, which may be administered in a single dose or in the form of individual divided doses, such as from 1 to 4 times per day. It will be understood that the specific dose level and frequency of dosage for any particular
25 subject may be varied and will depend upon a variety of factors including the activity of the specific compound employed, the metabolic stability and length of action of that compound, the species, age, body weight, general health, sex and diet of the subject, the mode and time of administration, rate of excretion, drug combination, and severity of the particular
30 condition. Preferred subjects for treatment include animals, most preferably mammalian species such as humans, and domestic animals

such as dogs, cats and the like, subject to endothelin-dependent or angiotensin II-dependent disorders.

The present invention also provides pharmaceutical compositions comprising at least one of the compounds of the formula I capable of
5 treating an endothelin-dependent or angiotensin II-dependent disorder in an amount effective therefor, and a pharmaceutically acceptable vehicle or diluent. The compositions of the present invention may contain other therapeutic agents as described below, and may be formulated, for example, by employing conventional solid or liquid vehicles or diluents, as
10 well as pharmaceutical additives of a type appropriate to the mode of desired administration (for example, excipients, binders, preservatives, stabilizers, flavors, etc.) according to techniques such as those well known in the art of pharmaceutical formulation or called for by accepted pharmaceutical practice.

15 The compounds of the formula I may be administered by any suitable means, for example, orally, such as in the form of tablets, capsules, granules or powders; sublingually; buccally; parenterally, such as by subcutaneous, intravenous, intramuscular, or intrasternal injection or infusion techniques (e.g., as sterile injectable aqueous or non-aqueous
20 solutions or suspensions); nasally such as by inhalation spray; topically, such as in the form of a cream or ointment; or rectally such as in the form of suppositories; in dosage unit formulations containing non-toxic, pharmaceutically acceptable vehicles or diluents. The present compounds may, for example, be administered in a form suitable for immediate
25 release or extended release. Immediate release or extended release may be achieved by the use of suitable pharmaceutical compositions comprising the present compounds, or, particularly in the case of extended release, by the use of devices such as subcutaneous implants or osmotic pumps. The present compounds may also be administered liposomally. For example,
30 the active substance can be utilized in a composition such as tablet, capsule, solution or suspension containing about 5 to about 500 mg per

unit dosage of a compound or mixture of compounds of formula I or in topical form for wound healing (0.01 to 5% by weight compound of formula I, 1 to 5 treatments per day). They may be compounded in a conventional manner with a physiologically acceptable vehicle or carrier, excipient, binder, preservative, stabilizer, flavor, etc., or with a topical carrier. The compounds of formula I can also be formulated in compositions such as sterile solutions or suspensions for parenteral administration. About 0.1 to 500 milligrams of a compound of formula I may be compounded with a physiologically acceptable vehicle, carrier, excipient, binder, preservative, stabilizer, etc., in a unit dosage form as called for by accepted pharmaceutical practice. The amount of active substance in these compositions or preparations is preferably such that a suitable dosage in the range indicated is obtained.

Exemplary compositions for oral administration include suspensions which may contain, for example, microcrystalline cellulose for imparting bulk, alginic acid or sodium alginate as a suspending agent, methylcellulose as a viscosity enhancer, and sweeteners or flavoring agents such as those known in the art; and immediate release tablets which may contain, for example, microcrystalline cellulose, dicalcium phosphate, starch, magnesium stearate and/or lactose and/or other excipients, binders, extenders, disintegrants, diluents and lubricants such as those known in the art. Molded tablets, compressed tablets or freeze-dried tablets are exemplary forms which may be used. Exemplary compositions include those formulating the present compound(s) with fast dissolving diluents such as mannitol, lactose, sucrose and/or cyclodextrins. Also included in such formulations may be high molecular weight excipients such as celluloses (avicel) or polyethylene glycols (PEG). Such formulations may also include an excipient to aid mucosal adhesion such as hydroxy propyl cellulose (HPC), hydroxy propyl methyl cellulose (HPMC), sodium carboxy methyl cellulose (SCMC), maleic anhydride copolymer (e.g., Gantrez), and agents to control release such as polyacrylic

copolymer (e.g., Carbopol 934). Lubricants, glidants, flavors, coloring agents and stabilizers may also be added for ease of fabrication and use.

Exemplary compositions for nasal aerosol or inhalation administration include solutions in saline which may contain, for example,
5 benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, and/or other solubilizing or dispersing agents such as those known in the art.

Exemplary compositions for parenteral administration include injectable solutions or suspensions which may contain, for example,
10 suitable non-toxic, parenterally acceptable diluents or solvents, such as mannitol, 1,3-butanediol, water, Ringer's solution, an isotonic sodium chloride solution, or other suitable dispersing or wetting and suspending agents, including synthetic mono- or diglycerides, and fatty acids, including oleic acid.

15 Exemplary compositions for rectal administration include suppositories which may contain, for example, a suitable non-irritating excipient, such as cocoa butter, synthetic glyceride esters or polyethylene glycols, which are solid at ordinary temperatures, but liquify and/or dissolve in the rectal cavity to release the drug.

20 Exemplary compositions for topical administration include a topical carrier such as Plastibase (mineral oil gelled with polyethylene). For example, the compounds of the invention may be administered topically to treat peripheral vascular diseases and as such may be formulated as a cream or ointment.

25 The compounds of the present invention may be employed alone or in combination with each other and/or other suitable therapeutic agents useful in the treatment of endothelin-dependent or angiotensin II-dependent disorders. For example, the compounds of this invention can be formulated in combination with endothelin converting enzyme
30 (ECE) inhibitors, such as phosphoramidon; thromboxane receptor antagonists such as ifetroban; potassium channel openers; thrombin

inhibitors (e.g., hirudin and the like); growth factor inhibitors such as modulators of PDGF activity; platelet activating factor (PAF) antagonists; anti-platelet agents such as GPIIb/IIIa blockers (e.g., abciximab, eptifibatide, and tirofiban), P2Y(AC) antagonists (e.g.,

5 clopidogrel, ticlopidine and CS-747), and aspirin; anticoagulants such as warfarin, low molecular weight heparins such as enoxaparin, Factor VIIa inhibitors, and Factor Xa inhibitors such as those described in U.S. Serial No. 09/496,571 filed February 2, 2000 (attorney docket HA 723); renin inhibitors; angiotensin converting

10 enzyme (ACE) inhibitors such as captopril, zofenopril, fosinopril, ceranapril, alacepril, enalapril, delapril, pentopril, quinapril, ramipril, lisinopril and salts of such compounds; neutral endopeptidase (NEP) inhibitors; vasopepsidase inhibitors (dual NEP-ACE inhibitors) such as omapatrilat and gemopatrilat; HMG CoA reductase inhibitors such

15 as pravastatin, lovastatin, atorvastatin, simvastatin, NK-104 (a.k.a. itavastatin, or nisvastatin or nisbastatin) and ZD-4522 (a.k.a. rosuvastatin, or atavastatin or visastatin); squalene synthetase inhibitors; fibrates; bile acid sequestrants such as questran; niacin; anti-atherosclerotic agents such as ACAT inhibitors; MTP inhibitors

20 such as those described in U.S. Serial No. 09/007,938 filed January 16, 1998 (attorney docket HX 91); calcium channel blockers such as amlodipine besylate; potassium channel activators; alpha-adrenergic agents, beta-adrenergic agents such as carvedilol and metoprolol; antiarrhythmic agents; diuretics, such as chlorothiazide,

25 hydrochlorothiazide, flumethiazide, hydroflumethiazide, bendroflumethiazide, methylchlorothiazide, trichloromethiazide, polythiazide or benzothiazide as well as ethacrynic acid, tricrynafen, chlorthalidone, furosemide, musolimine, bumetanide, triamterene, amiloride and spironolactone and salts of such compounds;

30 thrombolytic agents such as tissue plasminogen activator (tPA),

recombinant tPA, streptokinase, urokinase, prourokinase and anisoylated plasminogen streptokinase activator complex (APSAC); anti-diabetic agents such as biguanides (e.g. metformin), glucosidase inhibitors (e.g., acarbose), insulins, meglitinides (e.g., repaglinide),
5 sulfonylureas (e.g., glimepiride, glyburide, and glipizide), biguanide/glyburide combinations such as those described in U.S. Serial No. 09/432,465 filed November 3, 1999 (attorney docket LA 46) and U.S. Serial No. 09/460,920 filed December 14, 1999 (attorney docket LA 46a); thiozolidinediones (e.g. troglitazone, rosiglitazone and
10 pioglitazone), and PPAR-gamma agonists; mineralocorticoid receptor antagonists such as spironolactone and eplerenone; growth hormone secretagogues such as those described in U.S. Serial No. 09/417,180 filed October 12, 1999 (attorney docket LA 25) and U.S. Serial No. 09/506,749 filed February 18, 2000 (attorney docket LA 26); aP2
15 inhibitors such as those described in U.S. Serial No. 09/391,053 filed September 7, 1999 (attorney docket LA 24a) and U.S. Serial No. 09/390,275 filed September 7, 1999 (attorney docket LA 24b); digitalis; ouabian; non-steroidal antiinflammatory drugs (NSAIDS) such as aspirin and ibuprofen; phosphodiesterase inhibitors such as
20 PDE III inhibitors (e.g., cilostazol) and PDE V inhibitors (e.g., sildenafil); protein tyrosine kinase inhibitors; antiinflammatories; antiproliferatives such as methotrexate, FK506 (tacrolimus, Prograf), mycophenolate and mofetil; chemotherapeutic agents; immunosuppressants; anticancer agents and cytotoxic agents (e.g.,
25 alkylating agents, such as nitrogen mustards, alkyl sulfonates, nitrosoureas, ethylenimines, and triazenes); antimetabolites such as folate antagonists, purine analogues, and pyrimidine analogues; antibiotics, such as anthracyclines, bleomycins, mitomycin, dactinomycin, and plicamycin; enzymes, such as L-asparaginase;
30 farnesyl-protein transferase inhibitors; hormonal agents, such as

glucocorticoids (e.g., cortisone), estrogens/antiestrogens, androgens/antiandrogens, progestins, and luteinizing hormone-releasing hormone antagonists, octreotide acetate; microtubule-disruptor agents, such as ecteinascidins or their analogs and
5 derivatives; microtubule-stabilizing agents such as paclitaxel (Taxol®), docetaxel (Taxotere®), and epothilones A-F or their analogs or derivatives; plant-derived products, such as vinca alkaloids, epipodophyllotoxins, taxanes; and topoisomerase inhibitors; prenyl-protein transferase inhibitors; and miscellaneous agents such as,
10 hydroxyurea, procarbazine, mitotane, hexamethylmelamine, platinum coordination complexes such as cisplatin and carboplatin); cyclosporins; steroids such as prednisone or dexamethasone; gold compounds; cytotoxic drugs such as azathioprine and cyclophosphamide; TNF-alpha inhibitors such as tenidap;
15 anti-TNF antibodies or soluble TNF receptor such as etanercept (Enbrel) rapamycin (sirolimus or Rapamune), leflunimide (Arava); and cyclooxygenase-2 (COX-2) inhibitors such as celecoxib (Celebrex) and rofecoxib (Vioxx).

If formulated as a fixed dose, such combination products employ the
20 compounds of this invention within the dosage range described below and the other pharmaceutically active agent within its approved dosage range. The compounds of this invention may also be formulated with, or useful in conjunction with, antifungal and immunosuppressive agents such as amphotericin B, cyclosporins and the like to counteract the glomerular
25 contraction and nephrotoxicity secondary to such compounds. The compounds of this invention may also be used in conjunction with hemodialysis.

The above other therapeutic agents, when employed in combination with the compounds of the present invention, may be used,
30 for example, in those amounts indicated in the Physicians' Desk Reference (PDR) or as otherwise determined by one of ordinary skill in the art.

The following assays may be employed in ascertaining the degree of activity of a compound ("drug") as an endothelin and angiotensin II receptor antagonist. Compounds described in the following Examples
5 have been tested in these assays, and have shown activity.

ET_{A/B} Attached cell binding assay

CHO-K1 cells expressing either the human endothelin A or endothelin B receptor were cultured in Ham's F12 media (Gibco/BRL, Grand Island, NY) with 10% fetal bovine serum (Hyclone), supplemented
10 with 300 µg/mL Geneticin (G-418 Gibco BRL Products, Grand Island, NY) and maintained at 37°C with 5% CO₂ in a humidified incubator. Twenty four hours prior to assay, the cells were treated with 0.25% trypsin-EDTA and were seeded in Falcon, 96 well tissue culture plates at a density of 1.8
15 x 10⁴ cells/ well (the monolayer should reach 80-90% confluency by the day of assay).

In the attached cell assay, culture media was aspirated from each well and the monolayers were washed with 50 µl of PBS (Mg⁺⁺, Ca⁺⁺ free). The binding assay was performed in a total volume of 125 µl consisting of
20 assay buffer (50 mM Tris, pH 7.4, including 1% BSA, and 2 µM phosphoramidon), and 25 µl of either 500 nM ET-1 (to define nonspecific binding) or competing drug. The reaction was initiated with the addition of 25 µl of 0.25 nM [¹²⁵I]-ET-1 (New England Nuclear). Incubation was carried out with gentle orbital shaking, at 4°C, reaching equilibrium at 4
25 hours. The reaction was terminated by aspiration of the reaction buffer and two subsequent washes with cold PBS (Mg⁺⁺, Ca⁺⁺ free). The cells were dissociated by the addition of 100 µl of 0.5N NaOH followed by incubation for 40 minutes. Samples were then transferred from the 96 well format into tubes for counting in a Cobra gamma counter (Packard). Data was
30 analyzed with curve fitting software by Sigma plot.

RASMC binding assay

Assays were conducted in a total volume of 250 μ L in 96 well microtitre plates. The incubation mixture contained 50 μ L [125]I-Sar-Ile-Angiotensin II (0.2 nM), 25 μ L of drug dissolved in DMSO, or angiotensin II (1 μ M) to define non-specific binding. Binding to rat aortic smooth muscle cells (RASMCs) was conducted in RPMI media (Gibco BRL Products, Grand Island, NY) containing 0.1% BSA for 2 hours at room temperature with continuous shaking. Unbound radioligand was washed from the wells. The RASMCs with bound radioligand are lysed with 1% Triton X and 0.1% BSA in distilled water for 15 minutes at room temperature with continuous shaking. The solution in each well was transferred to tubes and placed in a gamma counter.

Compounds within the scope of this invention include compounds that have an IC_{50} concentration of less than 100 micromolar versus either or both [125]I-Sar-Ile-Angiotensin II or [125]I-ET-1, ideally against both ligands. Preferred compounds within the scope of this invention are compounds that have an IC_{50} concentration of less than 5 micromolar versus either or both [125]I-Sar-Ile-Angiotensin II or [125]I-ET-1, ideally against both ligands. More preferred compounds within the scope of this invention are compounds that have an IC_{50} concentration of less than 1 micromolar versus either or both [125]I-Sar-Ile-Angiotensin II or [125]I-ET-1, ideally against both ligands.

All documents cited in the present specification are incorporated herein by reference in their entirety.

The following Examples illustrate embodiments of the present invention, and are not intended to limit the scope of the claims. Abbreviations employed herein are defined below. Compounds of the Examples are identified by the example and step in which they are prepared (for example, "1A" denotes the title compound of step A of Example 1), or by the example only where the compound is the title

compound of the example (for example, "4" denotes the title compound of Example 4). Compounds prepared for use as synthetic intermediates are identified by the Preparation number and step in which they appear, prefaced by the letter "P." For example, "P1A" denotes the compound
5 generated in step A of Preparation 1, while "P1" denotes the title compound of Preparation 1.

Abbreviations

- 10 Ac = acetyl
(S)-BINAP = (S)-(-)-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl
BOC = t-butoxycarbonyl
n-Bu = *n*-butyl
BSA = bovine serum albumin
15 CDI = 1,1' carbonyldiimidazole
d = days
DBU = 1,8-diazabicyclo[5.4.0]undec-7-ene
DIBAL-H = diisobutylaluminum hydride
DMF = N,N-dimethylformamide
20 DMSO = dimethylsulfoxide
EDCI = 1-ethyl-3-[3-(dimethylamino)propyl]carbodiimide hydrochloride
EDTA = ethylenediaminetetraacetic acid
eq = equivalents
Et = ethyl
25 ET = endothelin
ET-1 = endothelin-1
EtOAc = ethyl acetate
EtOH = ethanol
h = hours
30 Me = methyl
MEM = methoxyethoxymethyl

MeOH = methanol

min = minutes

mp = melting point

Ms = methanesulfonyl

5 NBS = N-bromosuccinimide

PBS = phosphate buffered saline

Ph = phenyl

n-Pr = *n*-propyl

SEM = 2-(trimethylsiloxy)ethoxymethyl

10 Rochelle's salt = potassium sodium tartrate tetrahydrate

RT = room temperature

TFA = trifluoroacetic acid

THF = tetrahydrofuran

General Methods

15

The following General Methods were employed in the Preparations and Examples.

General Method 1: Suzuki Coupling of Aryl

20

Bromides with Arylboronic Acids



R = H, alkyl

25 A solution of 1.0 eq of an arylboronic acid (or ester) and the appropriate aryl bromide (1.0 eq) in 2:1 toluene:ethanol (0.1 M concentration for each reagent) was sparged with nitrogen for 15 minutes. Tetrakis (triphenylphosphine)palladium (0) (0.05 eq) and 2 M aqueous sodium carbonate (3 eq) were added and the mixture was heated at 85°C for 3 h under a nitrogen atmosphere. The mixture was cooled and ethyl
30 acetate and water were added. The organic layer was washed once with

saturated aqueous sodium carbonate, dried over sodium sulfate, and concentrated. The residue was chromatographed on silica gel using hexanes/ethyl acetate as eluant to yield the biaryl product.


- 5 Arylboronic acids used: [2-[[[(3,4-dimethyl-5-isoxazolyl)](2-methoxyethoxy)-methyl]amino]-sulfonyl]phenyl]boronic acid (or the corresponding SEM-protected compound, both of which were prepared as described in U.S. Patent No. 5,612,359); [2-[[[(4,5-dimethyl-3-isoxazolyl)](2-methoxyethoxy)methyl]amino]-sulfonyl]phenyl]boronic acid (prepared as
 10 described in U.S. Patent No. 5,612,359 and U.S. Patent Application Serial No. 09/013,952, filed January 27, 1998); [2-(*N*-*tert*-butylsulfamoyl)phenyl]boronic acid (prepared according to Chang, L. L. et al., *J. Med. Chem.*, 38, 3741-3758 (1995)).
- 15 Arylboronate ester used: N-[(2-methoxyethoxy)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzenesulfonamide (prepared as described in WO 97/29747).

20 General Method 2: Conversion of Primary Alcohols to Alkyl Bromides




- 25 To a 0.2 M solution of the alcohol in DMF at 0 °C was added carbon tetrabromide (1.5 eq) followed by triphenylphosphine (1.5 eq). The mixture was stirred at 0 °C for 4 h, diluted with 10 parts 2:1 hexanes/ethyl acetate, and washed with water and brine. The solution was dried over sodium sulfate and concentrated, and the residue chromatographed on
 30 silica gel using hexanes/ethyl acetate as eluant to yield the alkyl bromide product.

General Method 3: Conversion of Primary
Alcohols to Alkyl Methanesulfonates



To a 0.15 M solution of the alcohol in dichloromethane at 0 °C was added *N,N*-diisopropylethylamine (1.5 eq) followed by methanesulfonyl chloride (1.1 eq). The mixture was stirred at 0 °C for 1 to 10 3 h, and was then treated with 10% aqueous potassium dihydrogensulfate. The aqueous layer was extracted once with dichloromethane and the combined organic layers were dried over sodium sulfate and concentrated to yield the crude alkyl methanesulfonate.














15 General Method 4: Alkylation of Heterocycles or Aliphatic Alcohols




Sodium hydride (60% dispersion in mineral oil, 1.2 eq) was added at 20 0 °C to a 1.0 M solution or suspension of an appropriate heterocycle or alcohol (1.5 eq) in DMF. The mixture was allowed to warm to RT, was stirred for 20 min, and was then cooled back to 0 °C. To the heterocycle mixture was added a solution of the appropriate alkyl bromide or alkyl methanesulfonate (1.0 eq) in a minimal amount of DMF. The resultant 25 mixture was allowed to warm to RT and was stirred for 16-24 h. The reaction mixture was diluted with EtOAc and washed with water and brine. The organic layer was dried over sodium sulfate and concentrated, and the residue chromatographed on silica gel with hexanes/ethyl acetate as eluant to yield the alkylation product.

30


Heterocycles Used:

	2-ethyl-5,7-dimethyl-3H-imidazo[4,5-b]pyridine	Senanayake, C. H., et. al. <i>Heterocycles</i> 1996 , 42, 821-830.
	2-butyl-1,3-diazaspiro[4.4]non-1-en-4-one	Bernhart, C. A., et. al. <i>J. Med. Chem.</i> , 1993 , 36, 3371-3380.
	3,5-di-n-butyl-1,2,4-triazole	Reitz, D.B., et al. <i>Biorganic & Medicinal chemistry Letters</i> , 1994 , 4(1), 99-104
	2-propyl-1,3-diazaspiro[4.4]non-1-en-4-one	Bernhart, C. A., et. al. <i>J. Med. Chem.</i> , 1993 , 36, 3371-3380.
	7-oxo-2,4-dimethyl-5,6,7,8-tetrahydropyrido[2,3-d]pyrimidine	Hullar, T. L.; French, W. C. <i>J. Med. Chem.</i> , 1969 , 12, 424-426.
	2-ethyl-4(1H)-quinolone	Bradbury, R.H.; et. al. <i>J. Med. Chem.</i> 1992 , 35, 4027-4038.
	2-methyl-4(1H)-quinolone	commercially available
	2-propyl-4(1H)-quinolone	Bradbury, R.H.; et. al. <i>J. Med. Chem.</i> 1992 , 35, 4027-4038.
	2-ethyl-5,6,7,8-tetrahydro-4(1H)-quinolone	Bradbury, R.H.; et. al. <i>J. Med. Chem.</i> 1993 , 36, 1245-1254.
	methyl 2-(N-propylamino)pyridine-3-carboxylate	De, B.; et. al. <i>J. Med. Chem.</i> 1992 , 35, 3714-3717
	2,7-diethyl-5H-pyrazolo[1,5-b][1,2,4]triazole	U,S, Patent No. 5,475,114
	3-methoxy-2,6-dimethyl-4(4H)-pyridinone	Voss, G.; et. al. <i>Liebigs Ann Chem.</i> 1982 , 1466-1477
	2-butyl-4(3H)-quinazolinone	Allen, E. C. ; et. al <i>Biorganic & Medicinal Chemistry Let.</i> 1993 , 3(6), 1293-1298

	N- (2-butyl-3,4-dihydro-4-oxo-6-quinazolinonyl)-N'-isopropyl-N-methylurea	Laszlo, S. E.; et al <i>Biorganic & Medicinal Chemistry Let.</i> 1993 , 3(6), 1299-1304
---	---	---

General Method 5: Reductive Amination


5



To a mixture of an aromatic aldehyde (1.0 eq) and a primary amine (1.2 eq) in dichloromethane (0.1 M aldehyde concentration) was added 4 Å molecular sieves (5g per mmol aldehyde). [Alternately, a primary amine hydrochloride (1.2 eq) and triethylamine (1.2 eq) could be substituted for the primary amine free base.] The mixture was stirred vigorously for 1 h, after which sodium triacetoxyborohydride (1.5 eq) was added. The mixture was stirred vigorously at RT, while the course of the reaction was monitored by HPLC. If the reaction had not reached completion within several hours, additional sodium triacetoxyborohydride (1.0 eq) was added and monitoring was continued. When the reaction was complete the mixture was filtered through celite, aqueous sodium bicarbonate solution was added to the filtrate, and the aqueous layer was extracted with dichloromethane. The combined organic extracts were dried over sodium sulfate and evaporated. The crude residue was carried on without further purification.

In general, reductive amination with a 4-aminobutanoic acid resulted in a lactam product. In a few cases, cyclization was promoted by treatment of a 0.1 M solution of the crude amino acid product in dichloromethane with 1.0 eq of diisopropylcarbodiimide for 1 h at RT.

General Method 6: Amine Acylation



A 0.15 M solution of a primary or secondary amine (1.0 eq) and *N,N*-diisopropylethylamine (2.0 eq) in dichloromethane was treated at RT
5 with an acyl chloride (1.5 eq). After 1.5 h, methanol (10 eq) was added, followed by aqueous sodium carbonate solution. The aqueous layer was extracted with dichloromethane and the combined organic extracts were combined, dried over sodium sulfate, and concentrated. The residue was chromatographed on silica gel with hexanes/ethyl acetate as eluant to
10 provide the product tertiary amide.

General Method 7: Hydrolysis of SEM or MEM Sulfonamide
Protecting Groups Using Hydrochloric Acid/Ethanol



R = MEM, SEM

5

To a 0.1 M solution of a SEM- or MEM-protected N-heteroaryl sulfonamide in one volume of 95% EtOH was added an equal volume of 6N aqueous HCl, and the resulting solution was heated at reflux for 1 h. The reaction mixture was concentrated and the pH of the solution was
10 adjusted to pH 8 using aqueous sodium bicarbonate solution. It was then reacidified to pH 5 with glacial acetic acid. The mixture was extracted with three portions of ethyl acetate. The combined organic extracts were washed with water and brine, dried over sodium sulfate, and concentrated. The residue was purified by reverse-phase preparative
15 HPLC, or by silica gel chromatography using chloroform/methanol or hexanes/acetone as eluant.

General Method 8: Hydrolysis of SEM or MEM Sulfonamide
Protecting
Groups Using Hydrogen Chloride in Dioxane/Alcohol

20



R = MEM, SEM

A solution of a SEM- or MEM-protected N-heteroaryl sulfonamide
25 in one volume of absolute methanol or ethanol was treated with two volumes of 4 N hydrogen chloride/dioxane solution (final substrate

concentration 0.05 M). The resulting solution was heated at 55 °C for 16 h and was then concentrated. The residue was purified by reverse-phase preparative HPLC, or by extraction with ethyl acetate from aqueous potassium phosphate adjusted to pH 5-6, followed by silica gel

5 chromatography.

General Method 9: Cleavage of SEM or MEM Sulfonamide

Protecting

Groups Using Trimethylsilyl Iodide

10




R = MEM, SEM

To a 0.1 M solution of a SEM- or MEM-protected N-heteroaryl sulfonamide in acetonitrile was added trimethylsilyl chloride (8 eq) followed by sodium iodide (8 eq). The mixture was stirred at RT for 30 min and was then poured onto water and ethyl acetate. The organic layer was washed with saturated sodium sulfite and brine, and was then dried over sodium sulfate and concentrated. The residue was purified by reverse-phase preparative HPLC or by silica gel chromatography.

20

General Method 10: Cleavage of SEM Sulfonamide Protecting Groups Using Tetrabutylammonium Fluoride




25

To a 0.05 M solution of a SEM-protected N-heteroaryl sulfonamide in THF was added freshly activated 4Å molecular sieves (20 g per mmol

sulfonamide), followed by tetrabutylammonium fluoride (1.0 M solution in THF, 3 eq). The mixture was heated at 55 °C for 1-2 h, then was cooled and filtered through celite. The filter cake was rinsed with methanol, then aqueous potassium dihydrogen phosphate solution was added to the filtrate and the mixture partially concentrated. The residue was adjusted to pH 4-5 using dilute hydrochloric acid, and the mixture was extracted with two portions of ethyl acetate. The combined organic extracts were dried over sodium sulfate and concentrated. The residue was purified by reverse-phase preparative HPLC or by silica gel chromatography.

10

General Method 11: Reduction of Aryl Aldehydes to Benzylic Alcohols Using Sodium Borohydride




15

Sodium borohydride (0.5 eq) was added at 0 °C to a 0.2 M solution of an aromatic aldehyde in absolute ethanol or methanol. The mixture was allowed to warm to RT and stirred for 1-2 h. Aqueous potassium dihydrogen phosphate solution (or dilute hydrochloric acid) was added and the mixture was stirred for an additional 15 min. The mixture was partially concentrated and the residue partitioned between ethyl acetate and water. The aqueous layer was extracted twice with ethyl acetate and the combined organic extracts were dried over sodium sulfate and concentrated. The crude benzylic alcohol was either used directly or was purified by silica gel chromatography using hexanes/ethyl acetate as eluant.

25

General Method 12: Amide Formation using 1,1'-Carbonyldiimidazole


30



1,1'-Carbonyldiimidazole (2.0 eq) was added to a 0.1 M solution or suspension of an appropriate carboxylic acid (1.0 eq) in THF. The mixture was heated at 50 °C for 1h, and was then cooled to RT. An appropriate amine (5-10 eq) was then added, and the mixture was stirred at RT for 12 h. Ethyl acetate and aqueous sodium bicarbonate solution were added and the aqueous layer was extracted with ethyl acetate. The combined organic layers were dried over sodium sulfate and concentrated. The residue was purified by reverse-phase preparative HPLC or by silica gel chromatography.


General Method 13: Benzylic Bromination

Using N-Bromosuccinimide



To a 0.4 M solution of a methyl-substituted aromatic compound in carbon tetrachloride was added N-bromosuccinimide (1.05 eq) and benzoyl peroxide (0.03 eq), and the mixture was heated at reflux for 8-16 h. The mixture was cooled and filtered and the filtrate concentrated. The residue was purified by trituration with 3:1 hexanes/ethyl acetate, or by silica gel chromatography using hexanes/ethyl acetate as eluant to provide the mono-brominated product.

General Method 14: Reduction of an Aromatic Nitrile to an Aromatic Aldehyde Using DIBAL-H



DIBAL-H (1.5 M solution in toluene, 1.5 eq) was added dropwise at 0 °C to a 0.5 M solution of an aromatic nitrile (1.0 eq) in toluene or 9:1 toluene/dichloromethane. The solution was stirred at 0 °C for 1-4 h, and was then treated with excess methanol. After 15 min, 2N hydrochloric acid was added and the mixture was stirred vigorously for an additional 15 min. The layers were separated, and the aqueous layer was extracted with ethyl acetate. The combined organic layers were dried over sodium sulfate and concentrated to yield the crude aldehyde, which was either carried on crude or purified via silica gel chromatography using hexanes/ethyl acetate as eluant.

General Method 15: Ester Hydrolysis Using Lithium Hydroxide




15

A 0.25 M solution of an alkyl ester in 1:1 THF/water was treated with lithium hydroxide hydrate (1.5 eq) at RT. The mixture was stirred for 8-16 h and was then acidified with dilute hydrochloric acid. The product was either isolated by direct filtration from the reaction mixture, or by extraction with ethyl acetate, followed by drying of the organic layers with sodium sulfate, concentration, and silica gel chromatography using methanol/chloroform or hexanes/acetone as eluant.

General Method 16: Displacement of a Benzylic Bromide or Mesylate with Cyanide

25



Sodium cyanide (1.2 eq) was added at RT to a 1.0 M solution of a benzylic bromide or mesylate in DMF. The mixture was stirred at RT for

16 h. The reaction mixture was diluted with ethyl acetate and partitioned against aqueous sodium bicarbonate. The organic layer was dried over sodium sulfate and concentrated, and the residue chromatographed on silica gel with hexanes/ethyl acetate as eluant to yield the nitrile product.

5

General Method 17: Swern Oxidation of a Benzylic Alcohol to an Aromatic Aldehyde




10

Oxalyl chloride (1.5 eq) was added dropwise to a solution of DMSO (2.0 eq) in dichloromethane at $-78\text{ }^{\circ}\text{C}$. After 5 min, a solution of benzylic alcohol substrate (1.0 eq) in dichloromethane was added and the mixture (0.2 M final substrate concentration) was stirred at $-78\text{ }^{\circ}\text{C}$ for 15 min.

15 Triethylamine (4.0 eq) was added and the mixture was stirred and allowed to warm to RT. Aqueous sodium bicarbonate solution was added, the layers were separated, and the aqueous layer was extracted with one portion of dichloromethane. The combined organic layers were dried over sodium sulfate, concentrated, and the residue was purified by silica gel
20 chromatography using hexanes/ethyl acetate as eluant.

General Method 18: Reduction of an Aromatic Nitro Group to an Aromatic Amine Using Tin (II) Chloride Dihydrate

25



Tin (II) chloride dihydrate (4.0 eq) was added to a 0.05 M solution of an aromatic nitro compound in ethyl acetate and the resulting mixture was heated at $70\text{ }^{\circ}\text{C}$ for 45 min. The mixture was cooled, half-saturated
30 aqueous sodium carbonate solution was added, and the layers were separated. The aqueous layer was extracted once with ethyl acetate, and

the combined organic layers were dried over sodium sulfate and concentrated. The residue was chromatographed on silica gel using hexanes/ethyl acetate as eluant to provide the product aromatic amine.

5

General Method 19: Hydrolysis of a
2-Aryl-1,3-Dioxolane to an Aromatic Aldehyde




10

A 0.2 M solution of a 2-aryl-1,3-dioxolane (1.0 eq) in THF was treated with 1N hydrochloric acid (1.5 eq), and the resulting solution was heated at 55 °C for 16 h. The mixture was cooled and neutralized with aqueous sodium bicarbonate solution, then extracted with three portions of ethyl acetate. The combined organic extracts were dried over sodium sulfate and concentrated, and the crude aldehyde was used directly with no further purification.

15

General Method 20: Ester Formation Using
1,1'-Carbonyldiimidazole and DBU

20




25

1,1'-Carbonyldiimidazole (2.0 eq) was added to a 0.1 M solution or suspension of an appropriate carboxylic acid (1.0 eq) in THF. The mixture was heated at 50 °C for 1h. An appropriate alcohol (3.0 eq) was then added, followed by DBU (3.0 eq). The mixture was heated at 50 °C for 16 h and was then cooled. Ethyl acetate and 35% aqueous citric acid solution were added, and the organic layer was dried over sodium sulfate and

concentrated. The residue was purified by reverse-phase preparative HPLC or by silica gel chromatography.

General Method 21: Phase-Transfer Alkylation of Imidazoles




5

A solution of an appropriate imidazole (0.1 M) in toluene was treated with 50% aqueous sodium hydroxide solution (0.5 ml per mmol imidazole), tetrabutylammonium hydrogen sulfate (0.05 eq), and an appropriate benzylic alkyl bromide or mesylate (0.95 eq). The mixture was stirred vigorously at 40 °C for 24 h and was then cooled and filtered. Water was added and the aqueous layer was extracted with two portions of ethyl acetate. The combined organic extracts were dried over sodium sulfate and concentrated to provide the crude product, which was either purified by silica gel chromatography or was carried on crude. Imidazole used: 2-propyl-4,5,6,7-tetrahydro-8-oxocycloheptimidazole (Yanagisawa, T.; et. al. *Biorg. Med. Chem. Lett.* **1993**, 3, 1559-1564).

20




General Method 22: Imidazole or Phenol Alkylation




A solution of an appropriate imidazole (0.5 M) in DMF was treated with potassium carbonate (2.0 eq) and a benzylic alkyl bromide or mesylate (1.0 eq) at RT. The mixture was stirred at RT for 16 to 24 h. The solvent was evaporated and the residue was partitioned between

ethyl acetate and water. The ethyl acetate layer was dried over sodium sulfate and concentrated to provide the crude product, which was purified by silica gel chromatography or was carried on crude. When mixtures of N-1 and N-3 alkylation products were obtained, the regiochemistry of the alkylation was determined by NOESY spectroscopy.


Imidazoles Used:

	2-n-butyl-4-chloro-5-formylimidazole	Watson, S.P. <i>Synth. Comm.</i> , 1992 , 22, 2971-2977
	ethyl 2-n-propyl-4-ethylimidazole-5-carboxylate	Carini, D. J., WO 92/00977
	2-n-propyl-4-chloro-5-formylimidazole	Watson, S.P. <i>Synth. Comm.</i> , 1992 , 22, 2971-2977

General Method 23: Cleavage of SEM Sulfonamide Protecting Groups Using Cesium Fluoride



To a 0.05 M solution of a SEM-protected N-heteroaryl sulfonamide in DMF was added cesium fluoride (5.0 eq), and the resulting mixture was heated at 130 °C for 3 h. The reaction mixture was cooled and the solvent evaporated. Aqueous potassium dihydrogen phosphate solution was added (pH 4-5) and the mixture was extracted with three portions of ethyl acetate. The combined organic extracts were dried over sodium sulfate and concentrated. The residue was purified by reverse-phase preparative HPLC or by silica gel chromatography.

General Method 24: Sulfonylation of Aromatic Amines

5 To a 0.1 M solution of an aromatic amine (1.0 eq.) in dichloromethane at -30 °C was added triethylamine (2.6 eq), followed by a sulfonyl chloride (1.4 eq). The mixture was allowed to warm to RT over 3 hr. Aqueous sodium bisulfate was added (final pH 5) and the aqueous layer was extracted with dichloromethane. The combined organic extracts
10 were washed with water and brine, and were then dried over sodium sulfate and concentrated. The residue was chromatographed on silica gel using dichloromethane/methanol as eluant.

15 General Method 25: Oxidation of Aromatic Aldehydes to Carboxylic Acids



A 0.1 M solution of an aromatic aldehyde in 1:1 THF/water was
20 treated at 0 °C with sulfamic acid (1.5 eq) and sodium chlorite (1.5 eq). After 1 h aqueous potassium bisulfate solution was added and the mixture was extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate and concentrated to provide the crude carboxylic acid, which was used without further purification.

25

General Procedure: Purification by Anion Exchange Chromatography

Anion exchange chromatography was performed on Varian SAX cartridges (acetate form, 1.5-3 g) or United Chemical Technologies
30 CUQAX13M6-AC cartridges (acetate form, 3 g). Following a methanol rinse, the cartridge was loaded with a dichloromethane solution of crude

product. Elution of impurities with dichloromethane, followed by elution of the desired product with 1-3% TFA in dichloromethane or dichloromethane/methanol, provided the purified product.

5 General Procedure: Purification by Reverse-Phase Preparative HPLC

Reverse-phase preparative HPLC was performed with Shimadzu 8A liquid chromatographs using YMC S5 ODS columns (20 x 100, 20 x 250, or 30 x 250 mm). Gradient elution was performed with methanol/water mixtures in the presence of 0.1% TFA. In some cases a product eluting as a TFA salt was subsequently converted to the corresponding free base by extraction from aqueous sodium bicarbonate or sodium carbonate solution.

15 Analytical HPLC Methods Employed in Characterization of
 Examples

Analytical HPLC was performed on Shimadzu LC10AS liquid chromatographs using the following methods:

- 20 A. Linear gradient of 0 to 100% solvent B over 4 min, with 1 min hold
 at 100% B;
 UV visualization at 220 nm
 Column: YMC S5 ODS Ballistic 4.6 x 50 mm
 Flowrate: 4 ml/min
- 25 Solvent A: 0.1% trifluoroacetic acid, 90% water, 10% methanol
 Solvent B: 0.1% trifluoroacetic acid, 90% methanol, 10% water
- 30 B. Linear gradient of 0 to 100% solvent B over 30 min, with 5 min hold
 at 100% B;
 UV visualization at 254 nm
 Column: YMC S3 ODS 6 x 150 mm

Flowrate: 1.5 ml/min

Solvent A: 0.2% phosphoric acid, 90% water, 10% methanol

Solvent B: 0.2% phosphoric acid, 90% methanol, 10% water

- 5 C. Linear gradient of 0 to 100% solvent B over 4 min, with 1 min hold at 100% B
UV visualization at 220 nm
Column: YMC S5 ODS Ballistic 4.6 x 50 mm
Flowrate: 4 ml/min
- 10 Solvent A: 0.2% phosphoric acid, 90% water, 10% methanol
Solvent B: 0.2% phosphoric acid, 90% methanol, 10% water
- D. Linear gradient of 45 to 100% solvent B over 2 min, with 1 min hold at 100% B;
- 15 UV visualization at 220 nm
Column: Phenomenex Primesphere 4.6 x 30 mm
Flowrate: 5 ml/min
Solvent A: 0.2% phosphoric acid, 90% water, 10% methanol
Solvent B: 0.2% phosphoric acid, 90% methanol, 10% water
- 20
- E. Same conditions as (B), but with a linear gradient of 40 to 100% solvent B over 30 min, with 5 min hold at 100% B
- F. Same conditions as (B), but with a linear gradient of 70 to 100% solvent B over 30 min, with 5 min hold at 100% B
- 25
- G. Same conditions as (D), but with a linear gradient of 40 to 100% solvent B over 2 min, with 1 min hold at 100% B
- H. Linear gradient of 0 to 100% solvent B over 2 min, with 1 min hold at 100% B;
- 30

UV visualization at 220 nm

Column: Phenomenex Primesphere 4.6 x 30 mm

Flowrate: 5 ml/min

Solvent A: 0.1% trifluoroacetic acid, 90% water, 10% methanol

5 Solvent B: 0.1% trifluoroacetic acid, 90% methanol, 10% water

I. Same conditions as (B), but with a linear gradient of 50 to 100% solvent B over 30 min, with 5 min hold at 100% B

10 J. Same conditions as (C), but with a linear gradient of 0 to 100% solvent B over 8 min, with 1 min hold at 100% B

K. Same conditions as (D), but with a linear gradient of 0 to 100% solvent B over 2 min, with a 1 minute hold at 100% B.


15

Preparations

Preparation 1:

N-(3,4-Dimethyl-5-isoxazolyl)-4'-formyl-2'-(hydroxymethyl)-N-[(2-trimethylsiloxy)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

20



A. 4-Bromo-3-(bromomethyl)benzonitrile

25 The product was prepared according to General Method 13 starting from 12.0 g 4-bromo-3-methylbenzonitrile. Partial purification of the crude product was performed by trituration with 3:1 hexanes/ethyl acetate to afford 7.3 g of a slightly yellow solid, which was contaminated with approximately 20 mol% of the starting material.

B. 4-Bromo-3-(acetoxymethyl)benzonitrile

A mixture of **P1A** (7.3 g), potassium acetate (3.4 g), and DMF (10 ml) was stirred at RT for 16 h. Ethyl acetate was added and the mixture was washed with four portions of water, followed by one portion of brine.

- 5 The ethyl acetate layer was dried over sodium sulfate and concentrated. The solid residue was partially purified by crystallization from ethyl acetate, yielding 4.5 g of a slightly yellow solid.

C. 4-Bromo-3-(hydroxymethyl)benzaldehyde

- 10 **P1B** (4.4 g) was treated with DIBAL-H according to General Method 14, using 3.5 eq of the reducing agent rather than 1.5 eq. The crude product was an orange oil (4.8 g), judged by ¹H NMR to be approximately 75% pure by weight.

- 15 D. N-(3,4-Dimethyl-5-isoxazolyl)-4'-formyl-2'-(hydroxymethyl)-N-[(2-trimethylsiloxy)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

P1C (4.7 g) was subjected to Suzuki coupling according to General Method 1, yielding 7.6 g of the product as an orange oil following silica gel chromatography (2:1 hexanes/ethyl acetate eluant).

20


Preparation 2:

N-(3,4-Dimethyl-5-isoxazolyl)-4'-bromomethyl-2'-

(methoxymethyl)-N-

[(2-methoxyethoxy)methyl][1,1'-biphenyl]-2-sulfonamide

25



A. 4-Bromo-3-(bromomethyl)benzonitrile

The product was prepared according to General Method 13 starting from 19.6 g 4-bromo-3-methylbenzonitrile. After cooling the mixture was filtered and the filtrate was washed with H₂O and brine, dried and

5 concentrated. The residue was chromatographed on silica gel using 100:3 and then 100:10 hexane/EtOAc to afford **P2A** (16 g, 58%). R_f=0.15, silica gel, 10:1 hexane/EtOAc.

B. 4-Bromo-3-(Methoxymethyl)benzonitrile

10 To a solution of **P2A** (6.95 g, 25.28 mmol) in 10 ml DMF, NaOMe (25 wt. % in MeOH, 6.94 ml, 30.3 mmol) was added dropwise. The reaction mixture was stirred at RT for 3 h. Ethyl acetate (100 ml) and hexanes (50 ml) were added, and the mixture was washed twice with water and once with brine. The organic layer was dried over sodium
15 sulfate and concentrated. The residue was chromatographed on silica gel using 100:6 hexane/EtOAc to afford **P2B** (4.70 g, 82%). R_f=0.5, silica gel, 5:1 hexane/EtOAc.

C. 4-Bromo-3-(methoxymethyl)benzaldehyde

20 **P2B** (7.0 g) was treated with DIBAL-H according to General Method 14, using THF instead of toluene as solvent. The crude product was purified by silica gel chromatography using 11:1 hexanes/ethyl acetate as eluant to give 6.2 g **P2C** as a colorless gum. R_f=0.4, silica gel, 5:1 Hexane/EtOAc.

25

D. N-(3,4-Dimethyl-5-isoxazolyl)-4'-formyl-2'-(methoxymethyl)-N-[(2-methoxyethoxy)methyl][1,1'-biphenyl]-2-sulfonamide

P2C (6.2 g) was subjected to Suzuki coupling according to General Method 1, giving **P2D** as an oil in 83% yield after silica gel
30 chromatography.

E. N-(3,4-Dimethyl-5-isoxazolyl)-4'-hydroxymethyl-2'-(methoxymethyl)-N-[(2-methoxyethoxy)methyl][1,1'-biphenyl]-2-sulfonamide


P2D (2.8 g) was reduced with sodium borohydride according to
5 General Method 11, to provide 2.8 g **P2E**.

F. N-(3,4-Dimethyl-5-isoxazolyl)-4'-bromomethyl-2'-(methoxymethyl)-N-[(2-methoxyethoxy)methyl][1,1'-biphenyl]-2-sulfonamide

P2E (2.8 g) was treated with triphenylphosphine and carbon
10 tetrabromide according to General Method 2, providing the title compound (2.3 g) in 72% yield.

Preparation 3:

2'-Cyano N-(3,4-dimethyl-5-isoxazolyl) -4'-(hydroxymethyl)-N-[(2-trimethylsiloxy)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide



A. 4'-(Acetoxymethyl)-N-(3,4-dimethyl-5-isoxazolyl) -2'-formyl-N-[(2-trimethylsiloxy)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide acetoxime

Hydroxylamine hydrochloride (1.13 g) was added to a solution of 7.0 g **P4** in 20 ml pyridine and the mixture was stirred at RT for 2 h. Acetic anhydride (5.1 ml) was added and the mixture was stirred for 1 h at RT.
25 Ethanol (5 ml) was added and the mixture was concentrated under reduced pressure. The residue was taken up in ethyl acetate and washed twice with 0.1 N hydrochloric acid, twice with half-saturated aqueous sodium carbonate solution, and once with brine. The ethyl acetate layer

was dried over sodium sulfate and concentrated to provide **P3A** as an orange oil.

B. 4'-(Acetoxymethyl)-2'-cyano-N-(3,4-dimethyl-5-isoxazolyl)-N-[(2-trimethylsiloxy)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

P3A was dissolved in 75 ml acetonitrile, DBU (4.0 ml) was added, and the mixture was stirred at RT for 14 h. The mixture was concentrated, and the residue was taken up in ethyl acetate and washed twice with 0.1 N hydrochloric acid, then once with half-saturated aqueous sodium carbonate solution. The ethyl acetate layer was dried over sodium sulfate and concentrated to give **P3B** as an orange oil.

C. 2'-Cyano N-(3,4-dimethyl-5-isoxazolyl) -4'-(hydroxymethyl)-N-[(2-trimethylsiloxy)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

P3B was dissolved in 150 ml methanol, potassium carbonate (1.5 g) was added, and the mixture was stirred at RT for 2 h. 2N Hydrochloric acid (5.5 ml) was added and the mixture was concentrated. The residue was taken up in ethyl acetate and partitioned against aqueous sodium bicarbonate solution. The ethyl acetate layer was dried over sodium sulfate and concentrated to yield 5.7 g of the crude title compound, which was used without further purification.




Preparations 4 through 22 were performed by application of the General Methods and are listed in the following Table.

Preparations by General Methods

No.	Structure	Name	Starting Material	Starting Material Reference	General Methods Applied (yield,%)
P4		N-(3,4-Dimethyl-5-iso-xazolyl)-2'-formyl-4'-(hydroxymethyl)-N-[[2-(trimethylsilyl)ethoxy]methyl][1,1'-biphenyl]-2-sulfonamide		Zhang, H.-Y. et al., <i>Tetrahedron</i> , 1994 , 50, 11339-11362.	19 (95); 1 (81)
P5		2'-Chloro-N-(3,4-dimethyl-5-iso-xazolyl)-4'-formyl-N-[[2-(trimethylsilyl)ethoxy]methyl][1,1'-biphenyl]-2-sulfonamide		Casida, J. E.; Elliott, M.; Pullmann, D. A., EP 294229	1 (83)
P6		N-(3,4-Dimethyl-5-isoxazolyl)-4'-formyl-2'-(trifluoromethyl)-N-[[2-(trimethylsilyl)ethoxy]methyl][1,1'-biphenyl]-2-sulfonamide		DoAmaral, J. R.; French, F. A.; Blanz, E. J. Jr.; French, D. A. <i>J. Med. Chem.</i> 1971 , 9, 862-866.	1 (11)
P7		N-(3,4-Dimethyl-5-isoxazolyl)-4'-formyl-N-[(2-methoxyethoxy)methyl][1,1'-biphenyl]-2-sulfonamide		Pine, S. H.; et. al. <i>J. Org. Chem.</i> 1971 , 36, 984-91.	1 (77)
P8		N-(3,4-Dimethyl-5-isoxazolyl)-2'-fluoro-4'-formyl-N-[(2-methoxyethoxy)methyl][1,1'-biphenyl]-2-sulfonamide		Palmer, C. J.; Casida, J. E.; Larkin, J. P. US Patent 5,061,726	1 (50)

P9		2'-[[[(3,4-Dimethyl-5-isoxazolyl)][2-(trimethylsilyl)ethoxy]methyl]amino]sulfonyl]-2-(trifluoromethyl)[1,1'-biphenyl]-4-methanol methanesulfonate	P6		11 (90); 3 (75)
P10		2-Chloro-2'-[[[(3,4-dimethyl-5-isoxazolyl)][2-(trimethylsilyl)ethoxy]methyl]amino]sulfonyl][1,1'-biphenyl]-4-methanol methanesulfonate	P5		11 (90); 3 (83)
P11		4'-(Bromomethyl)-N-(3,4-dimethyl-5-isoxazolyl)-2'-fluoro-N-[(2-methoxyethoxy)methyl][1,1'-biphenyl]-2-sulfonamide	P8		11 (98); 2 (67)
P12		2'-[[[(3,4-Dimethyl-5-isoxazolyl)][2-(trimethylsilyl)ethoxy]methyl]amino]-sulfonyl]-4-(hydroxymethyl)[1,1'-biphenyl]-2-methanol methanesulfonate	P1		3 (99); 11 (76)
P13		4'-[(Methanesulfonyloxy)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-N-[(2-methoxyethoxy)methyl]-2'-methyl[1,1'-biphenyl]-2-sulfonamide	P7		11; 3 (62)
P14		2'-[[[(3,4-Dimethyl-5-isoxazolyl)][2-(trimethylsilyl)ethoxy]methyl]amino]sulfonyl]-2-(hydroxymethyl)[1,1'-biphenyl]-4-methanol methanesulfonate	P4		3 (90); 11 (35)


P1 5		N-(3,4-Dimethyl-5-isoxazolyl)-4'-formyl-2'-methyl-N-[[2-(trimethylsilyl)ethoxy]methyl][1,1'-biphenyl]-2-sulfonamide		Pine, S. H.; et. al. <i>J. Org. Chem.</i> 1971 , 36, 984-91.	1 (52)
P1 6		N-(3,4-Dimethyl-5-isoxazolyl)-4'-formyl-N-[[2-(trimethylsilyl)ethoxy]methyl][1,1'-biphenyl]-2-sulfonamide	4-bromo-benz-aldehyde		1 (67)
P1 7		N-(3,4-Dimethyl-5-isoxazolyl)-4'-formyl-N-[[2-methoxyethoxy]methyl][1,1'-biphenyl]-2-sulfonamide	4-bromo-benz-aldehyde		1 (80)
P1 8		4'-Bromomethyl-N-(3,4-dimethyl-5-isoxazolyl)-N-(2-methoxyethoxymethyl)[1,1'-biphenyl]-2-sulfonamide	P17		11,2 (80)
P1 9		N-(3,4-Dimethyl-5-isoxazolyl)-4'-[(methanesulfonyl)oxy]methyl-N-[[2-(trimethylsilyl)ethoxy]methyl][1,1'-biphenyl]-2-sulfonamide	P16		11 (83); 3 (90)
P2 0		N-(3,4-Dimethyl-5-isoxazolyl)-4'-formyl-N-[[2-(trimethylsilyl)ethoxy]methyl][1,1'-biphenyl]-2-sulfonamide	P2C		1 (72)

P2 1		N-(3,4-Dimethyl-5-isoxazolyl)-2'-formyl-4'-(hydroxymethyl)-N-[[2-methoxyethoxy)methyl][1,1'-biphenyl]-2-sulfonamide		Zhang, H.-Y. et al., <i>Tetrahedron</i> , 1994 , <i>50</i> , 11339-11362.	19 (95); 1 (77)
P2 2		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-di-methyl-5-isoxazolyl)-N-[[2-methoxyethoxy)methyl]-2'-nitro[1,1'-biphenyl]-2-sulfonamide	4-bromo-3-nitro-toluene		13 (53); 4 (74); 1 (50)

Examples

Example 1

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-
dimethyl-3-isoxazoly)[1,1'-biphenyl]-2-sulfonamide



A. N-(4,5-Dimethyl-3-isoxazoly)-4'-(hydroxymethyl)-N-[(2-
methoxyethoxy)methyl][1,1'-biphenyl]-2-sulfonamide

4-Bromobenzyl alcohol (750 mg, 4.0 mmol) was coupled with [2-
 [(4,5-dimethyl-3-isoxazoly)(2-
 methoxyethoxy)methyl]amino)sulfonyl]phenyl]boronic acid (1.0 g, 2.7
 mmol) according to General Method 1. The crude residue was
 chromatographed on silica gel using 3:4 hexane/EtOAc to afford **1A** (730
 mg, 66%) as a colorless gum: $R_f=0.26$, silica gel, 2:3 hexane/EtOAc.

B. 4'-Bromomethyl-N-(4,5-dimethyl-3-isoxazoly)-N-[(2-
methoxyethoxy)methyl][1,1'-biphenyl]-2-sulfonamide

1A (730 mg, 1.64 mmol) was converted to the corresponding
 bromide according to General Method 2. The crude residue was
 chromatographed on silica gel using 4:1 hexane/EtOAc to afford **1B** (750
 mg, 90%) as a colorless gum: $R_f=0.66$, silica gel, 1:1 hexane/EtOAc.

C. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-N-(2-methoxyethoxymethyl)[1,1'-biphenyl]-2-sulfonamide


1B (255 mg, 0.5 mmol) was reacted with 2-butyl-1,3-diazaspiro[4.4]non-1-en-4-one hydrochloride according to General Method 4. The crude residue was chromatographed on silica gel using 3:4 hexane/EtOAc to afford **1C** as a gum: $R_f=0.32$, silica gel, 1:1 hexane/EtOAc.

D. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)[1,1'-biphenyl]-2-sulfonamide

1C was subjected to deprotection according to General Method 7. The crude residue was purified by preparative HPLC to provide the title compound as a white solid (130 mg, 49%, for two steps): mp 77-81°C. Analysis calculated for $C_{29}H_{34}N_4O_4S \cdot 1.0H_2O$: Calc'd: C, 63.02; H, 6.57; N, 10.14; S, 5.80. Found: C, 62.75; H, 6.16; N, 9.85; S, 5.54.

Example 2

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-[(methylamino)methyl][1,1'-biphenyl]-2-sulfonamide



A. 4-Bromo-3-formyl-benzonitrile

To a solution of 4-bromo-3-methyl benzonitrile (14.0 g, 71.4 mmol) in carbon tetrachloride (200 mL), N-bromosuccinimide (13.98 g, 78.55 mmol) and benzoyl peroxide (700 mg) were added, and the mixture was heated at reflux for 8 h while illuminating the solution with a sun lamp. The mixture was cooled and filtered. The filtrate was concentrated to provide a light yellow solid (21 g) which was used in the next step without any further purification.

To a solution of the crude compound (21 g) obtained above in anhydrous DMSO (30 mL) under argon, was added anhydrous trimethylamine N-oxide (6.97 g, prepared as described in Soderquist et al., *Tetrahedron Letters*, 27, 3961 (1986)), and the mixture was stirred at 55°C for 48 h. The mixture was then cooled and added to ice/water (150 mL) and the resulting aqueous mixture was extracted with EtOAc (3 x 100 mL). The combined organic extracts were washed once with brine (100 mL), dried and evaporated. The residue was chromatographed on silica gel using 8:1 hexanes/EtOAc to afford **2A** as a white solid (6.1 g, 47% for two steps).

B. 4'-Cyano-2'-formyl-N-(3,4-dimethyl-5-isoxazolyl)-N-(2-methoxyethoxymethyl) [1,1'-biphenyl]-2-sulfonamide

2A (3.0 g, 14 mmol) was subjected to Suzuki coupling with N-[(2-methoxyethoxy)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzenesulfonamide according to General Method 1. The crude residue was chromatographed on silica gel using 2:1 hexane/EtOAc to afford **3B** (4.5 g, 68%) as a colorless gum.

C. 4'-Cyano-2'-(N-methylaminomethyl)-N-(3,4-dimethyl-5-isoxazolyl)-N-(2-methoxyethoxymethyl) [1,1'-biphenyl]-2-sulfonamide

To **2B** (2.2 g, 4.69 mmol), MeNH₂ (8.03 M in EtOH, 2.33 mL, 18.74 mmol) and 3A molecular sieves in CH₂Cl₂ (47 mL), glacial acetic acid (1.13

g, 18.74 mmol) was added followed by NaB(AcO)₃H (3.98 g, 18.74 mmol). The reaction mixture was stirred at RT overnight, diluted with EtOAc and filtered through celite. The filtrate was washed with H₂O, dried and concentrated. The residue was chromatographed on silica gel using 100:5 CH₂Cl₂/MeOH to afford **2C** (1.24 g, 55%) as a colorless gum: R_f=0.2, silica gel, 100:5 CH₂Cl₂/MeOH.

D. 2'-(N-*t*-Butoxycarbonyl-N-methylaminomethyl)-4'-cyano-N-(3,4-dimethyl-5-isoxazolyl)-N-(2-methoxyethoxymethyl) [1,1'-biphenyl]-2-sulfonamide

To **2C** (1.3 g, 2.68 mmol), triethylamine (434 mg, 4.3 mmol) and 4-dimethylaminopyridine (33 mg, 0.27 mmol) in THF (10 mL) was added a solution of di-*t*-butyl dicarbonate (703 mg, 3.22 mmol) in THF (10 mL) dropwise. The reaction mixture was stirred at RT for 3 h. 10% aqueous NH₄Cl was added, and the mixture was extracted with EtOAc. The extracts were washed with H₂O and brine, dried and concentrated. The residue was chromatographed on silica gel using 7:4 hexane/EtOAc to afford **2D** (1.1 g, 70%) as a colorless gum: R_f=0.57, silica gel, 2:3 hexane/EtOAc.

E. 2'-(N-*t*-Butoxycarbonyl-N-methylaminomethyl)-N-(3,4-dimethyl-5-isoxazolyl)-4'-formyl-N-(2-methoxyethoxymethyl) [1,1'-biphenyl]-2-sulfonamide

To **2D** (1.03 g, 1.76 mmol) in THF (18 mL), diisobutylaluminum hydride (1M in CH₂Cl₂, 5.29 mL, 5.29 mmol) was added dropwise. The reaction mixture was stirred at RT for 4 h. MeOH (20 mL) was added, and the mixture was stirred for 20 min. The mixture was then added into H₂O and extracted with EtOAc. The combined extracts were washed with H₂O and brine, dried and concentrated. The residue was chromatographed on silica gel using 5:6 hexane/EtOAc to afford **2E** (325 mg, 31%) as a colorless gum: R_f=0.37, silica gel, 1:1 hexane/EtOAc.

F. 2'-(N-*t*-Butoxycarbonyl-N-methylaminomethyl)-4'-Hydroxymethyl-N-(3,4-dimethyl-5-isoxazolyl)-N-(2-methoxyethoxymethyl) [1,1'-biphenyl]-2-sulfonamide

5 **2E** was reduced using sodium borohydride according to General Method 11 to provide **2F**, which was used in the next reaction step without any purification.

G. 4'-Bromomethyl-2'-(N-*t*-butoxycarbonyl-N-methylaminomethyl)-N-(3,4-dimethyl-5-isoxazolyl)-N-(2-methoxyethoxymethyl) [1,1'-biphenyl]-2-sulfonamide

2F was treated with carbon tetrabromide and triphenylphosphine according to General Method 2 to provide **2G** in 78% yield.

15 H. 2'-(N-*t*-Butoxycarbonyl-N-methylaminomethyl)-4'-[(2-butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-N-(2-methoxyethoxymethyl) [1,1'-biphenyl]-2-sulfonamide


20 **2G** was treated with 2-*n*-butyl-1,3-diazaspiro[4.4]non-1-en-4-one hydrochloride according to General Method 4 to provide **2H** in 79% yield.

I. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-[(methylamino)methyl][1,1'-biphenyl]-2-sulfonamide

25 **2H** (170 mg, 0.22 mmol) was deprotected according to General Method 7 to provide the title compound in 67% yield: R_f =0.39, silica gel, 10:1 CH₂Cl₂/MeOH; mp: 194-200 °C; LRMS (m/z) 578 (MH⁺).

Example 3

30 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-formyl-N-(3,4-dimethyl-5-isoxazolyl)-[1,1'-biphenyl]-2-sulfonamide



A. 4'-Cyano-2'-(1,3-dioxolan-2-yl)-N-(3,4-dimethyl-5-isoxazolyl)-N-(2-methoxyethoxymethyl) [1,1'-biphenyl]-2-sulfonamide

A mixture of **2B** (1.28 g, 2.73 mmol), ethylene glycol (1.69 g, 27.3 mmol) and *p*-toluenesulfonic acid (38 mg) in toluene (30 mL) was heated at 130°C for 5 h, while a Dean-Stark water separator was used. After cooling, the mixture was diluted with EtOAc. The organic liquid was separated and washed with H₂O and brine, dried and concentrated. The residue was chromatographed on silica gel using 5:4 hexane/EtOAc to afford **3A** (1.1 g, 79%) as a colorless gum: *R_f*=0.57, silica gel, 1:2 hexane/EtOAc.

B. 2'-(1,3-Dioxolan-2-yl)-4'-formyl-N-(3,4-dimethyl-5-isoxazolyl)-N-(2-methoxyethoxymethyl) [1,1'-biphenyl]-2-sulfonamide

To **3A** (1.1 g, 2.14 mmol) in THF (21 mL) at 0°C was added DIBAL-H (1M in CH₂Cl₂, 4.28 mL 4.28 mmol) dropwise. The reaction was stirred at RT overnight. MeOH (20 mL) was added and the reaction was stirred for 5 min. The mixture was poured into cold 0.1 N HCl solution (150 mL), shaken for 5 min, and then extracted with 3:1 EtOAc/hexane. The combined organic extracts were washed with H₂O and brine, dried and concentrated. The residue was chromatographed on silica gel using 3:4 hexane/EtOAc to afford **3B** (710 mg, 64%) as a colorless gum: *R_f*=0.45, silica gel, 2:3 hexane/EtOAc.

C. 2'-(1,3-Dioxolan-2-yl)-4'-hydroxymethyl-N-(3,4-dimethyl-5-isoxazolyl)-N-(2-methoxyethoxymethyl) [1,1'-biphenyl]-2-sulfonamide

3B (710 mg, 1.4 mmol) was subjected to sodium borohydride reduction according to General Method 11 to afford **3C**, which was used for the next reaction step without further purification.

D. 4'-Bromomethyl-2'-(1,3-dioxolan-2-yl)-N-(3,4-dimethyl-5-isoxazolyl)-N-(2-methoxyethoxymethyl) [1,1'-biphenyl]-2-sulfonamide

3C was treated with carbon tetrabromide and triphenylphosphine according to General Method 2. The crude residue was chromatographed on silica gel using 3:2 hexane/EtOAc to afford **3D** (750 mg, 94%) as a colorless gum: $R_f=0.74$, silica gel, 1:2 hexane/EtOAc.

E. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-(1,3-dioxolan-2-yl)-N-(3,4-dimethyl-5-isoxazolyl)-N-(2-methoxyethoxymethyl) [1,1'-biphenyl]-2-sulfonamide

3D (750 mg, 1.3 mmol) was treated with 2-*n*-butyl-1,3-diazaspiro[4.4]non-1-en-4-one hydrochloride (387 mg, 1.68 mmol) according to General Method 4. The crude residue was chromatographed on silica gel using 100:1.7 $\text{CH}_2\text{Cl}_2/\text{MeOH}$ to afford **3E** as a gum (830 mg, 93%): $R_f=0.40$, silica gel, 100:5 $\text{CH}_2\text{Cl}_2/\text{MeOH}$.


F. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-formyl-N-(3,4-dimethyl-5-isoxazolyl)-[1,1'-biphenyl]-2-sulfonamide

3E (830 mg, 1.20 mmol) was subjected to deprotection according to General Method 7. The crude residue was chromatographed on silica gel using 100:1.5 and then 100:4 $\text{CH}_2\text{Cl}_2/\text{MeOH}$ to afford the title compound as a gum (480 mg, 72%): $R_f=0.16$, silica gel, 100:5 $\text{CH}_2\text{Cl}_2/\text{MeOH}$.

Example 4

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-
dimethyl-5-isoxazolyl)-2'-[(3,3-dimethyl-2-oxo-1-
pyrrolidinyl)methyl][1,1'-biphenyl]-2-sulfonamide

5



To **3F** (110 mg, 0.20 mmol) in CH_2Cl_2 (4 mL) was added 4-amino-2,2-dimethylbutanoic acid hydrochloride (98 mg, 0.59 mmol) [Scheinmann, et al., *J. Chem. Research (S)*, 414-415 (1993)] and 3Å molecular sieves, followed by glacial acetic acid (35 mg, 0.59 mmol) and then sodium acetate (48 mg, 0.59 mmol). The mixture was stirred for 8 minutes, and $\text{NaB}(\text{AcO})_3\text{H}$ (124 mg, 0.59 mmol) was then added. The reaction mixture was stirred at RT for 2 h, diluted with EtOAc and filtered through celite. The filtrate was washed with H_2O and brine, dried and concentrated. This material was dissolved in CH_2Cl_2 (6 mL) and 1,3-diisopropylcarbodiimide (32 mg, 0.25 mmol) was added. The reaction mixture was stirred at RT for 2 h and diluted with CH_2Cl_2 , washed with H_2O and brine, dried and concentrated. The residue was purified by preparative HPLC to provide the title compound as a white solid (40 mg, 31%, for two steps): mp 104-110°C. Analysis calculated for $\text{C}_{36}\text{H}_{45}\text{N}_5\text{O}_5\text{S} \cdot 0.8 \text{ H}_2\text{O}$: Calc'd: C, 64.13; H, 6.97; N, 10.39; S, 4.75. Found: C, 64.18; H, 6.60; N, 10.23; S, 4.50.

Example 54'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-formyl-N-(3,4-dimethyl-5-isoxazolyl)-[1,1'-biphenyl]-2-sulfonamide (Alternative Preparation for 3F)

5

A. 2-[(2'-Bromo-5'-formyl)phenyl])-1,3-dioxolane

DIBAL-H (1.0 M solution in toluene, 445 mL, 445 mmol, 1.1 eq) was added over 30 minutes to a solution of 2-[(2'-bromo-5'-cyano)phenyl])-1,3-dioxolane (103 g, 404 mmol, 1.0 eq) [Zhang, H.-Y. et al., *Tetrahedron*, 50, 11339-11362 (1994)] in toluene (2.0 L) at -78 °C. The solution was allowed to warm to 0 °C. After 1 hour, a solution of Rochelle's salt (125 g) in water (200 mL) was added, and the mixture was allowed to warm to room temperature and was stirred vigorously for 16 h. The organic layer was concentrated and the residue partitioned between ethyl acetate (1 L) and 1 N hydrochloric acid (800 mL). The organic layer was washed with saturated aqueous sodium bicarbonate (800 mL), dried over sodium sulfate, and then concentrated to give 70.5 g of crude **5A** as a yellow solid, which was used without further purification.

20 B. 2-[(2'-Bromo-5'-hydroxymethyl)phenyl])-1,3-dioxolane

Sodium borohydride (3.66 g, 96.7 mmol, 0.5 eq) was added to a solution of crude **5A** (49.7 g, approximately 193 mmol, 1.0 eq) in absolute ethanol (1300 mL) at 0 °C. After 2 hours, a solution of 10% aqueous sodium dihydrogen phosphate (50 mL) was added and the mixture was stirred and allowed to warm to room temperature. The mixture was concentrated, then partitioned between ethyl acetate (800 mL) and saturated aqueous sodium bicarbonate (500 mL). The organic layer was dried over sodium sulfate and concentrated to give 49.0 g of crude **5B** as a yellow oil, which was used without further purification.

30

C. 2-[(2'-Bromo-5'-bromomethyl)phenyl]-1,3-dioxolane

Triphenylphosphine (52.7 g, 199 mmol, 1.05 eq) was added in portions over 15 minutes to a solution of crude **5B** (49.0 g, approximately 189 mmol, 1.0 eq) and carbon tetrabromide (69.0 g, 208 mmol, 1.1 eq) in THF at 0 °C. After 2 hours, saturated aqueous sodium bicarbonate solution (20 mL) was added, and the mixture was allowed to warm to room temperature and was then concentrated. Ether (500 mL) was added, and the resulting mixture was filtered. The filtrate was dried over magnesium sulfate and concentrated. The residue was chromatographed on silica gel (8:1 hexanes/ethyl acetate as eluant) to give **5C** as a white solid (31.1 g, 51% yield from 2-[(2'-bromo-5'-cyano)phenyl]-1,3-dioxolane).

D. 2-(1,3-Dioxolan-2-yl)-4-[(2-*n*-butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]bromobenzene

Sodium hydride (60% dispersion in mineral oil, 9.65 g, 241 mmol, 2.5 eq) was added in portions over 15 minutes to a mixture of 2-*n*-butyl-1,3-diazaspiro[4.4]non-1-en-4-one hydrochloride (18.7 g, 96.5 mmol, 1.0 eq) in DMF (400 mL) at 0°C. The mixture was stirred and allowed to warm to room temperature over 15 minutes. To this mixture was added via canula a solution of **5C** (31.1 g, 96.5 mmol, 1.0 eq) in DMF (100 mL). After 14 hours, the mixture was concentrated *in vacuo* and partitioned between ethyl acetate (500 mL) and 10% aqueous sodium dihydrogen phosphate (300 mL). The organic layer was dried over sodium sulfate and concentrated to give crude **5D** as an orange oil (42.7 g), which was used without further purification.

E. 4-[(2-*n*-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2-formyl-bromobenzene

A solution of crude **5D** (6.0 g, approximately 13.6 mmol, 1.0 eq) in THF (180 mL) and 1N hydrochloric acid (30 mL) was heated at 65°C for 1.5 hours. The mixture was cooled and then treated with saturated

aqueous sodium carbonate solution (75 mL) and ethyl acetate (200 mL). The organic layer was removed and dried over sodium sulfate, concentrated, and then further dried azeotropically with toluene to give **5E** as a crude yellow oil (8.2 g) which contained a small amount of toluene.

5 This material was used without further purification.

F. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-formyl-N-(3,4-dimethyl-5-isoxazolyl)-N-(2-methoxyethoxymethyl)[1,1'-biphenyl]-2-sulfonamide

10 Palladium catalyzed Suzuki coupling of **5E** and [2-[(3,4-dimethyl-5-isoxazolyl)[(2-methoxyethoxy)methyl]amino]sulfonyl]phenyl]boronic acid was performed according to General Method 1 to yield **5F** in 60% yield.

G. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-formyl-N-(3,4-dimethyl-5-isoxazolyl)-[1,1'-biphenyl]-2-sulfonamide

15 Deprotection of **5F** according to General Method 7 provided the title compound (**5G** = **3F**) in 73% yield: $R_f=0.2$ (silica gel using $\text{CH}_2\text{Cl}_2/\text{MeOH}$ [100:5]).

20

Example 6

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-[(3,3-dimethyl-2-oxo-1-pyrrolidinyl)methyl][1,1'-biphenyl]-2-sulfonamide (Alternative Preparation for 4)

25


The title compound (**6** = **4**) was obtained as a white solid from **5G** using a procedure similar to that described in Example 4 (945 mg, 35%): mp 104-110°C. Analysis calculated for $\text{C}_{36}\text{H}_{45}\text{N}_5\text{O}_5\text{S} \cdot 0.8 \text{ H}_2\text{O}$: Calc'd: C, 64.13; H, 6.97; N, 10.39; S, 4.75. Found: C, 64.18; H, 6.60; N, 10.23; S, 4.50.

30

Example 7

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-
dimethyl-5-isoxazolyl)-2'-[(3-methyl-2-oxo-1-
imidazolidinyl)methyl][1,1'-biphenyl]-2-sulfonamide

5



A. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-
dimethyl-5-isoxazolyl)-N-(2-methoxyethoxymethyl)-2'-[(3-methyl-2-
oxo-1-imidazolidinyl)methyl][1,1'-biphenyl]-2-sulfonamide

10

To a solution of 0.3 g (0.46 mmol) of **5F** in CH₂Cl₂ (15 mL) was added 3Å molecular sieves (1 mL), N-methyl ethylenediamine (0.051 g, 0.69 mmol) and glacial acetic acid (0.828 g, 1.38 mmol), and the resulting mixture was stirred under argon for 15 min. Sodium triacetoxymethylborohydride (0.292 g, 1.38 mmol) was then added to the mixture, and the resulting mixture was stirred at room temperature for 3 h. The solution was then filtered through celite, and the celite was washed with CH₂Cl₂ (25 mL). The combined filtrates were washed with water (2x50 mL), dried and evaporated to afford a colorless gum (0.32 g). This gum was redissolved in CH₂Cl₂ (10 mL), and carbonyldiimidazole was then added (0.112 g, 0.69 mmol). The resulting mixture was stirred at room temperature for 24 h. The mixture was then washed with water (15 mL) and dried and evaporated to provide a colorless gum. Purification on silica gel using 1:1 hexane: EtOAc then provided **7A** (0.17 g, 50%) as a colorless gum.


25

B. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-[(3-methyl-2-oxo-1-imidazolidinyl)methyl][1,1'-biphenyl]-2-sulfonamide

To a solution of **7A** (0.165 g, 0.225 mmol) in 95% EtOH (2 mL) was
 5 added 6N aqueous hydrochloric acid (2 mL), and the resulting solution
 was heated at reflux for 1 h. The mixture was then neutralized with
 aqueous sodium bicarbonate to pH 7 and then reacidified to pH 6 using
 aqueous sodium bisulfate. The mixture was then extracted with EtOAc (3
 x 30 mL). The combined organic extracts were then washed once with
 10 water and dried and evaporated. The residue was purified on silica gel
 using 5% MeOH in methylene chloride to provide the title compound (0.13
 g, 89%) as a light yellow solid: ¹H NMR (CDCl₃) δ 0.89 (t, *J*=7.0 Hz, 3H),
 1.26-1.60 (m, 4H), 1.87 (s, 3H), 1.97 (m, 8H), 2.16 (s, 3H), 2.35 (t, *J*=7.6 Hz,
 2H), 2.72 (s, 3H), 3.29 (br s, 4H), 4.10 (s, 2H), 4.72 (m, 2H), 7.13-7.92 (m,
 15 7H).

Example 8

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,3-dimethyl-2-oxo-1-pyrrolidinyl)methyl]-N-(2-pyrazinyl) [1,1'-biphenyl]-2-sulfonamide



A. N-tert-Butyl-4'-[(2-n-butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-formyl[1,1'-biphenyl]-2-sulfonamide

Crude **5E** (8.3 g, approximately 13.6 mmol) was subjected to Suzuki coupling with [2-(*N-tert*-butylsulfamoyl)phenyl]boronic acid according to General Method 1. The crude residue was chromatographed on triethylamine-deactivated silica gel (1:1 hexanes/ethyl acetate eluant) to yield **8A** (5.03 g, 71% from **5C**) as an orange oil.

B. *N-tert*-Butyl-4'-[(2-*n*-butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,3-dimethyl-2-oxo-1-pyrrolidinyl)methyl][1,1'-biphenyl]-2-sulfonamide

A mixture of **8A** (2.0 g, 3.81 mmol, 1.0 eq), 4-amino-2,2-dimethylbutanoic acid hydrochloride (1.28 g, 7.64 mmol, 2.0 eq), freshly-activated 3Å molecular sieves (5.0 g), and methanol (35 mL) was stirred at room temperature. Solid 85% potassium hydroxide (225 mg) was added to adjust the pH to 6.5. After 1.5 hours, sodium cyanoborohydride (120 mg, 1.91 mmol, 1.5 eq) was added, and the mixture was stirred for an additional two hours. Solid 85% potassium hydroxide (640 mg) was added, the mixture was filtered, and the filtrate concentrated. To the residue was added dry dichloromethane (35 mL) and EDCI (1.10 g, 5.72 mmol, 1.5 eq) and the mixture was stirred at room temperature for 17 hours. Saturated aqueous sodium carbonate solution (100 mL) was added and the aqueous layer was extracted with dichloromethane (3 x 150 mL). The combined organic extracts were dried over sodium sulfate and concentrated to give **8B** (2.50 g) as a crude yellow oil used directly without further purification.

C. 4'-[(2-*n*-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,3-dimethyl-2-oxo-1-pyrrolidinyl)methyl][1,1'-biphenyl]-2-sulfonamide

Crude **8B** (2.50 g) was dissolved in trifluoroacetic acid (16 mL) and the resulting solution was stirred at room temperature for 7 hours. The resulting mixture was concentrated, and saturated aqueous sodium

bicarbonate solution was added to the residue. This mixture was extracted with ethyl acetate (3 x 50 mL), and the combined organic extracts were dried over sodium sulfate and concentrated. Silica gel chromatography of the residue (5% methanol in dichloromethane as eluant) gave **8C** (0.98 g, 45% from **8A**) as a yellow solid.


D. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,3-dimethyl-2-oxo-1-pyrrolidinyl)methyl]-N-(2-pyrazinyl) [1,1'-biphenyl]-2-sulfonamide

A solution of **8C** (110 mg, 0.19 mmol, 1.0 eq) in DMF (2.5 mL) was treated at room temperature with sodium hydride (60% dispersion in mineral oil, 17 mg, 0.43 mmol, 2.2 eq). After 10 minutes, 2-chloropyrazine (68 µl, 0.76 mmol, 4.0 eq) was added via syringe, and the mixture heated at 120 °C for 2 hours and 130 °C for 6 hours. The solvent was evaporated, and the residue partially purified by preparative reverse-phase HPLC. The fractions containing product were evaporated, and the residue partitioned between dichloromethane and 10% aqueous sodium dihydrogen phosphate. The aqueous phase was adjusted to pH 5 and extracted with two additional portions of dichloromethane. The combined organic extracts were dried over sodium sulfate, evaporated, and the residue chromatographed on silica gel (3% methanol in dichloromethane as eluant) to yield still impure product (70 mg). This material was subjected to ion-exchange chromatography to give the pure title compound (8 mg, 6%) as a white powder after lyophilization: ¹H NMR (CDCl₃, 400 MHz) δ 8.52 (s, 1H), 8.29 (dd, *J*=1 and 8 Hz, 1H), 8.16 (d, *J*=3 Hz, 1H), 7.97 (s, 1H), 7.60 (dt, *J*=1 and 7 Hz, 1H), 7.55 (dt, *J*=1 and 7 Hz, 1H), 7.20 (dd, *J*=1 and 7 Hz, 1H), 7.01 (s, 2H), 6.96 (s, 1H), 4.60 (AB quartet, *J*=16 Hz, 2 H), 4.15 (AB quartet, *J*=16 Hz, 2 H), 3.16 (m, 2H), 2.34 (dd, *J*=7 and 8 Hz, 2H), 1.97 (m, 6H), 1.87 (t, *J*=7 Hz, 2H), 1.82 (m, 2H), 1.61 (m, 2H), 1.36 (sextet, *J*=7 Hz, 2 H), 1.17 (s, 3H), 1.16 (s, 3H), 0.90 (t, *J*=7 Hz, 3H); LRMS *m/z* 643 (ESI+ mode).

Example 9

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3-chloro-2-pyrazinyl)-2'-[(3,3-dimethyl-2-oxo-1-pyrrolidinyl)methyl][1,1'-biphenyl]-2-sulfonamide

5




A solution of **8C** (100 mg, 0.18 mmol, 1.0 eq) in DMF (1.8 mL) was
 10 treated at room temperature with sodium hydride (60% dispersion in mineral oil, 8.5 mg, 0.21 mmol, 1.2 eq). After 10 minutes, 2,3-dichloropyrazine (79 mg, 0.53 mmol, 3.0 eq) was added via syringe, and the mixture heated at 60°C for 3 hours, then at 85°C for 14 hours, and then at 120°C for 5 hours. Workup and purification as described in
 15 Example 8 yielded the title compound (7.4 mg, 10%) as a white powder after lyophilization: ¹H NMR (CDCl₃, 400 MHz) δ 8.40 (dd, *J*=1 and 8 Hz, 1H), 8.0 (m, 2H), 7.68 (dt, *J*= 1 and 7 Hz, 1H), 7.61 (dt, *J*=1 and 7 Hz, 1H), 7.22 (dd, *J*=1 and 7 Hz, 1H), 7.16 (s, 1H), 7.05 (d, *J* =8 Hz, 1H), 6.89 (d, *J*=8 Hz, 1H), 6.85 (br s, 2H), 4.87 (AB quartet, *J*=16 Hz, 2 H), 4.31 (d, *J*=16
 20 Hz, 1H), 4.15 (d, *J*=16 Hz, 1H), 3.16 (m, 2H), 2.82 (t, *J*=7 Hz, 2H), 1.90-2.20 (m, 8H), 1.82 (m, 2H), 1.66 (m, 2H), 1.38 (sextet, *J*=7 Hz, 2H), 1.22 (s, 3H), 1.14 (s, 3H), 0.92 (t, *J*=7 Hz, 3H) ppm; LRMS *m/z* 677, 679 (ESI+ mode); *m/z* 675, 677 (ESI- mode).

Example 10

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-[(2-oxo-1-pyrrolidinyl)methyl][1,1'-biphenyl]-2-sulfonamide


5



The title compound was prepared in 51% yield starting from **3F** and 4-aminobutyric acid as described in Example 4 as a white solid: mp 95-
 10 100°C; ¹H NMR (CDCl₃) δ 0.90 (t, *J*=7.3 Hz, 3H), 1.36 (m, 2H), 1.61 (m, 2H), 1.82-2.10 (m, 15H), 2.17 (s, 3H), 2.36 (m, 2H), 3.41 (m, 2H), 4.20 (m, 2H), 4.72 (s, 2H), 7.00-7.90 (m, 7H).

Example 11

15 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,3-dimethyl-2-oxo-1-pyrrolidinyl)methyl]-N-(3,6-dimethyl-2-pyrazinyl)[1,1'-biphenyl]-2-sulfonamide




20

A mixture of **8C** (200 mg, 0.36 mmol, 1.0 eq), palladium acetate (4 mg, 0.02 mmol, 0.05 eq), (S)-BINAP (2.5 mg, 0.02 mmol, 0.05 eq), and toluene (5 mL) was sparged with nitrogen for 10 minutes. 2-Chloro-3,6-dimethylpyrazine (83 μ l, 0.71 mmol, 2.0 eq) was added via syringe,
 5 followed by sodium hydride (60% dispersion in mineral oil, 28 mg, 0.71 mmol, 2.0 eq). The mixture was heated at 80°C for 15 hours. After cooling, the mixture was diluted with ethyl acetate and partitioned against 10% aqueous sodium phosphate adjusted to pH 5-6, and the aqueous layer was extracted with two additional portions of ethyl acetate.
 10 The combined organic extracts were dried over sodium sulfate, concentrated, and evaporated to a yellow oil, which was chromatographed on silica gel (1:3 hexanes/ethyl acetate eluant) to yield the title compound (50 mg, 21%) as an off-white solid: ^1H NMR (CD_3OD , 400 MHz) δ 8.14 (d, \underline{J} =7 Hz, 1H), 7.75 (br s, 1H), 7.49 (m, 2H), 7.11 (dd, \underline{J} = 1 and 7 Hz, 1H),
 15 6.87 (s, 2H), 6.70 (s, 1H), 4.65 (br s, 2 H), 4.11 (s, 2H), 3.95 (m, 2H), 2.92 (m, 2H), 2.29 (br s, 2H), 2.17 (s, 3H), 2.06 (s, 3H), 1.79 (m, 6H), 1.68 (m, 2H), 1.65 (m, 2H), 1.41 (m, 2H), 1.17 (m, 2H), 1.03 (s, 3H), 1.00 (s, 3H), 0.72 (t, \underline{J} = 7 Hz, 3H); LRMS m/z 671 (ESI+ mode); m/z 669 (ESI- mode).

20

Example 12

N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl]methyl]-N,N',N'-trimethylurea




25

To **2I** (34.7 mg, 0.06 mmol) in CH₂Cl₂ (0.6 mL) and DMF (0.15 mL) was added dimethylcarbamyl chloride (6.5 mg, 0.06 mmol) followed by triethyl amine (6.7 mg, 0.066 mmol). The resulting mixture was stirred at RT for two days. The mixture was purified by anion-exchange chromatography to provide the title compound (90%). Characterization recorded with Example 73.

Example 13

N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N'-(1,1-dimethylethyl)-N-methylurea




To **2I** (34.7 mg, 0.06 mmol) in CH₂Cl₂ (0.6 mL) and DMF (0.15 mL), *t*-butylisocyanate (6.0 mg, 0.06 mmol) was added, and the resulting mixture was stirred at RT for two days. The mixture was purified by anion-exchange chromatography to provide the title compound (90%).

Characterization recorded with Example 74.

Example 14


[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]methylcarbamic acid ethyl ester



The title compound was prepared from **2I** and ethyl chloroformate
 5 using a procedure similar to the one described in Example 12 (90%).
 Characterization recorded with Example 75.

Example 15

10 [[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-
dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-
yl)methyl]methylcarbamic acid 2-methylpropyl ester




15 The title compound was prepared from **2I** and isobutyl
 chloroformate using a procedure similar to the one described in Example
 12 (90%). Characterization recorded with Example 76.

Example 16

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,3-dimethyl-2-oxo-1-pyrrolidinyl)methyl]-N-(3-methoxy-2-pyrazinyl)[1,1'-biphenyl]-2-sulfonamide

5




The title compound was prepared from **8C** (486 mg, 0.86 mmol) and 2-chloro-3-methoxypyrazine (250 mg, 1.72 mmol) [Uchimaru, F. et al., *Chem. Pharm. Bull.*, 20, 2204-2208 (1972)] using a procedure similar to the one used in Example 9. Preparative reverse-phase HPLC of the crude product after extractive workup gave the title compound (24 mg, 4%) as an off-white solid after lyophilization: ¹H NMR (CDCl₃, 400 MHz) δ 10.8 (br s, 2H), 8.36 (dd, *J*=1 and 8 Hz, 1H), 7.50-7.65 (m, 4H), 7.19 (dd, *J*=1 and 7 Hz, 1H), 7.15 (s, 1H), 7.01 (s, 2H), 4.92 (AB quartet, *J*=16 Hz, 2H), 4.20 (AB quartet, *J*=16 Hz, 2H), 3.99 (s, 3H), 3.18 (m, 2H), 2.82 (m, 2H), 2.18 (m, 4H), 1.96 (m, 4H), 1.82 (m, 2H), 1.63 (m, 2H), 1.32 (m, 2H), 1.21 (s, 3H), 1.12 (s, 3H), 0.87 (t, 3H, *J*=7 Hz, 3H); LRMS *m/z* 673 (ESI+ mode); *m/z* 671 (ESI- mode); HPLC retention time 25.52 minutes (Method B).

20

Example 17

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-formyl-N-(4,5-dimethyl-3-isoxazolyl)-[1,1'-biphenyl]-2-sulfonamide



- A. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-formyl-N-(4,5-dimethyl-3-isoxazolyl)-N-(2-methoxyethoxymethyl)[1,1'-biphenyl]-2-sulfonamide

Palladium catalyzed Suzuki coupling of **5E** and [2-[[[(4,5-dimethyl-3-isoxazolyl)(2-methoxyethoxy)methyl]amino]sulfonyl]phenyl]boronic acid was performed according to General Method 1 to afford **17A** (81%) following silica-gel chromatography.


- B. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-formyl-N-(4,5-dimethyl-3-isoxazolyl)-[1,1'-biphenyl]-2-sulfonamide

Treatment of **17A** with 6N aqueous hydrochloric acid according to General Method 7 provided the title compound (85%): $R_f = 0.38$, 5% MeOH in methylene chloride.

Example 18


4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-
dimethyl-3-isoxazolyl)-2'-[(2-oxo-1-pyrrolidinyl)methyl][1,1'-
biphenyl]-2-sulfonamide

5



The title compound was obtained as a white solid using **17B** and 4-aminobutyric acid as described in Example 4 (25%): mp 97-103°C; ¹H NMR (CDCl₃) δ 0.90 (t, *J*=7 Hz, 3H), 1.36 (m, 2H), 1.62 (m, 2H), 1.80-2.10 (m, 13H), 2.26 (s, 3H), 2.39 (m, 4H), 3.31 (m, 2H), 4.20 (s, 2H), 4.73 (s, 2H), 7.05-8.10 (m, 7H).

(S)-N-[[2'-[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-
yl)methyl]-N-[2-methyl-1-(3-methyl-1,2,4-oxadiazol-5-
yl)propyl]pentanamide



20

A. (S)-5-(1-amino-2-methylpropyl)-3-methyl-1,2,4-oxadiazole

A solution of BOC-L-valine (4.34 g, 20.0 mmol) in dichloromethane (20 mL) was cooled to 0°C and treated dropwise with a solution of dicyclohexylcarbodiimide (2.06 g, 10.0 mmol) in dichloromethane (5 mL).

- 5 After 1 hour the white solid which had formed was removed by filtration and the filtrate concentrated under reduced pressure. The residue was dissolved in pyridine (15 mL) and treated with acetamidoxime (488 mg, 6.6 mmol) in pyridine (5 mL) and the mixture heated at reflux for 1 hour. The pyridine was evaporated under reduced pressure and the residue
- 10 partitioned between ethyl acetate and water. The organic phase was washed several times with water, dried over magnesium sulfate and concentrated. The residue was chromatographed on silica gel using 10% ethyl acetate in hexanes to elute the product (0.62 g, 25%) as a white solid.

- A solution of the above product in 3 M hydrochloric acid (10 mL)
- 15 was stirred at room temperature for 1 hour. The mixture was evaporated to dryness under reduced pressure and then azeotroped with toluene. This gave **19A** as a white solid (100%).

B. 2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-carboxaldehyde

P17 (0.35 g, 0.79 mmol) was subjected to deprotection according to General Method 7. Extraction and concentration gave 200 mg (71%) of **19B** as a yellow solid which was used without purification.

C. (S)-[[1-[2'-[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-ylmethylamino]-2-methylpropyl]-3-methyl-1,2,4-oxadiazole

- A mixture of **19B** (100 mg, 0.28 mmol) and **19A** (53 mg, 0.28 mmol) in dichloroethane (8 mL) was treated with sodium triacetoxyborohydride (74 mg, 0.35 mmol) in one portion and the mixture stirred under argon at
- 30 room temperature. After 1 hour more sodium triacetoxyborohydride (32 mg, 0.15 mmol) was added to the mixture which stirred for 45 minutes

more and the mixture was adjusted to pH 6 with saturated sodium bicarbonate solution. The mixture was diluted with 50 mL dichloromethane and the organic phase washed with water, dried over magnesium sulfate and concentrated to give crude **19C** (110 mg, 79%)


5 which was used without purification in the next step.

D. (S)-N-[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-[2-methyl-1-(3-methyl-1,2,4-oxadiazol-5-yl)propyl]pentanamide

10 A mixture of **19C** (100 mg, 0.20 mmol), dichloromethane (5 mL) and triethylamine (70 mL, 0.5 mmol) was treated with valeryl chloride (60 mg, 0.5 mmol) and the mixture stirred under argon for 16 hours. The reaction mixture was concentrated under reduced pressure and the residue treated with 2 M sodium hydroxide (1 mL) and 2-propanol (2 mL) and stirred at
15 room temperature for 1.5 hours. Citric acid was added to adjust the pH to 5 - 6 and the mixture extracted with dichloromethane. The residue obtained after drying the extract over magnesium sulfate and concentrating under reduced pressure was chromatographed on silica gel using 30 to 50% ethyl acetate in hexanes to elute the title product (73 mg,
20 63%) as a white foam: ESIMS (NH₃) m/z 580 (MH⁺ 100), 597 (M + NH₄⁺, 20), 1176 (2M + NH₄⁺, 10); HRMS calcd for C₃₀H₃₇N₅O₅S (MH⁺) 580.2593, found 580.2619.

Example 20

25 N-(3,4-Dimethyl-5-isoxazolyl)-2'-[[[(3,3-dimethyl-2-oxo-1-pyrrolidinyl)methyl]-4'-[(2-ethyl-5,7-dimethyl-3H-imidazo[4,5-b]pyridin-3-yl)methyl][1,1'-biphenyl]-2-sulfonamide



A. N-(3,4-Dimethyl-5-isoxazolyl)-2'-[(3,3-dimethyl-2-oxo-1-pyrrolidinyl)methyl]-4'-hydroxymethyl-N-(2-methoxyethoxymethyl)[1,1'-biphenyl]-2-sulfonamide

Preparation **P21** (4.25 g, 9.0 mmol) was reacted with ethyl 4-amino-2,2-dimethylbutanoate hydrochloride (2.13 g, 10.9 mmol) according to General Method 5. The crude residue was chromatographed on silica gel (5% methanol in dichloromethane eluant) to yield 3.05 g of **20A** (59%) as an orange oil.

B. N-(3,4-Dimethyl-5-isoxazolyl)-2'-[(3,3-dimethyl-2-oxo-1-pyrrolidinyl)methyl]-4'-[(2-ethyl-5,7-dimethyl-3H-imidazo[4,5-b]pyridine-3-yl)methyl]-N-(2-methoxyethoxymethyl)[1,1'-biphenyl]-2-sulfonamide

A solution of **20A** (500 mg, 0.87 mmol, 1.0 eq), triphenylphosphine (344 mg, 1.3 mmol, 1.5 eq), and 2-ethyl-5,7-dimethyl-3H-imidazo[4,5-b]pyridine (184 mg, 1.1 mmol, 1.2 eq) in tetrahydrofuran (5 mL) was treated at 0°C with diethylazodicarboxylate (206 µl, 1.3 mmol, 1.5 eq).


The mixture was allowed to warm to room temperature, was stirred for 16 hr, and then concentrated. The residue was chromatographed on silica gel (3:2 hexanes/acetone as eluant) to give 320 mg of a mixture containing **20B** and 2-ethyl-5,7-dimethyl-3H-imidazo[4,5-b]pyridine (approximately 3:1 ratio by weight).

C. N-(3,4-Dimethyl-5-isoxazolyl)-2'-[(3,3-dimethyl-2-oxo-1-pyrrolidinyl)methyl]-4'-[(2-ethyl-5,7-dimethyl-3H-imidazo[4,5-b]pyridine-3-yl)methyl][1,1'-biphenyl]-2-sulfonamide

20B (320 mg) was deprotected according to General Method 8 (ethanol). The crude residue was purified by reverse-phase preparative HPLC followed by extraction (3x50 mL ethyl acetate) of the product from brine adjusted to pH 4 with hydrochloric acid, to provide 135 mg of the title compound (24% from **20A**) as a white solid after lyophilization; mp 95-104 °C; MS m/e 641 (ESI+ mode); MS m/e 639 (ESI- mode); HPLC retention time 3.43 minutes (Method C).

Example 21

N-(3,4-Dimethyl-5-isoxazolyl)-4'-[2-(2-methoxyethyl)-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl][1,1'-biphenyl]-2-sulfonamide



A. Methyl 1-[(3-methoxy-1-oxopropyl)amino]cyclopentane-1-carboxylate

3-Methoxypropanoic acid (1.65 mL, 17.6 mmol, 2.1 eq) was added to a mixture of EDCI (1.77 g, 9.24 mmol, 1.1 eq), triethylamine (3.5 mL, 25.2 mmol, 3.0 eq), 4-dimethylaminopyridine (20 mg, 0.18 mmol, 0.02 eq), and dichloromethane (100 mL) at room temperature. After 5 minutes, methyl 1-aminocyclopentane-1-carboxylate hydrochloride (1.50 g, 8.4 mmol, 1.0 eq) was added and the mixture was stirred at room temperature for 18 hours. The reaction mixture was partitioned against 1N hydrochloric acid

and the aqueous phase was extracted once with dichloromethane. The combined organic extracts were dried over sodium sulfate and concentrated. Silica gel chromatography of the residue (1:3 hexanes/ethyl acetate as eluant) yielded 1.30 g of **21A** (67%) as a colorless oil.

5

B. 1-[(3-Methoxy-1-oxopropyl)aminol]cyclopentane-1-carboxylic acid

A mixture of **21A** (1.25 g, 5.5 mmol, 1.0 eq), lithium hydroxide hydrate (300 mg, 7.1 mmol, 1.3 eq), THF (10 mL), and water (10 mL) was stirred at room temperature for 5.5 hours. 1N hydrochloric acid (10 mL)
10 was added and the mixture was saturated with solid sodium chloride, then extracted with ethyl acetate (3x30 mL). The combined organic extracts were dried over sodium sulfate and concentrated to provide 0.95 g of **21B** (80%) as a white solid.

15 C. 1-[(3-Methoxy-1-oxopropyl)aminol]cyclopentane-1-carboxamide

A suspension of **21B** (0.95 g, 4.4 mmol, 1.0 eq) in THF (20 mL) was treated with 1,1'-carbonyldiimidazole (930 mg, 5.7 mmol, 1.3 eq) at room temperature. After 30 minutes, the mixture was cooled to -78 °C and ammonia gas was introduced. The resulting heterogeneous mixture was
20 allowed to warm to room temperature and was stirred for 16 hours. The solvent was evaporated and 1N hydrochloric acid saturated with sodium chloride was added to the residue. The aqueous mixture was extracted with ethyl acetate (6x50 mL), and the combined organic layers were dried over sodium sulfate and concentrated to yield 500 mg of **21C** (53%) as a
25 white solid.

D. 2-(2-Methoxyethyl)-1,3-diazaspiro[4.4]non-1-en-4-one

A solution of **21C** (460 mg, 2.15 mmol, 1.0 eq), potassium hydroxide (288 mg, 4.30 mmol, 2.0 eq) and methanol (15 mL) was heated at reflux for
30 20 hours. The mixture was cooled, solid ammonium chloride was added, and the solvent was evaporated. Water was added and the mixture was

extracted with ethyl acetate (3x30 mL). The combined organic extracts were dried over sodium sulfate, concentrated, and the residue chromatographed on silica gel (4% methanol in dichloromethane as eluant) to yield 162 mg of **21D** (38%) as an amber oil.

5

E. N-(3,4-Dimethyl-5-isoxazolyl)-4'-[2-(2-methoxyethyl)-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl][1,1'-biphenyl]-2-sulfonamide


21D (75 mg, 0.38 mmol, 1.0 eq) was alkylated with **P18** (195 mg, 0.38 mmol, 1.0 eq) according to General Method 4. The resulting crude orange oil was dissolved in methanol (7 mL). Concentrated hydrochloric acid (7 mL) was added and the solution was heated at 55°C for 14 hours. The reaction mixture was concentrated and then partitioned between ethyl acetate and pH 5 sodium phosphate buffer. The organic layer was dried over sodium sulfate and concentrated to provide a crude residue.

15 The title compound (7 mg, 3% yield) was obtained as a white powder following silica gel chromatography (4% methanol in chloroform as eluant), preparative reverse-phase HPLC, and lyophilization: MS m/e 537 (ESI+ mode); MS m/e 535 (ESI- mode); HPLC retention time 3.35 minutes (Method A).

20

Example 22

N-(3,4-Dimethyl-5-isoxazolyl)-4'-[2-(ethoxymethyl)-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl][1,1'-biphenyl]-2-sulfonamide



25

A. Methyl 1-[(2-ethoxyethanoyl)amino]cyclopentane-1-carboxylate

2-Ethoxyacetic acid (3.3 mL, 35 mmol, 2.1 eq) was added to a mixture of EDCI (3.38 g, 18 mmol, 1.1 eq), triethylamine (7.0 mL, 50 mmol, 3.0 eq), 4-dimethylaminopyridine (20 mg, 0.18 mmol, 0.01 eq), and
5 dichloromethane (100 mL) at room temperature. After 5 minutes, methyl 1-aminocyclopentane-1-carboxylate hydrochloride (3.0 g, 17 mmol, 1.0 eq) was added and the mixture was stirred at room temperature for 18 hours. The solvent was evaporated and the residue partitioned between ether and 1N hydrochloric acid. The organic layer was washed once with 1N
10 hydrochloric acid, then twice with saturated aqueous sodium bicarbonate, and once with brine. The organic layer was dried over magnesium sulfate and concentrated to give 2.0 g of **22A** (52%) as a pink oil.

B. 2-(Ethoxymethyl)-1,3-diazaspiro[4.4]non-1-en-4-one


15 A mixture of **22A** (2.0 g, 8.7 mmol, 1.0 eq), lithium hydroxide hydrate (440 mg, 10.5 mmol, 1.2 eq), THF (10 mL), and water (3 mL) was stirred at room temperature for 16 hours. 2N hydrochloric acid (6 mL) was added and the solvents were evaporated. The residue was dried azeotropically with toluene, then dissolved in THF (15 mL) and treated
20 with 1,1'-carbonyldiimidazole (2.8 g, 17.3 mmol, 2.1 eq) at room temperature. After 2 hours, the mixture was cooled to -78 °C and ammonia gas was introduced. The resulting mixture was allowed to warm to room temperature and was stirred for 16 hours, after which the solvent was evaporated. The residue was suspended in methanol (15 mL), solid
25 potassium hydroxide (2.9 g, 43 mmol, 5 eq) was added, and the mixture was heated at reflux for 72 hours. After cooling, the mixture was treated with solid ammonium chloride and the solvent was evaporated. Water was added and the mixture was extracted with ethyl acetate (3x30 mL). The combined organic extracts were dried over sodium sulfate,
30 concentrated, and the residue chromatographed on silica gel (4% methanol in chloroform as eluant) to yield 370 mg of **22B** (22%) as an orange oil.

C. N-(3,4-Dimethyl-5-isoxazolyl)-4'-[2-(ethoxymethyl)-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl][1,1'-biphenyl]-2-sulfonamide

The title compound was prepared from **22B** (80 mg, 0.41 mmol) and **P18** (209 mg, 0.41 mmol) according to the procedure described in Example 61, Step E, substituting ethanol for methanol in the deprotection reaction. The crude residue was chromatographed twice on silica gel (4% methanol in chloroform as eluant, followed by 25:75:1 hexanes/ethyl acetate/ acetic acid as eluant), then was further purified by reverse-phase preparative HPLC to provide 42 mg of the title compound (19%) as a white solid after lyophilization; MS m/e 537 (ESI+ mode); MS m/e 535 (ESI- mode); HPLC retention time 3.51 minutes (Method A).

Example 23

15 4'-[(2-Ethyl-5,7-dimethyl-3H-imidazo[4,5-b]pyridin-3-yl)methyl]-2'-[(2-oxo-1-pyrrolidinyl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)[1,1'-biphenyl]-2-sulfonamide



20

A. 2'-Formyl-N-(3,4-dimethyl-5-isoxazolyl)-4'-[(methanesulfonyl)oxy]-N-(2-methoxyethoxymethyl)[1,1'-biphenyl]-2-sulfonamide

25

P21 (2.4 g) was converted to the corresponding mesylate according to General Method 3. The crude product (2.7 g) was carried on without further purification.

B. 4'-[(2-Ethyl-5,7-dimethyl-3H-imidazo[4,5-b]pyridin-3-yl)methyl]-2'-formyl-N-(3,4-dimethyl-5-isoxazolyl)-N-(2-methoxyethoxymethyl)[1,1'-biphenyl]-2-sulfonamide

5 **23A** (2.7 g) was used to alkylate 2-ethyl-5,7-dimethyl-3H-imidazo[4,5-b]pyridine according to General Method 4. The crude product was chromatographed on silica gel using 1:3 hexanes/ethyl acetate as eluant to provide 1.8 g **23B** as a colorless oil.

10 C. 4'-[(2-Ethyl-5,7-dimethyl-3H-imidazo[4,5-b]pyridin-3-yl)methyl]-2'-[(2-oxo-1-pyrrolidinyl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-N-(2-methoxyethoxymethyl)[1,1'-biphenyl]-2-sulfonamide


15 **23B** (1.5 g) was subjected to reductive amination with ethyl 4-aminobutanoate hydrochloride according to General Method 5. The reaction mixture was allowed to stir for 24 h to allow time for cyclization of the amino ester to the corresponding lactam. The crude product after workup was purified by silica gel chromatography (1:1 hexanes/acetone eluant) to provide **23C** (0.50 g) as a yellow oil.

20 D. 4'-[(2-Ethyl-5,7-dimethyl-3H-imidazo[4,5-b]pyridin-3-yl)methyl]-2'-[(2-oxo-1-pyrrolidinyl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)[1,1'-biphenyl]-2-sulfonamide

25 **23C** (0.50 g) was subjected to sulfonamide deprotection using HCl/dioxane/ethanol according to General Method 8. The crude product was purified by silica gel chromatography (90:9:1 dichloromethane/methanol/ammonium hydroxide eluant), and the purified product was partitioned between ethyl acetate and pH 5 potassium phosphate buffer. The ethyl acetate layer was dried over sodium sulfate and concentrated to provide 310 mg of the title compound as a white solid;
30 mp 98-102 °C; MS m/e 613 (ESI+ mode); HPLC retention time 3.09 min (Method A); HPLC purity 97%.

Example 24

4'-[(2-Ethyl-5,7-dimethyl-3H-imidazo[4,5-b]pyridin-3-yl)methyl]-2'-[(3-methyl-2-oxo-1-imidazolidinyl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)[1,1'-biphenyl]-2-sulfonamide



A. 4'-[(2-Ethyl-5,7-dimethyl-3H-imidazo[4,5-b]pyridin-3-yl)methyl]-2'-[(3-methyl-2-oxo-1-imidazolidinyl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-N-(2-methoxyethoxymethyl)[1,1'-biphenyl]-2-sulfonamide


23B (2.0 g) was subjected to reductive amination with *N*-methylethylenediamine according to General Method 5. The crude product following extractive workup was dissolved in dichloromethane (25 ml) and treated with CDI (0.77 g). The mixture was stirred at RT for 24 h, and was then washed once with water and once with brine. The dichloromethane layer was dried over sodium sulfate and concentrated. The residue was chromatographed on silica gel using 95:5 chloroform/methanol as eluant to give **24A** (0.53 g) as a slightly yellow oil.

B. 4'-[(2-Ethyl-5,7-dimethyl-3H-imidazo[4,5-b]pyridin-3-yl)methyl]-2'-[(3-methyl-2-oxo-1-imidazolidinyl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)[1,1'-biphenyl]-2-sulfonamide

24A (0.50 g) was subjected to sulfonamide deprotection according to General Method 8. The crude product was purified by reverse-phase preparative HPLC to yield 140 mg of the title compound as a white solid following lyophilization; MS m/e 628 (ESI+ mode); HPLC retention time 3.03 min (Method A); HPLC purity 97%.

Example 25

(S)-2-[N-[2'-[[N-(3-Methyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-(1-oxopentyl)amino]-3,N-dimethylbutanamide



15 A. N-(3-Methyl-5-isoxazolyl)-2-bromobenzenesulfonamide

5-Amino-3-methylisoxazole (3.84 g) was added at RT in portions to a solution of 2-bromobenzenesulfonyl chloride (10.0 g) in pyridine (40 ml). The mixture was heated at 60 °C for 16 h, then the solvent was evaporated. The residue was taken up in ethyl acetate and washed three times with 1 N hydrochloric acid. The ethyl acetate layer was dried over sodium sulfate and concentrated to give **25A** (8.8 g).

B. N-[(2-Trimethylsilyl)ethoxymethyl]-N-(3-methyl-5-isoxazolyl)-2-bromobenzenesulfonamide

25 2-(Trimethylsilyl)ethoxymethyl chloride (5.2 ml) was added to a mixture of **25A** (8.8 g), potassium carbonate (7.7 g), and DMF (40 ml) at 0 °C. The mixture was allowed to warm to RT and was then stirred for 16 h. The solvent was evaporated, and the residue was taken up in ethyl acetate

and washed with water and brine. The ethyl acetate layer was dried over sodium sulfate and concentrated. The residue was purified by silica gel chromatography using 3:2 hexanes/ethyl acetate as eluant to provide **25B** (6.6 g) as an oil.

5

C. 4'-Formyl-N-(3-methyl-5-isoxazolyl)-N-[(2-trimethylsilyl)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

25B (2.0 g) and 4-formylphenylboronic acid (1.5 g) were subjected to Suzuki coupling according to General Method 1. The crude product was chromatographed on silica gel using 9:1 hexanes/ethyl acetate as eluant to yield 0.75 g **25C** as a colorless oil.

10

D. (S)-2-[N-[2'-[[N-(3-Methyl-5-isoxazolyl)-N-[(2-trimethylsilyl)ethoxymethyl]amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-amino]-3,N-dimethylbutanamide

15

25C (0.75 g) was subjected to reductive amination with L-valine N-methyl amide hydrochloride according to General Method 5. Crude **25D** (0.93 g) was obtained as an orange oil.

E. (S)-2-[N-[[2'-[[N-(3-Methyl-5-isoxazolyl)-N-[(2-trimethylsilyl)ethoxymethyl]amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-(1-oxopentyl)amino]-3,N-dimethylbutanamide

20

25D (0.93 g) was subjected to acylation with valeryl chloride according to General Method 6. The crude product was chromatographed on silica gel using 2:3 hexanes/ethyl acetate as eluant to yield 0.78 g **25E** as a colorless oil.

25


F. (S)-2-[N-[2'-[[N-(3-Methyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-(1-oxopentyl)amino]-3,N-dimethylbutanamide

30

25E (0.78 g) was deprotected with HCl/methanol according to General Method 8. The crude product was purified by silica gel chromatography using 3:7 hexanes/ethyl acetate as eluant, providing the title compound (210 mg) as a white solid; MS m/e 541 (ESI+ mode); HPLC retention time 31.32 min (Method B); HPLC purity >98%.

Example 26

(S)-2-[N-[2'-[N-(4-Bromo-3-methyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-(1-oxopentyl)amino]-3,N-dimethylbutanamide




15

A solution of **25** (75 mg) in chloroform (1.5 ml) was treated with NBS (25 mg) at RT. After 1h, the mixture was diluted with dichloromethane and partitioned against water. The organic layer was dried over sodium sulfate and concentrated, and the residue was chromatographed on silica gel using 3:7 hexanes/ethyl acetate as eluant to provide the title compound (25 mg) as a white solid; MS m/e 619, 621 (ESI+ mode); HPLC retention time 31.83 min (Method B); HPLC purity >99%.

25

Example 27

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-propyl[1,1'-biphenyl]-2-sulfonamide



5 A. 4-Bromo-3-(1-propen-1-yl)benzonitrile

n-Butyllithium (2.5M solution in hexane, 7.6 ml, 19 mmol) was added dropwise to a solution of ethyltriphenylphosphonium bromide (6.42 g, 17.3 mmol) in 100 ml of 1:1 THF/ether at -15 °C. The mixture was stirred for 3h at RT and then was cooled to -50 °C. **2A** (4.0 g, 19.0 mmol) in THF (10 ml) was added and the mixture was allowed to warm to RT and was stirred for 16 h. The mixture was added to water and extracted with EtOAc (3 x 50 mL) and the combined organic extracts were washed with water, dried over magnesium sulfate, and evaporated. The residue was chromatographed on silica gel using 95:5 hexane/EtOAc to afford **27A** as an E/Z mixture (3.5 g, 83%).

B. 4-Bromo-3-propylbenzonitrile

A mixture of **27A** (1.5 g) and 150 mg of PtO₂ in 40 ml EtOH was hydrogenated at 35 PSI for 40 min. Filtration and concentration gave 1.44 g of **27B** (85%).

C. 4-Bromo-3-propylbenzaldehyde

27B (1.44 g) was treated with DIBAL-H according to General Method 14 to provide crude **27C** (1.4 g, 97%) as an oil.

D. N-(3,4-Dimethyl-5-isoxazolyl)-4'-formyl-2'-propyl-N-[(2-methoxyethoxy)methyl][1,1'-biphenyl]-2-sulfonamide

27C (1.4 g) was subjected to Suzuki coupling according to General Method 1 to provide **27D** (27%) as an oil.

E. N-(3,4-Dimethyl-5-isoxazolyl)-4'-hydroxymethyl-2'-propyl-N-[(2-methoxyethoxy)methyl][1,1'-biphenyl]-2-sulfonamide

27D (810 mg) was reduced with sodium borohydride in methanol according to General Method 11 to provide **27E** (32%) as an oil.

F. N-(3,4-Dimethyl-5-isoxazolyl)-4'-(methanesulfonyloxymethyl-2'-propyl-N-[(2-methoxyethoxy)methyl][1,1'-biphenyl]-2-sulfonamide


27E (250 mg) was converted to the corresponding methanesulfonate ester according to General Method 3 to provide **27F** (68%) as an oil.

G. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-propyl-N-[(2-methoxyethoxy)methyl]-N-(3,4-dimethyl-5-isoxazolyl)[1,1'-biphenyl]-2-sulfonamide

27F (100 mg) was used to alkylate 2-butyl-1,3-diazaspiro[4.4]non-1-en-4-one according to General Method 4. **27G** (100 mg, 85%) was produced as an oil.

H. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-propyl-N-(3,4-dimethyl-5-isoxazolyl)[1,1'-biphenyl]-2-sulfonamide

27G (100 mg) was deprotected according to General Method 7. The crude product was purified by preparative HPLC to provide the title compound (57 mg, 66%) as a solid; MS m/e 577 (ESI+ mode); HPLC retention time 29.11 min (Method B); HPLC purity >98%.

Example 284'-[(7-Methoxycarbonyl-2-ethoxybenzimidazol-1-yl)methyl]-
N-(3,4-dimethyl-5-isoxazolyl)-[1,1'-biphenyl]-2-sulfonamide

5

A. N-(2-Methoxycarbonyl-6-nitrophenyl)-4-bromobenzylamine

Triethylamine (5.2 ml, 37 mmol) was added to a mixture of methyl
 10 2-chloro-3-nitrobenzoate (3.2 g, 15 mmol) and 4-bromobenzylamine
 hydrochloride (3.4 g, 15 mmol) in acetonitrile (75 ml). The mixture was
 heated at reflux for 48 hr, then was cooled and concentrated. 10%
 Aqueous sodium dihydrogen phosphate solution was added and the
 mixture was extracted with 2 portions of ethyl acetate. The combined
 15 organic extracts were dried over sodium sulfate and concentrated, and the
 residue was crystallized from ethyl acetate to give **28A** (3.2 g) as a yellow
 solid.

B. 4'-[(2-Methoxycarbonyl-6-nitrophenyl)aminomethyl]-N-(3,4-
 20 dimethyl-5-isoxazolyl)-N-(2-trimethylsiloxyethyl)[1,1'-biphenyl]-2-
 sulfonamide

28A (3.5 g, 9.6 mmol) was subjected to Suzuki coupling according to
 General Method 1, providing **28B** as a yellow oil (3.1 g) following silica gel
 chromatography using 5:1 hexanes/ethyl acetate as eluant.

25

C. 4'-[(2-Methoxycarbonyl-6-aminophenyl)aminomethyl]-N-(3,4-
 dimethyl-5-isoxazolyl)-N-(2-trimethylsiloxyethyl)[1,1'-biphenyl]-2-
 sulfonamide

28B (2.4 g, 3.6 mmol) was treated with tin (II) chloride dihydrate (3.3 g) in ethyl acetate (80 ml) according to General Method 18. The crude product was purified by silica gel chromatography using 2:1 hexanes/ethyl acetate as eluant to provide **28C** (1.3 g) as a pale yellow oil.

5

D. 4'-[(7-Methoxycarbonyl-2-ethoxybenzimidazol-1-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-N-(2-trimethylsiloxymethyl)[1,1'-biphenyl]-2-sulfonamide

10 **28C** (1.3 g), tetraethylorthocarbonate (6 ml), and acetic acid (0.2 ml) was heated under a nitrogen atmosphere at 70 °C for 2 h. The mixture was cooled and concentrated, and the residue was chromatographed on silica gel using 1:1 hexanes/ethyl acetate as eluant to provide **28D** (1.1 g) as a brown oil.

15


E. 4'-[(7-Methoxycarbonyl-2-ethoxybenzimidazol-1-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-[1,1'-biphenyl]-2-sulfonamide

28D (1.1 g) was subjected to sulfonamide deprotection using TBAF
20 in THF according to General Method 10. The crude product was purified by silica gel chromatography using 2:1 hexanes/acetone as eluant to provide 0.75 g of the title compound as a white solid; mp 105-110 °C; MS m/e 561 (ESI+ mode); HPLC retention time 3.96 min (Method A); HPLC purity 96%.

25

Example 29

4'-[(7-Carboxy-2-ethoxybenzimidazol-1-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-[1,1'-biphenyl]-2-sulfonamide




28 (0.70 g) was subjected to ester hydrolysis according to General Method 15 to provide the crude product (0.66 g). Purification of a portion by reverse-phase preparative HPLC provided the title compound (9 mg, white solid); MS m/e 547 (ESI+ mode); HPLC retention time 3.79 min (Method A); HPLC purity 91%.

10

Example 30

4'-[(7-Methoxycarbonyl-2-ethylbenzimidazol-1-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-[1,1'-biphenyl]-2-sulfonamide



15

A solution of **28C** (3.8 g, 5.9 mmol) and triethylamine (1.7 ml, 12 mmol) in dichloromethane (25 ml) was treated at 0 °C with propionyl chloride (0.67 ml, 10 mmol), and the mixture was allowed to come to RT. After 2.5 h, aqueous sodium bicarbonate solution was added to the mixture and the aqueous layer was extracted with two portions of dichloromethane. The combined organic extracts were dried over sodium sulfate and concentrated to provide an orange oil.

To this residue was added methanolic hydrogen chloride (prepared from 100 ml methanol and 11 ml (200 mmol) acetyl chloride), and the


resulting solution was heated at 50 °C for 16 h. The mixture was cooled and concentrated, and the residue was extracted with two portions of ethyl acetate from sodium phosphate buffer adjusted to pH 4. The combined ethyl acetate extracts were dried over sodium sulfate and concentrated.

- 5 The residue was chromatographed on silica gel using 1:2 hexanes/acetone as eluant to provide the title compound (2.6 g) as a slightly orange solid; MS m/e 545 (ESI+ mode); HPLC retention time 3.32 min (Method C); HPLC purity 95%.

10

Example 31

4'-[(7-Carboxy-2-ethylbenzimidazol-1-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-[1,1'-biphenyl]-2-sulfonamide




15

- 30 (2.6 g, 4.8 mmol) was subjected to ester hydrolysis according to General Method 15. The THF was evaporated and the residue was treated with 6 ml of 2N hydrochloric acid, resulting in the precipitation of a white solid. The solid was collected on a filter, rinsed with water, and dried to provide 2.4 g of the title compound; MS m/e 531 (ESI+ mode); HPLC retention time 2.94 min (Method A); HPLC purity 95%.

25

Example 32

2'-[(3,3-Dimethyl-2-oxopyrrolidin-1-yl)methyl]-4'-[(2-ethoxy-7-(methoxycarbonyl)benzimidazol-1-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-[1,1'-biphenyl]-2-sulfonamide



A. 2-[(5'-Aminomethyl-2'-bromo)phenyl]-1,3-dioxolane

5 Borane•THF (100 ml of a 1.0 M solution in THF, 100 mmol) was added at 0 °C to a solution of 2-[(2'-Bromo-5'-cyano)phenyl]-1,3-dioxolane [Zhang, H.-Y. et al., *Tetrahedron*, **50**, 11339-11362 (1994)] (10.8 g, 43 mmol) in 25 ml THF. The mixture was allowed to warm to RT and was stirred for 18 h. After cooling to 0 °C, the mixture was treated carefully
10 with 10 ml methanol and was then evaporated. The residue was taken up in 150 ml ethyl acetate and was washed with 1N aqueous sodium hydroxide, followed by water and brine. The organic layer was dried over magnesium sulfate and concentrated to give crude **32A** (11.0 g) as an amber oil, which was estimated (HPLC) to be 80% pure.

15

B. N-(2-Methoxycarbonyl-6-nitrophenyl)-4-bromo-3-(1,3-dioxolan-2-yl)benzylamine

Triethylamine (4.8 ml, 34 mmol) was added to a mixture of methyl
20 2-chloro-3-nitrobenzoate (4.9 g, 23 mmol) and **32A** (5.9 g of a 65% pure mixture, 15 mmol) in acetonitrile (150 ml). The mixture was heated at reflux for 24 h and was then cooled and concentrated. Ethyl acetate was added and the solution was washed twice with 10% aqueous potassium dihydrogen phosphate solution and once with aqueous sodium bicarbonate
25 solution. The organic layer was concentrated and the residue purified by silica gel column chromatography using 3:1 hexanes/ethyl acetate as eluant, followed by trituration with 3:1 hexanes/ethyl acetate. **32B** was a yellow solid (6.6 g).

C. N-(2-Methoxycarbonyl-6-nitrophenyl)-4-bromo-3-(formyl)benzylamine

5 **32B** (6.6 g, 15 mmol) was subjected to acetal hydrolysis according to General Method 19, providing **32C** as a crude yellow solid following extractive workup.

10 D. 2'-Formyl-4'-[(2-methoxycarbonyl-6-nitrophenyl)aminomethyl]-N-(3,4-dimethyl-5-isoxazolyl)-N-(2-trimethylsiloxymethyl)[1,1'-biphenyl]-2-sulfonamide

15 Crude **32C** was subjected to Suzuki coupling according to General Method 1, providing **32D** (5.9 g) as a yellow oil following silica gel chromatography using 2:1 hexanes/ethyl acetate as eluant.

20 E. 2'-[(3,3-Dimethyl-2-oxopyrrolidin-1-yl)methyl]-4'-[(2-methoxycarbonyl-6-nitrophenyl)aminomethyl]-N-(3,4-dimethyl-5-isoxazolyl)-N-(2-trimethylsiloxymethyl)[1,1'-biphenyl]-2-sulfonamide

25 **32D** (4.3 g) was subjected to reductive amination with ethyl 4-amino-2, 2-dimethylbutanoate hydrochloride according to General Method 5. The reaction mixture was allowed to stir for 60 h at RT to allow time for cyclization of the amino ester to the corresponding lactam. The crude product after workup was purified by silica gel chromatography, using 3:2 hexanes/ethyl acetate as eluant, to provide **32E** (2.8 g) as a yellow oil.

30 F. 2'-[(3,3-Dimethyl-2-oxopyrrolidin-1-yl)methyl]-4'-[(2-methoxycarbonyl-6-aminophenyl)aminomethyl]-N-(3,4-dimethyl-5-

isoxazolyl)-N-(2-trimethylsiloxymethyl)[1,1'-biphenyl]-2-sulfonamide

5 **32E** (2.8 g, 3.6 mmol) was treated with tin (II) chloride dihydrate (3.2 g) in ethyl acetate (200 ml) according to General Method 18. The crude product (3.5 g) was used without further purification.

G. 2'-[(3,3-Dimethyl-2-oxopyrrolidin-1-yl)methyl]-4'-[(7-methoxycarbonyl)benzimidazol-1-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-N-(2-trimethylsiloxymethyl)[1,1'-biphenyl]-2-sulfonamide

10

A mixture of **32F** (1.5 g, 2.0 mmol), tetraethylorthocarbonate (6 ml), and acetic acid (0.15 ml) was heated under a nitrogen atmosphere at 70 °C
15 for 2 h. The mixture was cooled and concentrated, and the residue chromatographed on silica using 1:2 hexanes/ethyl acetate as eluant to provide **32G** (0.80 g) as a yellow oil.

H. 2'-[(3,3-Dimethyl-2-oxopyrrolidin-1-yl)methyl]-4'-[(2-ethoxy-7-(methoxycarbonyl)benzimidazol-1-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-[1,1'-biphenyl]-2-sulfonamide


20

32G (0.80 g) was subjected to sulfonamide deprotection using TBAF in THF according to General Method 10. The crude product was purified
25 by silica gel chromatography using 3:2 hexanes/acetone as eluant to provide 0.55 g of the title compound as a white solid; mp 101-103 °C (decomp); MS m/e 686 (ESI+ mode); HPLC retention time 3.91 min (Method A); HPLC purity >98%.

Example 33

2'-[(3,3-Dimethyl-2-oxopyrrolidin-1-yl)methyl]-4'-[(2-ethoxy-7-(carboxy)benzimidazol-1-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-[1,1'-biphenyl]-2-sulfonamide

5




32 (0.31 g) was subjected to ester hydrolysis according to General Method 15. Purification by reverse-phase preparative HPLC provided the title compound (14 mg) as a white solid; MS m/e 672 (ESI+ mode); HPLC retention time 3.61 min (Method A); HPLC purity 82%.

10

Example 34

2'-[(3,3-Dimethyl-2-oxopyrrolidin-1-yl)methyl]-4'-[(2-ethoxy-7-(N-methylcarbamoyl)benzimidazol-1-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-[1,1'-biphenyl]-2-sulfonamide

15




20

33 (80 mg) was subjected to amide formation according to General Method 12 using methylamine as the amine component. The product was purified by reverse-phase preparative HPLC: white solid (7 mg); MS m/e 685 (ESI+ mode); HPLC retention time 3.39 min (Method A); HPLC purity 81%.

25

Example 35


2'-[(3,3-Dimethyl-2-oxopyrrolidin-1-yl)methyl]-4'-[(2-ethoxy-7-(N,N-
dimethylcarbamoyl)benzimidazol-1-yl)methyl]-N-(3,4-dimethyl-5-
isoxazolyl)-[1,1'-biphenyl]-2-sulfonamide



- 10 **33** (80 mg) was subjected to amide formation according to General
 Method 12 using dimethylamine as the amine component. The crude
 product was subjected to reverse-phase preparative HPLC to give the title
 compound as a white solid (6 mg); MS m/e 699 (ESI+ mode); HPLC
 retention time 3.46 min (Method A); HPLC purity 66% [contaminant
 15 (34%) is the corresponding imidazolin-2-one].

Example 36

4'-[(2-Ethylquinolin-4-yl)oxymethyl]-N-(1,3,5-trimethylpyrazol-4-yl)
[1,1'-biphenyl]-2-sulfonamide



A. 4'-[(4-Bromophenyl)methoxy]-2-ethylquinoline

A mixture of 2-ethyl-4-quinolone (1.0 g, 5.8 mmol), 4-bromobenzyl bromide (1.7 g, 6.9 mmol), potassium carbonate (1.6 g, 11.6 mmol), and DMF (10 ml) was stirred at RT for 16 h. Ethyl acetate (100 ml) was added and the mixture was washed four times with water, then once with brine.

- 5 The organic layer was dried over sodium sulfate and evaporated, and the residue was triturated with 1:1 hexanes/ethyl acetate to provide 1.6 g **36A** as a white solid.

10 B. 4'-[(2-Ethylquinolin-4-yl)oxymethyl]-N-(tert-butyl)-[1,1'-biphenyl]-2-sulfonamide

- A mixture of **36A** (1.5 g, 4.4 mmol) and [2-(*N*-tert-butylsulfamoyl)phenyl]boronic acid (2.3 g, 8.7 mmol) was subjected to Suzuki coupling according to General Method 1. The crude product was chromatographed on silica gel using 1:1 hexanes/ethyl acetate as eluant to
15 provide the **36B** (1.8 g) as a yellow solid.

C. 4'-[(2-Ethylquinolin-4-yl)oxymethyl]-[1,1'-biphenyl]-2-sulfonamide

- A solution of **36B** (1.8 g, 3.8 mmol) in 4 ml dichloromethane and 8 ml TFA was stirred at RT for 14 h, then was heated at reflux for 8 h. The
20 solvent was evaporated and the residue partitioned between ethyl acetate and aqueous sodium bicarbonate. A small amount of a solid precipitate was retained with the organic layer. The organic layer was washed twice with water, then was concentrated. The crude product was crystallized from 1:1 toluene/ethyl acetate to provide **36C** (1.2 g) as a yellow solid.

25

D. 4'-[(2-Ethylquinolin-4-yl)oxymethyl] [1,1'-biphenyl]-2-sulfonic acid

- A suspension of **36C** (1.1 g, 2.7 mmol) in 25 ml acetonitrile was treated at 0 °C with nitrosonium tetrafluoroborate (370 mg, 3.2 mmol). After 30 min the mixture was allowed to warm to RT and was stirred at
30 RT for 4 h. The mixture was concentrated to provide **36D** (1.2 g) as a crude white solid.


E. 4'-[(2-Ethylquinolin-4-yl)oxymethyl]-N-(1,3,5-trimethylpyrazol-4-yl)[1,1'-biphenyl]-2-sulfonamide

A suspension of **36D** (100 mg, 0.24 mmol) in thionyl chloride (4 ml) was treated with DMF (10 μ l). The resulting mixture was refluxed for 45 min, then the solvent was evaporated. The residue was twice taken up in toluene and evaporated to dryness. Pyridine (2.5 ml) and 4-amino-1,3,5-trimethylpyrazole (90 mg, 0.72 mmol) were added and the mixture was stirred at RT for 14 h, then the solvent was evaporated. The residue was purified by silica gel chromatography (100:1 chloroform/methanol eluant), followed by silica gel thin-layer chromatography (1:1 hexanes/acetone eluant) to provide the title compound (24 mg) as an amorphous white solid: mp 212-215 $^{\circ}$ C (decomp); MS m/e 527 (ESI+ mode); HPLC retention time 3.18 min (Method C); HPLC purity 94%.

15

Example 37

4'-[(2-Ethylquinolin-4-yl)oxymethyl]-N-(3-methylisoxazol-5-yl)[1,1'-biphenyl]-2-sulfonamide




20

A suspension of the product of **36D** (370 mg, 0.88 mmol) was subjected to the procedure used for Example 36, Step E, substituting 5-amino-3-methylisoxazole as the amine component. The crude product was purified by reverse-phase preparative HPLC to provide the title compound (8 mg) as an amorphous tan solid: MS m/e 500 (ESI+ mode); HPLC retention time 3.32 min (Method A); HPLC purity >98%.

25

Example 38

4'-[(5-Acetyl-2-n-propyl-4-chloroimidazol-1-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-[1,1'-biphenyl]-2-sulfonamide



5

A. 5-(1-Hydroxyethyl)-2-n-propyl-4-chloroimidazole

A solution of 2-n-propyl-4-chloroimidazole-5-carboxaldehyde (Watson, S.P. *Synth. Comm.*, **1992**, 22, 2971-2977) (1.5 g, 8.7 mmol) in THF (50 ml) was treated dropwise at 0 °C with methylmagnesium bromide (8.7 ml of a 3.0 M solution in ether). Upon completion of the addition, the mixture was allowed to warm to RT and was stirred for 2h. The mixture was cooled again to 0 °C and was quenched with the addition of 1N hydrochloric acid. The mixture was adjusted to pH 8-9 by the addition of aqueous dipotassium hydrogen phosphate solution, then was partitioned against ethyl acetate. The ethyl acetate layer and an accompanying precipitate were collected and the solvent was evaporated to provide **38A** (1.6 g) as a slightly yellow solid.

20

B. 5-Acetyl-4-chloro-2-n-propyl-imidazole

A mixture of **38A** (1.6 g), activated manganese dioxide (5.7 g), and dioxane (20 ml) was heated at 55 °C for 48 h. The mixture was cooled and filtered through celite, and the filter cake was rinsed with dichloromethane. The combined filtrates were evaporated and the residue was chromatographed on silica gel using 3:1 hexanes/ethyl acetate as eluant. The product was further purified by trituration with 9:1 hexanes/ethyl acetate, providing 0.6 g **38B** as an orange solid.

25

C. 4'-[(5-Acetyl-2-n-propyl-4-chloroimidazol-1-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-N-(2-trimethylsiloxymethyl)[1,1'-biphenyl]-2-sulfonamide


5 **38B** (148 mg, 0.79 mmol) was alkylated with **P19** (300 mg, 0.53 mmol) according to General Method 22. The crude product was purified by silica gel chromatography using 2:1 hexanes/ethyl acetate as eluent to provide **38C** (89 mg) as a yellow oil.

10 D. 4'-[(5-Acetyl-2-n-propyl-4-chloroimidazol-1-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-[1,1'-biphenyl]-2-sulfonamide

Deprotection of **38C** (60 mg) according to General Method 8, followed by preparative thin-layer chromatography using 1:1 hexanes/aceteone as eluant, provided the title compound (24 mg) as a
 15 white solid: MS m/e 528 (ESI+ mode); HPLC retention time 3.75 min (Method A); HPLC purity 98%.

Example 39

20 4'-[(5-Methoxycarbonyl-2-n-propyl-4-chloroimidazol-1-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-[1,1'-biphenyl]-2-sulfonamide



25 A. 4'-[(5-Formyl-2-n-propyl-4-chloroimidazol-1-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-N-(2-trimethylsiloxymethyl)[1,1'-biphenyl]-2-sulfonamide

P19 (300 mg) was used to alkylate 2-n-propyl-4-chloroimidazole-5-carboxaldehyde according to General Method 22. The crude product was chromatographed on silica gel using 4:1 hexanes/ethyl acetate as eluant to provide **39A** (200 mg) as a yellow oil.

5

B. 4'-[(5-Carboxy-2-n-propyl-4-chloroimidazol-1-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-N-(2-trimethylsiloxyethyl)[1,1'-biphenyl]-2-sulfonamide

Sodium chlorite (19 mg, 0.21 mmol) was added to a mixture of **39A** (90 mg, 0.14 mmol) and sulfamic acid (20 mg, 0.21 mmol) in 1:1 THF/water (8 ml) at 0 °C. The mixture was stirred at 0 °C for 1h, then saturated potassium bisulfate solution was added. The aqueous layer was extracted twice with ethyl acetate. The combined organic extracts were washed once with brine, then were dried over sodium sulfate and concentrated to provide **39B** (62 mg) as a yellow oil.

15

C. 4'-[(5-Carboxy-2-n-propyl-4-chloroimidazol-1-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl) 1,1'-biphenyl]-2-sulfonamide

39B (62 mg) was deprotected according to General Method 8, using water in place of an alcohol as co-solvent. Crude **39C** (54 mg) was a yellow oil.

20

D. 4'-[(5-Methoxycarbonyl-2-n-propyl-4-chloroimidazol-1-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl) 1,1'-biphenyl]-2-sulfonamide


39C (54 mg) was subjected to ester formation according to General Method 20. Reverse-phase preparative HPLC provided the title compound (9 mg) as a white solid: MS m/e 544 (ESI+ mode); HPLC retention time 3.94 min (Method A); HPLC purity >98%

25

Example 40

4'-[(5-(N,N-dimethylcarbamoyl)-2-n-propyl-4-chloroimidazol-1-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-[1,1'-biphenyl]-2-sulfonamide

5



A. 4'-[(5-Ethoxycarbonyl-2-n-propyl-4-ethylimidazol-1-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-N-(2-methoxyethoxymethyl)[1,1'-biphenyl]-2-sulfonamide

10

Ethyl 2-n-propyl-4-ethylimidazole-5-carboxylate (94 mg, 0.45 mmol) was alkylated with **P18** (380 mg, 0.37 mmol), according to General Method 22 to provide crude **40A** as a 3:1 mixture of N-1 and N-3 regioisomeric alkylation products.

15

B. 4'-[(5-Carboxy-2-n-propyl-4-ethylimidazol-1-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-[1,1'-biphenyl]-2-sulfonamide

20

A solution of **40A** (0.45 g) in dioxane (3 ml) and 6N hydrochloric acid (3 ml) was heated at 70 °C for 2 h. The mixture was cooled to RT and made basic (pH > 14) with the addition of 45% aqueous potassium hydroxide solution. Additional dioxane and water were added to obtain a clear solution and the mixture was stirred at RT for 16 h, followed by heating at 70 °C for 3 h. The pH was adjusted to 2-3 with the addition of 6N hydrochloric acid and solid trisodium phosphate, and the mixture was extracted with three portions of ethyl acetate. The combined organic extracts were dried over sodium sulfate and evaporated to give **40B** (0.33 g) as a crude oil.


25

C. 4'-[(5-(N,N-dimethylcarbamoyl)-2-n-propyl-4-ethylimidazol-1-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-[1,1'-biphenyl]-2-sulfonamide

- 5 **40B** (110 mg) was subjected to amide formation according to General Method 12 using dimethylamine as the amine component. The crude material was subjected to reverse-phase preparative HPLC to provide the title compound (19 mg) as a white solid: MS m/e 550 (ESI+ mode); HPLC retention time 19.54 min (Method B); HPLC purity 74%.
- 10 The contaminant (24%) is the isomeric product arising from N-3 alkylation of the imidazole in Step A (HPLC retention time 19.77 min).

Example 41

- 15 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-hydroxymethyl[1,1'-biphenyl]-2-sulfonamide



- 20 A. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-N-[(2-trimethylsilylethoxy)methyl]-2'-hydroxymethyl[1,1'-biphenyl]-2-sulfonamide

- P14** (243 mg, 0.41 mmol) was used to alkylate 2-butyl-4-oxo-1,3-diazaspiro[4.4]non-1-ene hydrochloride according to General Method 4.
- 25 **41A** (100 mg, 35% yield) was isolated as a slightly yellow oil after silica gel chromatography using 1:1 hexanes/ethyl acetate as eluant.


B. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-hydroxymethyl[1,1'-biphenyl]-2-sulfonamide

Deprotection of **41A** (100 mg, 0.14 mmol) according to General Method 8 (ethanol) gave the title compound as white solid in 46% yield following silica gel chromatography (96:4 methanol/chloroform eluant): MS m/e 565 (ESI+ mode); HPLC retention time 3.21 min (Method A); HPLC purity >98%.

10

Example 42

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-ethoxymethyl[1,1'-biphenyl]-2-sulfonamide



15

A. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-N-[(2-methoxyethoxy)methyl]-2'-hydroxymethyl[1,1'-biphenyl]-2-sulfonamide

Triethylsilane (6 ml) and TFA (6 ml) were added to a solution of **5F** (960 mg, 1.5 mmol) in 15 ml dichloromethane at RT. The mixture was stirred at RT for 2 h and was then concentrated. The residue was taken up in ethyl acetate and was washed successively with aqueous sodium bicarbonate, water, and brine. The organic layer was dried over sodium sulfate and concentrated. The residue was chromatographed on silica gel using 100:2 dichloromethane/methanol to afford **42A** (740 mg, 77%) as a colorless gum. Rf=0.13, silica gel, 100:5 dichloromethane/methanol.

B. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-N-[(2-methoxyethoxy)methyl]-2'-ethoxymethyl[1,1'-biphenyl]-2-sulfonamide


A mixture of **42A** (100 mg, 0.15 mmol), iodoethane (960 mg, 6.1 mmol) and silver (I) oxide (180 mg, 0.77 mmol) in 0.7 ml DMF was heated at 40 °C for 16 h.. Additional iodoethane (190 mg, 1.2 mmol) and silver (I) oxide (71 mg, 0.31 mmol) were added and the reaction mixture was heated at 40 °C for an additional 4 h. The mixture was diluted with 1:4 hexanes/ethylacetate and was then washed with water and brine. The organic layer was dried over sodium sulfate and was then concentrated. The residue was chromatographed on silica gel using 200:3 dichloromethane/methanol as eluant to afford **42B** (51mg, 49%) as a colorless gum. Rf=0.35, silica gel, 100:5 dichloromethane/methanol.

C. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-ethoxymethyl[1,1'-biphenyl]-2-sulfonamide

42B (51 mg) was deprotected according to General Method 7 to afford the title compound in 80% yield following preparative reverse-phase HPLC purification: white solid; m.p. 74-80 °C (amorphous); ¹H NMR (CDCl₃) δ0.87(tr, J=7Hz, 3H), 0.99(tr, J=7Hz, 3H), 1.32(m, 2H), 1.59(m, 2H), 1.75-2.02(m, 11H), 2.16(s, 3H), 2.35(m, 2H), 3.38 (m, 2H), 4.23(m, 2H), 4.73(s, 2H), 7.11-7.85 (m, 7H); MS m/e 593 (ESI+ mode); HPLC retention time 18.22 min. (Method E); HPLC purity >97%.

Example 43

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-propyl[1,1'-biphenyl]-2-sulfonamide



- A. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-[(2-methoxyethoxy)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-(2-methoxyvinyl)[1,1'-biphenyl]-2-sulfonamide


5F was treated with methoxymethyltriphenylphosphonium bromide according to the procedure used in Example 27, Step A. The product (34%) was generated as a mixture of E and Z isomers.

- B. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-methoxyethyl[1,1'-biphenyl]-2-sulfonamide

43A (18 mg) was treated with triethylsilane and TFA according to the procedure of Example 42, Step B, to provide the title compound (6 mg, 45%) as an oil: MS m/e 593 (ESI+ mode); HPLC retention time 24.74 min. (Method B); HPLC purity >98%.

Example 44

- 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,3-dimethyl-2-oxo-1-pyrrolidinyl)methyl]-N-(3-methoxy-5-methyl-2-pyrazinyl)[1,1'-biphenyl]-2-sulfonamide



- A. N-(3-Methoxy-5-methyl-2-pyrazinyl)-2-bromobenzenesulfonamide

2-amino-3-methoxy-5-methylpyrazine (1.50 g, 10.8 mmol; synthesized according to Bradbury, R. H., et. al. *J. Med. Chem.* **1997**, *40*, 996-1004) and 2-bromobenzenesulfonyl chloride (2.80 g, 11.0 mmol) were reacted according to the procedure of Example 25, Step A. **44A** was a pink solid, 2.0 g (52%).

B. N-(3-Methoxy-5-methyl-2-pyrazinyl)-N-[2-(trimethylsilyl)ethoxymethyl]-2-bromobenzenesulfonamide
44A (2.0 g) was reacted with 2-(trimethylsilyl)ethoxymethyl chloride (1.15 ml) according to the procedure of Example 25, Step B. The crude residue was chromatographed on silica gel using 4:1 hexanes/ethyl acetate to give **44B** (2.37 g, 86%) as a yellow oil.

C. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-(1,3-dioxolan-2-yl)-N-[2-(trimethylsilyl)ethoxymethyl]-N-(3-methoxy-5-methyl-2-pyrazinyl)[1,1'-biphenyl]-2-sulfonamide

A solution of **5E** (2.0 g, 4.6 mmol) in ether (45 ml) was treated at -78 °C with t-butyllithium (1.7 M in pentane, 5.9 ml, 10.1 mmol). After stirring at -78 °C for 10 min, the mixture was treated with trimethylborate (1.3 ml, 11.5 mmol) and was then allowed to warm to RT. The mixture was concentrated and the residue azeotroped twice with methanol to produce a pale yellow solid (3.5 g).

The crude solid was subjected to Suzuki coupling with **44B** (2.37 g) according to General Method 1. Silica gel chromatography using 1:1 hexanes/ethyl acetate as eluant provide 1.45 g **44C** (39%) as a yellow oil.

D. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-formyl-N-(3-methoxy-5-methyl-2-pyrazinyl)[1,1'-biphenyl]-2-sulfonamide

44C (1.45 g, 1.9 mmol) was treated with 2M sulfuric acid (11 ml) and ethanol (11 ml) at RT for 6 h. Aqueous sodium bicarbonate was added

(final pH 7) and the mixture was extracted with ethyl acetate. The combined organic extracts were dried over sodium sulfate and concentrated to provide crude **44D** (0.91 g) as an off-white solid.


5 E. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,3-dimethyl-2-oxo-1-pyrrolidinyl)methyl]-N-(3-methoxy-5-methyl-2-pyrazinyl)[1,1'-biphenyl]-2-sulfonamide

44D (0.91 g) was reacted with ethyl 4-amino-2,2-dimethylbutanoate hydrochloride according to General Method 5. The crude residue was
 10 purified by silica gel column chromatography using 1:3 hexanes/ethyl acetate as eluant to provide the title compound (230 mg, 18% over two steps) as a yellow solid: MS m/e 687 (ESI+ mode); HPLC retention time 3.58 min (Method A); HPLC purity 95%.

15

Example 45

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4-bromo-3-methyl-5-isoxazolyl)-2'-[(3,3-dimethyl-2-oxo-1-pyrrolidinyl)methyl][1,1'-biphenyl]-2-sulfonamide



20

A. N-(2-methoxyethoxymethyl)-N-(3-methyl-5-isoxazolyl)-2-bromobenzenesulfonamide

25 **25A** (10.0 g, 31.5 mmol) was reacted with MEM chloride according to the procedure of Example 25, Step B. The crude residue was chromatographed on silica gel using 2:1 hexanes/ethyl acetate to give 4.8 g **45A** (38%) as a yellow oil.

B. [2-[[[(3-methyl-5-isoxazolyl)](2-methoxyethoxy)-methyl]amino]sulfonyl]phenyl]boronic acid

n-Butyllithium (1.35 M solution in hexanes, 9.7 ml, 13 mmol) was added dropwise over 5 min to a 0.2 M solution of the **45A** (4.8 g, 12 mmol) in THF at -90 °C. After 10 min, trimethylborate (1.6 ml, 14 mmol) was added and the mixture was allowed to warm to RT and stirred for 30 min. The mixture was cooled to 0 °C and treated with 21 ml of 3N hydrochloric acid, after which it was allowed to warm to RT over 30 min. Brine was added and the mixture was extracted with dichloromethane. The combined extracts were dried over sodium sulfate and concentrated to give **45B** (5.3 g) as a yellow oil.

C. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(2-methoxyethoxymethyl)-N-(3-methyl-5-isoxazolyl)-2'-formyl[1,1'-biphenyl]-2-sulfonamide

45B (4.5 g) was subjected to Suzuki coupling with **5E** according to General Method 1. Silica gel chromatography using 1:2 hexanes/ethyl acetate as eluant provide **45C** (1.30 g, 17%) as a yellow oil.

D. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3-methyl-5-isoxazolyl)-N-[2-methoxyethoxy)methyl]-2'-[(3,3-dimethyl-2-oxo-1-pyrrolidiny)methyl][1,1'-biphenyl]-2-sulfonamide

45C (420 mg, 0.66 mmol) was reacted with ethyl 4-amino-2,2-dimethylbutanoate hydrochloride and sodium cyanoborohydride using a procedure similar to that of Example 8, Step B. The crude residue was chromatographed on silica gel using 1:3 hexanes/ethyl acetate as eluant to provide **45D** (150 mg) as a yellow oil.

E. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3-methyl-5-isoxazolyl)-2'-[(3,3-dimethyl-2-oxo-1-pyrrolidinyl)methyl][1,1'-biphenyl]-2-sulfonamide


45D (110 mg) was deprotected according to General Method 7. The crude residue was chromatographed on silica gel using 95:5 chloroform/methanol as eluant to provide **45E** (50 mg) as a white solid.








F. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4-bromo-3-methyl-5-isoxazolyl)-2'-[(3,3-dimethyl-2-oxo-1-pyrrolidinyl)methyl][1,1'-biphenyl]-2-sulfonamide








45E (33 mg) was brominated using NBS according to the procedure used for Example 26. Preparative TLC purification of the crude residue provided the title compound (4 mg) as a white powder: MS m/e 724, 726 (ESI+ mode); HPLC retention time 3.53 min (Method A); HPLC purity 98%.









Examples 46 to 97








The following compounds **46** to **97** were prepared by a solution phase combinatorial chemistry method using **2I** and the corresponding carboxylic acid in the presence of diisopropylcarbodiimide. The products were purified by ion-exchange chromatography according to the General Method. HPLC retention times were determined using HPLC Method C.




















Ex. No.	Compound Name	J	HPLC Retention Time (min)	LRMS m/z [MH ⁺]
46	4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-[(formylmethylamino)-methyl][1,1'-biphenyl]-2-sulfonamide		3.23	606
47	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N-methylpropanamide		3.28	634
48	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N-methylcyclopropanecarbox-amide		3.33	646
49	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N,2-dimethylpropanamide		3.41	648
50	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N-methylbutanamide		3.45	648
51	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-2-methoxy-N-methylacetamide		3.29	650
52	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N-methyl-4-pentynamide		3.32	658







53	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N-methylcyclobutanecarbox-amide		3.50	660
54	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N,3-dimethylbutanamide		3.57	662
55	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N,2,2-trimethylpropanamide		3.80	662
56	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-3-methoxy-N-methylpropanamide		3.38	664
57	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-2-ethoxy-N-methylacetamide		3.25	664
58	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N-methyl-2-furancarboxamide		3.30	672
59	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N,4-dimethylpentanamide		3.95	676

60	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N-methylbenzamide		3.66	682
61	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N-methyl-3-thiophenecarboxamide		3.59	688
62	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N-methylcyclopentaneacetamide		3.82	688
63	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N-methylcyclohexanecarboxamide		3.76	688
64	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N,3-dimethylbenzamide		3.82	696
65	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N-methylbenzeneacetamide		3.78	696
66	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-2-fluoro-N-methylbenzamide		3.46	700
67	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-3-fluoro-N-methylbenzamide		3.53	700





68	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-4-fluoro-N-methylbenzamide		3.72	700
69	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N-methylcyclohexaneacetamide		3.96	702
70	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-2-fluoro-N-methylbenzeneacetamide		3.79	714
71	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-3-fluoro-N-methylbenzeneacetamide		3.82	714
72	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-4-fluoro-N-methylbenzeneacetamide		3.82	714
73	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N,N',N'-trimethylurea		3.44	649
74	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N'-(1,1-dimethylethyl)-N-methylurea		3.74	677






75	[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]methylcarbamic acid ethyl ester		3.69	650
76	[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]methylcarbamic acid 2-methylpropyl ester		4.01	678
77	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N,3,3-trimethylbutanamide		3.85	676
78	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N-methyl-2-pyridinecarboxamide		3.33	683
79	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N-methyl-3-pyridinecarboxamide		3.16	683
80	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N-methyl-2-pyrazinecarboxamide		3.27	684
81	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N,1-dimethyl-1H-pyrrole-2-carboxamide		3.64	685
82	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N-methyl-1,2,3-thiadiazole-4-carboxamide		3.36	690






83	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N,5-dimethyl-2-pyrazinecarboxamide		3.37	698
84	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N,3,5-trimethyl-4-isoxazolecarboxamide		3.42	701
85	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N,3-dimethyl-2-thiophenecarboxamide		3.66	702
86	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N,5-dimethyl-2-thiophenecarboxamide		3.72	702
87	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-3-cyano-N-methylbenzamide		3.45	707
88	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-4-cyano-N-methylbenzamide		3.49	707
89	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-2-methoxy-N-methylbenzamide		3.63	712
90	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-2-chloro-N-methylbenzamide		3.77	717
91	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-3-chloro-N-methylbenzamide		3.84	717

92	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-4-chloro-N-methylbenzamide		3.87	717
93	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-2,3-difluoro-N-methylbenzamide		3.66	718
94	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-3,4-difluoro-N-methylbenzamide		3.76	718
95	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-3,5-difluoro-N-methylbenzamide		3.76	718
96	4-Acetyl-N-[[4-[(2-butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N-methylbenzamide		3.48	724
97	N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-3-ethoxy-N-methyl-2-thiophenecarboxamide		3.63	732

The following examples were synthesized by combinations of the General Methods.






E X A M P L E	S T R U C T U R E	N A M E	S T A R T I N G M A T E R I A L	General Methods Applied (yield, %)	M/z (MH ⁺)	HPLC % Puri- ty	HPLC ret time, min (meth- od)
98		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)[1,1'-biphenyl]-2-sulfonamide	P18	4, 7 (32)	535	>98	15.00 (I)
99		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-[(propylsulfonyl)amino][1,1'-biphenyl]-2-sulfonamide	123	24 (67)	656	98	14.58 (E)
100		N-[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N-(1-oxopentyl)-L-valine methyl ester	P17	5,6 (16); 9 (60)	556	>98	16.27 (F)
101		N-[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N-(1-oxopentyl)-L-valine	100	15 (40)	542	>98	13.55 (F)

102		N-(3,4-Dimethyl-5-isoxazolyl)-2'-[(3,3-dimethyl-2-oxo-1-pyrrolidinyl)methyl]-4'-[(2-propyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl][1,1'-biphenyl]-2-sulfonamide	5C	4 (65); 19, 1 (70); 5 (36); 9 (50)	646	97	7.92 (E)
103		N-[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl]-4-[(2-ethyl-5,7-dimethyl-3H-imidazo[4,5-b]pyridin-3-yl)methyl][1,1'-biphenyl]-2-yl)methyl]-N,3,3-trimethylbutanamide	2G	4 (32); 7, 6 (80)	657	>98	25.45 (B)
104		N ² -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N ² -(1-oxopentyl)-L-valinamide	101	12, NH ₃ (70)	541	98	11.39 (B)
105		N ² -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N-methyl-N ² -(1-oxopentyl)-L-valinamide	101	12, MeNH ₂ (67)	555	98	11.27 (F)
106		N ² -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N,N-dimethyl-N ² -(1-oxopentyl)-L-valinamide	101	12, Me ₂ NH (12)	569	94	12.28 (F)

107		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-[[[(2,2,2-trifluoroethyl)amino)methyl][1,1'-biphenyl]-2-sulfonamide	P4	3 (90); 4 (37); 5 (56); 8, EtOH (31)	646	>99	3.46 (A)
108		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[[[(3,4-dimethyl-5-isoxazolyl)amino)sulfonyl][1,1'-biphenyl]-2-carboxylic acid	110	15 (62)	578	95	3.16 (A)
109		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-(trifluoromethyl)[1,1'-biphenyl]-2-sulfonamide	P9	4 (60); 10 (90)	603	94	2.33 (D)
110		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[[[(3,4-dimethyl-5-isoxazolyl)amino)sulfonyl][1,1'-biphenyl]-2-carboxylic acid methyl ester	methyl 2-bro- mo-5- methyl- benzo- ate	13, 4 (61); 1 (50); 8, MeOH (34)	593	96	3.42 (A)
111		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-(methoxymethyl)[1,1'-biphenyl]-2-sulfonamide	P2	4 (53); 7 (58)	579	>98	16.26 (E)






112		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-fluoro[1,1'-biphenyl]-2-sulfonamide	P11	4 (82); 7 (85)	553	>99	25.90 (B)
113		2'-(Cyanomethyl)-N-(3,4-dimethyl-5-isoxazolyl)-4'-[(2-ethyl-5,7-dimethyl-3H-imidazo[4,5-b]pyridin-3-yl)methyl][1,1'-biphenyl]-2-sulfonamide	P12	16, 3, 4 (35); 8, EtOH (81)	555	97	3.16 (A)
114		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-(cyanomethyl)-N-(3,4-dimethyl-5-isoxazolyl)[1,1'-biphenyl]-2-sulfonamide	P12	16, 3, 4 (35); 8, EtOH (52)	574	96	3.36 (A)
115		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-cyano-N-(3,4-dimethyl-5-isoxazolyl)[1,1'-biphenyl]-2-sulfonamide	P3	3 (88); 4 (45); 10 (40)	560	>97	3.31 (C)
116		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-methyl[1,1'-biphenyl]-2-sulfonamide	P7	11 (83); 2 (87); 4 (70); 1 (35); 7 (20)	549	98	27.21 (B)

117		2'-Cyano-N-(3,4-dimethyl-5-isoxazolyl)-4'-[(2-ethyl-5,7-dimethyl-3H-imidazo[4,5-b]pyridin-3-yl)methyl][1,1'-biphenyl]-2-sulfonamide	P3	3 (88); 4 (45); 10 (38)	541	98	3.10 (A)
118		N ² -[[2'-[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl]-2-methyl[1,1'-biphenyl]-4-yl)methyl]-N-methyl-N ² -(1-oxopentyl)-L-valinamide	P15	5 (39); 6 (92); 10 (30)	569	>99	2.15 (H)
119		N-(3,4-Dimethyl-5-isoxazolyl)-4'-[(2-ethyl-5,7-dimethyl-3H-imidazo[4,5-b]pyridin-3-yl)methyl]-2'-[[2,2,2-trifluoroethyl)amino]methyl][1,1'-biphenyl]-2-sulfonamide	P4	3 (90); 4 (49); 5 (83); 10 (15)	627	89	3.10 (A)
120		N-[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl]benzeneacetamide	122	6 (27)	668	97	14.97 (I)
121		N-[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl]-3,3-dimethylbutanamide	122	6 (36)	648	>98	16.38 (I)






122		2'-Amino-4'-[(2-butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)[1,1'-biphenyl]-2-sulfonamide	123	18 (75)	550	>98	8.39 (I)
123		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-nitro[1,1'-biphenyl]-2-sulfonamide	P22	7 (93)	580	>98	11.17 (I)
124		N ² -[[2'-[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N-methyl-N ² -(1-oxopropyl)-L-isoleucinamide	P16	5 (85); 6, 10 (62)	541	96	1.00 (G)
125		N ² -(Cyclopropylcarbonyl)-N ² -[[2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N-methyl-L-isoleucinamide	P16	5 (85); 6, 10 (62)	553	96	1.09 (G)
126		N ² -[[2'-[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N-methyl-N ² -(1-oxo-3-phenylpropyl)-L-isoleucinamide	P16	5 (85); 6, 10 (64)	617	95	1.63 (G)

127		N ² -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N-methyl-N ² -(3-methyl-1-oxobutyl)-L-isoleucinamide	P16	5 (85); 6, 10 (58)	569	98	1.47 (G)
128		N ² -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N-methyl-N ² -(1-oxohexyl)-L-isoleucinamide	P16	5 (85); 6, 10 (75)	583	98	1.74 (G)
129		N ² -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N-methyl-N ² -(1-oxobutyl)-L-isoleucinamide	P16	5 (85); 6, 10 (62)	555	95	1.31 (G)
130		N ² -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N-methyl-N ² -(1-oxopropyl)-L-leucinamide	P16	5 (83); 6, 10 (75)	541	90	1.01 (G)
131		N ² - (Cyclopropylcarbonyl)-N ² -[[2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N-methyl-L-leucinamide	P16	5 (83); 6, 10 (79)	553	95	1.11 (G)






132		N ² -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N-methyl- N ² -(1-oxo-3-phenylpropyl)-L-leucinamide	P16	5 (83); 6, 10 (20)	617	95	1.08 (G)
133		N ² -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N-methyl- N ² -(phenylacetyl)-L-leucinamide	P16	5 (83); 6, 10 (20)	603	95	1.68 (G)
134		N ² -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N-methyl- N ² -(3-methyl-1-oxobutyl)-L-leucinamide	P16	5 (83); 6, 10 (62)	569	90	1.51 (G)
135		N ² -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N-methyl- N ² -(1-oxohexyl)-L-leucinamide	P16	5 (83); 6, 10 (47)	583	97	1.76 (G)
136		N ² -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N-methyl- N ² -(1-oxobutyl)-L-leucinamide	P16	5 (83); 6, 10 (64)	555	94	1.29 (G)
137		N ² -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N-methyl- N ² -(1-oxopropyl)-L-valinamide	P16	5 (89); 6, 10 (39)	527	90	0.73 (G)






138		N ² - (Cyclopropylcarbo- nyl)- N ² -[[2'-[[[(3,4- dimethyl-5- isoxazolyl)amino] sulfonyl][1,1'- biphenyl]-4- yl]methyl]-N- methyl-L- valinamide	P16	5 (89); 6, 10 (66)	539	98	0.89 (G)
139		N ² -[[2'-[[[(3,4- Dimethyl-5- isoxazolyl)amino] sulfonyl][1,1'- biphenyl]-4- yl]methyl]-N- methyl- N ² -(1-oxo- 3-phenylpropyl)- L-valinamide	P16	5 (89); 6, 10 (6)	603	95	1.63 (G)
140		N ² -[[2'-[[[(3,4- Dimethyl-5- isoxazolyl)amino] sulfonyl][1,1'- biphenyl]-4- yl]methyl]-N- methyl- N ² - (phenylacetyl)-L- valinamide	P16	5 (89); 6, 10 (54)	589	93	1.53 (G)
141		N ² -[[2'-[[[(3,4- Dimethyl-5- isoxazolyl)amino] sulfonyl][1,1'- biphenyl]-4- yl]methyl]-N- methyl- N ² -(3- methyl-1- oxobutyl)-L- valinamide	P16	5 (89); 6, 10 (45)	555	98	1.37 (G)
142		N ² -[[2'-[[[(3,4- Dimethyl-5- isoxazolyl)amino] sulfonyl][1,1'- biphenyl]-4- yl]methyl]-N- methyl- N ² -(1- oxohexyl)-L- valinamide	P16	5 (89); 6, 10 (61)	569	98	1.58 (G)

143		N ² -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-methyl-N ² -(1-oxobutyl)-L-valinamide	P16	5 (89); 6, 10 (43)	541	98	1.65 (G)
144		N ² -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-ethyl-N ² -(1-oxopentyl)-L-valinamide	P17	5 (87); 6 (93); 9 (65); 15 (85); 12 (65)	569	>97	12.94 (F)
145		N ² -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-hexyl-N ² -(1-oxopentyl)-L-valinamide	P17	5 (87); 6 (93); 9 (65); 15 (85); 12 (65)	625	>98	22.19 (F)
146		N ² -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl]-2-cyano[1,1'-biphenyl]-4-yl]methyl]-N-methyl-N ² -(1-oxopentyl)-L-valinamide	P3	17 (58); 5, 6 (49); 10 (14)	580	>98	3.78 (C)
147		N ² -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl]-2-hydroxymethyl[1,1'-biphenyl]-4-yl]methyl]-N-methyl-N ² -(1-oxopentyl)-L-valinamide	P1	5 (77); 6 (92); 10 (17)	585	94	3.82, 3.90 (A) (two diastereomers)

148		4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-carboxamide	108	12 (33)	578	99	3.16 (A)
149		N,N-Dimethyl-4-[(2-butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-carboxamide	108	12 (61)	606	99	3.22 (A)
150		N-Methyl-4-[(2-butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-carboxamide	108	12 (93)	592	99	3.17 (A)
151		N-(3,4-Dimethyl-5-isoxazolyl)-4'-[(1,4,5,6,7,8-hexahydro-8-oxo-2-propyl-1-cycloheptimidazolyl)methyl]-2'-(methoxymethyl)[1,1'-biphenyl]-2-sulfonamide	P2	11 (87); 3 (38); 8 (42)	577	>98	11.58 (E)
152		N-(3,4-Dimethyl-5-isoxazolyl)-4'-[[[(3-methoxy-2,6-dimethyl-4-pyridinyl)oxy]methyl]-2'-methyl[1,1'-biphenyl]-2-sulfonamide	P13	4 (20); 7 (27)	508	>98	22.83 (B)

153		N-(3,4-Dimethyl-5-isoxazolyl)-4'-[(1,4,5,6,7,8-hexahydro-8-oxo-2-propyl-1-cycloheptimidazolyl)methyl]-2'-methyl[1,1'-biphenyl]-2-sulfonamide	P13	21 (60); 7 (20)	547	>98	23.29 (B)
154		N ² -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl]-2-(methoxymethyl)][1,1'-biphenyl]-4-yl)methyl]-N-methyl-N ² -(1-oxopentyl)-L-valinamide	P20	5 (90); 6 (90); 7 (42)	599	>98	12.71, 12.95 (F), migrates as diastereomers
155		N-(3,4-Dimethyl-5-isoxazolyl)-4'-[(1,4,5,6,7,8-hexahydro-8-oxo-2-propyl-1-cycloheptimidazolyl)methyl]-2'-(hydroxymethyl)[1,1'-biphenyl]-2-sulfonamide	P14	21 (29); 8, EtOH (32)	563	>98	2.99 (A)
156		2'-Chloro-N-(3,4-dimethyl-5-isoxazolyl)-4'-[[[(5,6,7,8-tetrahydro-2-ethyl-4-quinolinyl)oxy]methyl][1,1'-biphenyl]-2-sulfonamide	P10	4, 8 (33)	553	97	1.69 (D)
157		N ² -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl]-2-fluoro[1,1'-biphenyl]-4-yl)methyl]-N-methyl-N ² -(1-oxopentyl)-L-valinamide	P8	5 (30); 6 (34); 7 (45)	573	>98	30.02 (B)

158		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-(phenoxy)methyl[1,1'-biphenyl]-2-sulfonamide	42A	3 (70); 22 (68); 7 (76)	641	94	19.91 (I)
159		N ² -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl]-2-(1H-pyrazol-1-yl)methyl][1,1'-biphenyl]-4-yl)methyl]-N-methyl-N ² -(1-oxopentyl)-L-valinamide	P1	5 (77); 6 (92); 3 (99); 4 (69); 23 (18)	635	95	4.08 (A)
160		N ² -(Cyclopropylcarbonyl)-N ² -[[2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N,N-dimethyl-L-valinamide	P16	5 (85); 6, 10 (28)	553	88	2.58 (D)
161		N ² -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N,N-dimethyl-N ² -(1-oxobutyl)-L-valinamide	P16	5 (85); 6, 10 (33)	555	93	2.82 (D)
162		N ² -(Cyclopropylcarbonyl)-N ² -[[2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl]-2-(methoxymethyl)[1,1'-biphenyl]-4-yl)methyl]-N,N-dimethyl-L-valinamide	P2	5 (50); 6, 8, EtOH (56)	597	>98	2.50, 2.63 (D) (migrates as diastereomers)

163		N ² -[[2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl]-2-(methoxymethyl)][1,1'-biphenyl]-4-yl)methyl]-N,N-dimethyl-N ² -(1-oxobutyl)-L-valinamide	P2	5 (50); 6, 8, EtOH (59)	599	96	2.92, 3.02 (D) (migrates as diastereomers)
164		N ² -[[2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N-methyl-N ² -(1-oxopentyl)-L-valinamide	P5	5 (69); 6, 10 (50)	590	96	3.03 (D)
165		N ² -[[2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl]-2-(trifluoromethyl)][1,1'-biphenyl]-4-yl)methyl]-N-methyl-N ² -(1-oxopentyl)-L-valinamide	P6	5 (80); 6, 10 (83)	623	95	2.93, 3.05 (D) (migrates as diastereomers)
166		N ² -(Cyclobutylcarbonyl)-N ² -[[2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N-methyl-L-valinamide	P16	5 (85); 6, 10 (35)	553	98	2.82 (D)
167		1-[[2-Chloro-2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-4-ethyl-2-propyl-1H-imidazole-5-carboxylic acid	P10	22 (60); 8, EtOH (90); 15 (99)	558	>98	3.04 (A)






168		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-[(methylsulfonyl)amino][1,1'-biphenyl]-2-sulfonamide	122	24 (25)	628	>98	11.48 (E)
169		(S)-N-[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N-[2-methyl-1-(4-methyl-1-piperazinyl)carbonyl]propyl]pentanamide	101	12 (43)	624	>98	28.26 (B)
170		(S)-N-[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N-(1-piperidinyl)carbonyl]propyl]pentanamide	101	12 (30)	609	>98	34.45 (B)
171		N-(3,3-Dimethylbutyl)-N ² -[[2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N ² -(1-oxopentyl)-L-valinamide	101	12 (38)	625	>98	33.97 (B)
172		N ² -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N-[(4-fluorophenyl)methyl]-N ² -(1-oxopentyl)-L-valinamide	101	12 (38)	649	>98	32.90 (B)







173		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-[(methylethoxy)methyl][1,1'-biphenyl]-2-sulfonamide	42A	2 (67); 4 (51); 7 (62)	607	97	13.59 (E)
174		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-(propoxymethyl)[1,1'-biphenyl]-2-sulfonamide	42A	2 (67); 4 (49); 7 (68)	607	>97	20.61 (E)
175		4-Chloro-1-[[2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-2-propyl-1H-imidazole-5-carboxamide	P19	22 (87); 25 (47); 8, H ₂ O (99); 12 (35)	529	>98	3.69 (A)
176		N-(3,4-Dimethyl-5-isoxazolyl)-2'-fluoro-4'-[(1,4,5,6,7,8-hexahydro-8-oxo-2-propyl-1-cycloheptimidazolyl)methyl][1,1'-biphenyl]-2-sulfonamide	P8	21 (15); 7 (16)	551	>98	5.78 (J)
177		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)[1,1'-biphenyl]-2-sulfonamide	42A	2 (67); 4 (72); 7 (21)	642	>97	13.59 (E)

178		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-(1H-pyrazol-1-ylmethyl)[1,1'-biphenyl]-2-sulfonamide	P1	3 (98); 4, 11 (35); 3, 4, 7 (13)	615	97	3.42 (A)
179		2-Butyl-4-chloro-1-[[2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-1H-imidazole-5-carboxamide]	P19	22 (43); 25; 8, H ₂ O;12 (19)	543	94	3.82 (A)
180		N-(3,4-Dimethyl-5-isoxazolyl)-4'-[[[(2-methyl-4-quinolinyl)oxy]methyl][1,1'-biphenyl]-2-sulfonamide	P18	4 (79); 8, EtOH (26)	500	98	22.4 (B)
181		N-(3,4-Dimethyl-5-isoxazolyl)-4'-[[[(2-ethyl-4-quinolinyl)oxy]methyl][1,1'-biphenyl]-2-sulfonamide	P18	4 (49); 8, EtOH (30)	514	93	23.25 (B)
182		N-(3,4-Dimethyl-5-isoxazolyl)-4'-[[[(2-ethyl-5,6,7,8-tetrahydro-4-quinolinyl)oxy]methyl][1,1'-biphenyl]-2-sulfonamide	P18	4 (79); 8, EtOH (3)	518	93	3.49 (A)
183		N-(3,4-Dimethyl-5-isoxazolyl)-4'-[[[(2-propyl-4-quinolinyl)oxy]methyl][1,1'-biphenyl]-2-sulfonamide	P18	4 (47); 8, EtOH (20)	528	97	3.50 (A)






184		N-(3,4-Dimethyl-5-isoxazolyl)-4'-[(5,6,7,8-tetrahydro-2,4-dimethyl-7-oxopyrido[2,3-d]pyrimidin-8-yl)methyl][1,1'-biphenyl]-2-sulfonamide	P18	4 (79); 8, EtOH (20)	518	>98	3.05 (A)
185		4'-[[[(2-Ethyl-4-quinolinyl)oxy)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-[(3,3-dimethyl-2-oxo-1-pyrrolidinyl)methyl][1,1'-biphenyl]-2-sulfonamide	P18	4 (20); 8, EtOH (33)	639	93	3.43 (A)
186		N-(3,4-Dimethyl-5-isoxazolyl)-2'-[(3,3-dimethyl-2-oxo-1-pyrrolidinyl)methyl]-4'-[[[(5,6,7,8-tetrahydro-2-ethyl-4-quinolinyl)oxy)methyl][1,1'-biphenyl]-2-sulfonamide	P18	4 (63); 8, EtOH (8)	643	92	3.79 (C)
187		3-[[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-2-ethyl-N-methyl-3H-benzimidazole-4-carboxamide	31	12 (12)	544	92	2.80 (A)
188		1-[[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-2-ethyl-1H-benzimidazole-7-carboxylic acid phenylmethyl ester	31	20 (55)	621	96	3.95 (C)






189		1-[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-2-ethyl-1H-benzimidazole-7-carboxylic acid 2-phenylethyl ester	31	20 (12)	635	>98	4.03 (C)
190		1-[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-2-ethyl-1H-benzimidazole-7-carboxylic acid 2-(2-oxo-1-pyrrolidinyl)ethyl ester	31	20 (45)	642	>98	3.15 (C)
191		1-[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-2-ethyl-1H-benzimidazole-7-carboxylic acid 3-(2-oxo-1-pyrrolidinyl)propyl ester	31	20 (30)	656	>98	3.21 (C)
192		2'-Cyano-N-(3,4-dimethyl-5-isoxazolyl)-4'-[[[(2-ethyl-4-quinolinyloxy)methyl][1,1'-biphenyl]-2-sulfonamide	P3	3, 4 (59); 10 (60)	539	>98	3.01 (C)
193		2'-(Cyanomethyl)-N-(3,4-dimethyl-5-isoxazolyl)-4'-[(1,4,5,6,7,8-hexahydro-8-oxo-2-propyl-1-cycloheptimidazolyl)methyl][1,1'-biphenyl]-2-sulfonamide	P12	16, 3 (73); 21 (46); 8, EtOH(16)	572	96	2.95 (A)

194		3-[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-5-ethyl-N-methyl-2-propyl-3H-imidazole-4-carboxamide	40B	12, MeNH ₂ (12)	536	>98	2.88 (C)
195		1-[[2-Chloro-2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-4-ethyl-2-propyl-1H-imidazole-5-carboxamide	167	12, NH ₃ (17)	557	>98	3.08 (A)
196		3-[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-5-ethyl-2-propyl-3H-imidazole-4-carboxamide	40B	12, NH ₃ (16)	522	>98	2.79 (C)
197		3-[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-2-ethoxy-N-methyl-3H-benzimidazole-4-carboxamide	29	12, MeNH ₂ (32)	560	>98	3.31 (A)
198		3-[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-2-ethoxy-N,N-dimethyl-3H-benzimidazole-4-carboxamide	29	12, Me ₂ NH (5)	574	>98	3.42 (A)





199		2-[[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]propylamino]-3-pyridinecarboxylic acid	P18	4, 15, 7 (34)	521	99	23.4 (B)
200		4'-[[[(3,5-Dibutyl-1H-1,2,4-triazol-1-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)[1,1'-biphenyl]-2-sulfonamide	P18	4, 7 (77)	522	99	28.4 (B)
201		N-(3,4-Dimethyl-5-isoxazolyl)-4'-[[[(1,4,5,6,7,8-hexahydro-8-oxo-2-propyl-1-cycloheptimidazolyl)methyl][1,1'-biphenyl]-2-sulfonamide	P18	21, 7 (55)	533	99	21.5 (B)
202		4'-[(2,7-Diethyl-5H-pyrazolo[1,5-b][1,2,4]triazol-5-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)[1,1'-biphenyl]-2-sulfonamide	P18	4, 7 (33)	505	99	21.8 (B)
203		N-[2-Butyl-3-[[[2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-3,4-dihydro-4-oxo-6-quinazolinyl]-N'-methyl-N'-(1-methylethyl)urea	P18	4, 7 (71)	657	99	28.7 (B)
204		2-[[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]propylamino]-N-methyl-3-pyridinecarboxamide	199	12 (98)	534	99	23.8 (B)

205		4'-[[[(3-Methoxy-2,6-dimethyl-4-pyridinyl)oxy)methyl]-N-(3,4-dimethyl-5-isoxazolyl)[1,1'-biphenyl]-2-sulfonamide	P18	4 (59); 8 (74)	494	98	6.99 (I)
206		N-(3,4-Dimethyl-5-isoxazolyl)-2'-[[[(3,3-dimethyl-2-oxo-1-pyrrolidinyl)methyl]-4'-[[[(3-methoxy-2,6-dimethyl-4-pyridinyl)oxy)methyl]-1,1'-biphenyl]-2-sulfonamide	20A	2 (82); 4 (82); 8 (31)	619	96	14.02 (E)
207		N-(3,4-Dimethyl-5-isoxazolyl)-2'-[[[(3,3-dimethyl-2-oxo-1-pyrrolidinyl)methyl]-4'-[(1,4,5,6,7,8-hexahydro-8-oxo-2-propyl-1-cycloheptimidazolyl)methyl]-1,1'-biphenyl]-2-sulfonamide	20A	2 (82); 21, 9 (54)	658	97	15.56 (E)
208		N-(3,4-Dimethyl-5-isoxazolyl)-4'-[[[(3-methoxy-2,6-dimethyl-4-pyridinyl)oxy)methyl]-1,1'-biphenyl]-2-sulfonamide	P2	4 (45); 8 (63)	538	99	10.60 (E)
209		3-[[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl]-1,1'-biphenyl]-4-yl)methyl]-2-ethyl-5-methyl-3H-imidazole-4-carboxamide	P19	22 (35); 7 (67)	494	>97	7.96 (E)

210		4'-[(2-Butyl-3,4-dihydro-4-oxo-3-quinazolinyl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)[1,1'-biphenyl]-2-sulfonamide	P18	4, 7 (13)	543	98	25.52 (I)
211		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-[(3,3-dimethyl-2-oxo-1-pyrrolidinyl)methyl][1,1'-biphenyl]-2-sulfonamide	17	5 (36)	660	98	14.51 (I)
212		N²-[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl]-2'-[(3,3-dimethyl-2-oxo-1-pyrrolidinyl)methyl][1,1'-biphenyl]-4-yl)methyl]-N-methyl-N²-(1-oxopentyl)-L-valinamide	20A	17 (80); 5 (95); 6 (95); 7 (42)	680	97	13.13, 14.66 (F) migrates as diastereomers
213		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-[(4,4-dimethyl-2-oxo-1-pyrrolidinyl)methyl][1,1'-biphenyl]-2-sulfonamide	3	5 (58)	660	>98	24.60 (B)
214		4'-[(3,5-Dibutyl-1H-1,2,4-triazol-1-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-[(3,3-dimethyl-2-oxo-1-pyrrolidinyl)methyl][1,1'-biphenyl]-2-sulfonamide	20A	4 (52); 9 (57)	647	97	23.53 (I)

215		N-[2-[[[2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl]-4-[(2-ethyl-5,7-dimethyl-3H-imidazo[4,5-b]pyridin-3-yl)methyl][1,1'-biphenyl]-2-yl)methyl]methylamino]ethyl]acetamide	23B	5, 12 (4); 8 (15)	644	89	2.69 (A)
216		4-[(2-Ethyl-5,7-dimethyl-3H-imidazo[4,5-b]pyridin-3-yl)methyl]-2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-acetic acid ethyl ester	113	8, EtOH (13)	602	90	3.41 (A)
217		N ² -[[2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N ² -(1-oxopentyl)-N-propyl-L-valinamide	101	12 (65)	583	>98	14.97 (F)
218		N ² -[[2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N ² -(1-oxopentyl)-N-[(tetrahydro-2-furanyl)methyl]-L-valinamide	101	12 (43)	625	>98	14.06 (F)
219		2'-Chloro-N-(3,4-dimethyl-5-isoxazolyl)-4'-[[[(2-ethyl-4-quinolinyloxy)methyl][1,1'-biphenyl]-2-sulfonamide	P10	4 (90); 8 (65)	549	98	1.68 (D)


220		N-(3,4-Dimethyl-5-isoxazolyl)-4'-[[[(2-ethyl-4-quinolinyl)oxy)methyl]-2'-(trifluoromethyl)][1,1'-biphenyl]-2-sulfonamide	P9	4, 8 (30)	582	97	1.75 (D)
221		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-chloro-N-(3,4-dimethyl-5-isoxazolyl)[1,1'-biphenyl]-2-sulfonamide	P10	4, 8 (42)	570	98	2.13 (D)
222		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-[(2-methylpropoxy)methyl][1,1'-biphenyl]-2-sulfonamide	42A	2 (67); 4 (49); 7 (85)	621	>98	23.07 (B)
223		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-[(ethylsulfonyl)amino][1,1'-biphenyl]-2-sulfonamide	123	24 (33)	642	98	14.32 (E)
224		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-[(2,2,2-trifluoroethoxy)methyl][1,1'-biphenyl]-2-sulfonamide	P2A	4, F ₃ CCH ₂ OH (89); 14 (76); 1 (67); 11 (95); 2 (90); 4 (85); 7 (85)	647	>97	20.15 (E)

225		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-(2-fluoroethoxy)methyl[1,1'-biphenyl]-2-sulfonamide	P2A	4, FCH ₂ C H ₂ OH (72); 14 (65); 1 (68); 11 (95); 2 (85); 4 (80); 7 (90)	611	>97	15.94 (E)
226		N-(3,4-Dimethyl-5-isoxazolyl)-4'-[[[(3-methoxy-2,6-dimethyl-4-pyridinyl)oxy]methyl]-2'-ethoxymethyl[1,1'-biphenyl]-2-sulfonamide	P2A	4, EtOH (77); 14 (80); 1 (75); 11 (95); 2 (77); 22 (43); 7 (74)	552	97	12.60 (E)
227		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-(ethoxymethyl)[1,1'-biphenyl]-2-sulfonamide	P2A	4, EtOH (77); 14 (80); 1 (70); 11 (98); 2 (80); 4 (83); 7 (86)	593	>98	18.75 (E)
228		N-(3,4-Dimethyl-5-isoxazolyl)-4'-[(1,4,5,6,7,8-hexahydro-8-oxo-2-propyl-1-cycloheptimidazolyl)methyl]-2'-(ethoxymethyl)[1,1'-biphenyl]-2-sulfonamide	P2A	4, EtOH (77); 14 (80); 1 (75); 11 (95); 2 (77); 21 (); 7 ()	591	>98	20.07 (B)

Example 229

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethylisoxazol-5-yl)-2'-(3,3,3-trifluoropropyl)[1,1'-biphenyl]-2-sulfonamide

5



A. 3-Bromo-4-methoxybenzaldehyde dimethyl acetal

5 A solution of 3-bromo-4-methoxybenzaldehyde (19.2 g, 89 mmol) and trimethylorthoformate (14.8 ml, 140 mmol) in 150 ml methanol was treated at RT with concentrated sulfuric acid (10 μ l). After stirring for 16 h, the mixture was treated with potassium carbonate (70 mg) and the solvent was evaporated. The residue was partitioned between
10 dichloromethane and aqueous sodium bicarbonate. The organic layer was dried over sodium sulfate and concentrated to provide **229A** (22.6 g) as a brown oil, which was used without further purification.

B. 4-Methoxy-3-(3,3,3-trifluoropropyl)benzaldehyde dimethyl acetal

15 A solution of **229A** (15.8 g, 60 mmol) in dry THF (120 ml) was added to magnesium turnings (4.1 g, 170 mmol) at RT. The mixture was heated to 60 °C for 5 min, which initiated an exothermic reaction. The heating bath was removed and the mixture was allowed to reflux gently. Upon subsidence of the reaction (approximately 5 min), the mixture was heated
20 to reflux for an additional 20 min and then was allowed to cool to RT. The clear green supernatant was transferred via canula away from excess magnesium and into a separate flask containing copper (I) iodide (575 mg, 3.0 mmol) under a nitrogen atmosphere. To this mixture was added 1-iodo-3,3,3-trifluoropropane (15 g, 67 mmol) and the resulting mixture was
25 allowed to stir for 16 h, during which time a thick white precipitate developed. Aqueous ammonium chloride solution, ethyl acetate, and hexanes were added, and the organic layer was dried over sodium sulfate. Evaporation followed by silica gel chromatography of the residue using 3:1

hexanes/ethyl acetate as eluant provided 7.7 g of a mobile, slightly yellow oil, which was a 6:1 (mol ratio) mixture of **229B** and 4-methoxybenzaldehyde dimethyl acetal, as determined by proton NMR. This mixture was used without further purification.

5

C. 4-Methoxy-3-(3,3,3-trifluoropropyl)benzaldehyde

The **229B** mixture (7.7 g) was subjected to hydrochloric acid hydrolysis according to General Method 19. The resulting crude yellow oil (8.0 g) was a 6:1 mixture (mol ratio) of **229C** and 4-methoxybenzaldehyde, and was used without further purification.

10

D. 4-Hydroxy-3-(3,3,3-trifluoropropyl)benzaldehyde

A solution of the **229C** mixture (8.0 g, 35 mmol) in dichloromethane (5 ml) was treated at 0 °C with boron tribromide (55 ml of a 1.0 M solution in dichloromethane). The mixture was allowed to warm to RT and was stirred for 2 h. Aqueous 10% dipotassium hydrogen phosphate solution was added, and the aqueous layer was extracted with two portions of dichloromethane. The combined organic extracts were dried over sodium sulfate and evaporated. The residue was chromatographed on silica gel using 3:1 hexanes/ethyl acetate as eluant to provide 2.0 g of **229D** as an orange solid (contaminated with 4-hydroxybenzaldehyde).

15

20

E. 4-(trifluoromethanesulfonyloxy)-3-(3,3,3-trifluoropropyl)benzaldehyde

A solution of the **229D** mixture (2.0 g, 9 mmol) and N,N-diisopropylethylamine (2.4 ml, 14 mmol) in dichloromethane (40 ml) was treated dropwise at -78 °C with trifluoromethanesulfonic anhydride (1.9 ml, 11 mmol). The mixture was stirred at that temperature for 15 min following the completion of the addition. Aqueous sodium bicarbonate solution was added and the mixture was stirred and allowed to warm to RT. The layers were separated and the aqueous layer was extracted with

25

30

two portions of dichloromethane. The combined organic extracts were dried over sodium sulfate, evaporated, and the residue was chromatographed on silica gel using hexanes/ethyl acetate as eluant to provide first recovered **229D** (540 mg), and then the product fraction (380 mg) as an orange oil. The product fraction was a 2:1 mixture (mol ratio) of **229E** and 4-(trifluoromethanesulfonyloxy)benzaldehyde, as determined by proton NMR.

F. N-(3,4-Dimethyl-5-isoxazolyl)-4'-formyl-2'-(3,3,3-trifluoropropyl)-N-[(2-trimethylsilyl)ethoxymethyl] [1,1'-biphenyl]-2-sulfonamide

A solution of **229E** (380 mg, 1.1 mmol) and [2-[(3,4-dimethyl-5-isoxazolyl)[(2-(trimethylsilyl)ethoxy)methyl]amino]-sulfonyl]phenyl]boronic acid (925 mg, 2.2 mmol) in dioxane (10 ml) was sparged with nitrogen for 15 min.

Tetrakis(triphenylphosphine)palladium(0) (130 mg, 0.11 mmol) was added, followed by powdered potassium phosphate (460 mg, 2.2 mmol). The mixture was heated at 85 °C for 5 h, then the solvent was evaporated. The residue was chromatographed on silica gel using hexanes/ethyl acetate as eluant to provide the product fraction (550 mg) as an orange oil.

The product fraction was a 2:1 mixture (mol ratio) of **229F** and the corresponding 2'-H product, as judged by proton NMR.

G. N-(3,4-Dimethyl-5-isoxazolyl)-4'-hydroxymethyl-2'-(3,3,3-trifluoropropyl)-N-[(2-trimethylsilyl)ethoxymethyl] [1,1'-biphenyl]-2-sulfonamide

The **229F** mixture (550 mg, 0.9 mmol) was subjected to sodium borohydride reduction according to General Method 14. The crude residue was chromatographed directly on silica gel using 3:1 hexanes/ethyl acetate as eluant to provide the product fraction (355 mg) as an amber oil. The product fraction was a 2:1 mixture (mol ratio) of **229G** and the corresponding 2'-H product, as judged by proton NMR.

H. N-(3,4-Dimethyl-5-isoxazolyl)-4'-(methanesulfonyloxy)methyl-2'-(3,3,3-trifluoropropyl)-N-[(2-trimethylsilyl)ethoxymethyl] [1,1'-biphenyl]-2-sulfonamide

5 The **229G** mixture (350 mg, 0.63 mmol) was converted to the corresponding methanesulfonate ester according to General Method 3. The entire crude product **229H** was used directly in the next step.

I. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-(3,3,3-trifluoropropyl)-N-[(2-trimethylsilyl)ethoxymethyl] [1,1'-biphenyl]-2-sulfonamide


10 **229H** was used to alkylate 2-butyl-1,3-diazaspiro[4.4]non-1-en-4-one according to General Method 4. The crude residue was chromatographed on silica gel using 2:1 hexanes/ethyl acetate as eluant to provide the product fraction (370 mg) as a slightly orange oil. The product
15 fraction was a 2:1 mixture (mol ratio) of **229I** and the corresponding 2'-H product, as judged by proton NMR

J. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethylisoxazol-5-yl)-2'-(3,3,3-trifluoropropyl) [1,1'-biphenyl]-2-sulfonamide

20 The **229I** mixture was deprotected according to General Method 8. The crude product (which contained the 2'-H contaminant) was purified by reverse-phase preparative HPLC, followed by extraction with ethyl acetate from pH 8 potassium phosphate buffer. Finally, silica gel chromatography
25 using 2:1 hexanes/acetone as eluant provided the title compound (120 mg) as a white foam; LRMS m/z 631 (ESI+ mode); HPLC retention time 3.78 min (Method C); HPLC purity >98%.

Example 230

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethylisoxazol-5-yl)-2'-(3-fluoropropyl) [1,1'-biphenyl]-2-sulfonamide



5

A. Methyl 3-(2-propenyl)-4-(trifluoromethanesulfonyloxy)benzoate

Methyl 4-hydroxy-3-(2-propenyl)benzoate (12.0 g, 62.4 mmol, prepared according to W.J. Greenlee, et. al., WO 91/11999) was treated with trifluoromethanesulfonic anhydride according to the procedure of Example 229, Step E. The crude product was chromatographed on silica gel using 3:1 hexanes/ethyl acetate as eluant to yield 17.4 g of **230A** as a yellow oil.

15

B. Methyl 3-(3-hydroxypropyl)-4-(trifluoromethanesulfonyloxy)benzoate

Borane-THF complex (1.0 M solution in THF, 32 ml, 32 mmol) was added to a solution of **230A** (8.5 g, 26 mmol) in THF at 0 °C. The mixture was stirred at RT for 16h and then cooled to 0 °C. A 1.0 M solution of sodium/potassium phosphite buffer (pH 7, 50 ml) was added, followed by 30% aqueous hydrogen peroxide (9.0 ml). The mixture was allowed to warm to RT, then water and EtOAc were added. The organic layer was dried over sodium sulfate and evaporated. The crude product was chromatographed on silica gel using 1:1 hexanes/ethyl acetate as eluant to yield 4.3 g of **230B** as a yellow oil.

25

C. N-(3,4-Dimethyl-5-isoxazolyl)-2'-(3-hydroxypropyl)-4'-(methoxycarbonyl)-N-[(2-trimethylsilyl)ethoxymethyl] [1,1'-biphenyl]-2-sulfonamide

Suzuki coupling of **230B** (1.3 g, 3.8 mmol) according to the
5 procedure of Example 229, step F, provided **230C** (750 mg) as an orange oil following silica gel chromatography using 1:1 hexanes/ethyl acetate as eluant.

D. N-(3,4-Dimethyl-5-isoxazolyl)-2'-(3-fluoropropyl)-4'-(methoxycarbonyl)-N-[(2-trimethylsilyl)ethoxymethyl] [1,1'-biphenyl]-2-sulfonamide

A solution of **230C** (750 mg, 1.3 mmol) in dichloromethane (3 ml)
was treated at -78 °C with (diethylamino)sulfur trifluoride (0.21 ml, 1.6
mmol). The mixture was allowed to warm to RT. After 15 min, water was
15 added and the aqueous layer was extracted with two portions of
dichloromethane. The organic extracts were dried over sodium sulfate,
concentrated, and the residue was chromatographed on silica gel using 2:1
hexanes/ethyl acetate as eluant to provide **230D** (175 mg) as a yellow oil.

E. N-(3,4-Dimethyl-5-isoxazolyl)-2'-(3-fluoropropyl)-4'-(hydroxymethyl)-N-[(2-trimethylsilyl)ethoxymethyl] [1,1'-biphenyl]-2-sulfonamide

A solution of **230D** (175 mg, 0.3 mmol) in THF (5 ml) was treated
with DIBAL-H (0.53 ml of a 1.5 M solution in toluene, 0.8 mmol) at -78
25 °C. The temperature was allowed to rise to -25 °C and the mixture was
stirred for 2 h. Water (2 ml) and ether (10 ml) were added and the
mixture was stirred at RT for 2 h. The mixture was filtered and the
filtrate concentrated to provide crude **230E** (110 mg) as a colorless oil.

F. N-(3,4-Dimethyl-5-isoxazolyl)-2'-(3-fluoropropyl)-4'-(methanesulfonyloxy)methyl-N-[(2-trimethylsilyl)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

230E (110 mg, 0.2 mmol) was converted to the corresponding methanesulfonate ester according to General Method 3. The entire crude product **230F** was used directly in the next step.

G. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-(3-fluoropropyl)-N-[(2-trimethylsilyl)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide


230F was used to alkylate 2-butyl-1,3-diazaspiro[4.4]non-1-en-4-one according to General Method 4, to provide **230G** (124 mg) as a colorless oil, which was used without further purification.

H. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethylisoxazol-5-yl)-2'-(3-fluoropropyl)[1,1'-biphenyl]-2-sulfonamide

230G (124 mg, 0.2 mmol) was deprotected according to General Method 8 (EtOH). The crude product was purified by silica gel chromatography using 1:1 hexanes/ethyl acetate as eluant provided the title compound (23 mg) as a white foam; LRMS m/z 595 (ESI+ mode); HPLC retention time 2.10 min (Method D); HPLC purity >98%.

Example 231

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-(1,1-difluoroethyl)-N-(3,4-dimethylisoxazol-5-yl)[1,1'-biphenyl]-2-sulfonamide



A. 5-Bromo-2-(trifluoromethanesulfonyloxy)acetophenone

5-Bromo-2-hydroxyacetophenone (3.3 g, 15 mmol) was treated with
5 trifluoromethanesulfonic anhydride according to the procedure of Example
229, Step E. The crude residue was taken up in ether and washed twice
with 10% aqueous potassium dihydrogenphosphate solution and once with
brine. The ether solution was dried over magnesium sulfate and
concentrated to provide crude **231A** (4.7 g) as an orange oil, which was
10 used without further purification.

B. 3-(1,1,-Difluoroethyl)-4-(trifluoromethanesulfonyloxy)bromobenzene

231A (4.4 g, 13 mmol) was treated at RT with neat
(diethylamino)sulfur trifluoride (2.5 ml, 19 mmol) and the resulting
15 solution was stirred at RT for 40 h. The mixture was poured onto ice,
aqueous sodium bicarbonate solution was added, and the mixture was
extracted with three portions of dichloromethane. The combined extracts
were dried over sodium sulfate and concentrated, and the residue was
chromatographed on silica gel using 5:1 hexanes/ethyl acetate as eluant to
20 provide **231B** (4.0 g) as an orange oil.

C. 3-(1,1,-Difluoroethyl)-4-(trifluoromethanesulfonyloxy)benzaldehyde

A solution of **231B** (3.8 g, 10 mmol) and DMF (1.2 ml, 15 mmol) in
dry THF (60 ml) was treated dropwise at -78 °C with n-butyllithium (5.0
25 ml of a 2.5 M solution in hexanes, 13 mmol). 10% aqueous potassium
dihydrogenphosphate solution was added and the resulting mixture was
allowed to warm to RT. 1:1 hexanes/ethyl acetate and brine were added
and the organic layer was collected, dried over sodium sulfate, and
evaporated. The residue was chromatographed on silica gel using 3:1
30 hexanes/ethyl acetate as eluant to provide recovered **231B** (1.4 g), followed
by **231C** (1.0 g), which was an orange oil.

D. 2'-(1,1-Difluoroethyl)-N-(3,4-dimethylisoxazol-5-yl)-4'-formyl-N-[(2-trimethylsilyl)ethoxymethyl] [1,1'-biphenyl]-2-sulfonamide

Suzuki coupling of **231C** (1.0 g, 3.2 mmol) according to the
5 procedure of Example 229, step F, provided **231D** (640 mg) as an orange oil following silica gel chromatography using 3:1 hexanes/ethyl acetate as eluant.

E. 2'-(1,1-Difluoroethyl)-N-(3,4-dimethylisoxazol-5-yl)-4'-hydroxymethyl-N-[(2-trimethylsilyl)ethoxymethyl] [1,1'-biphenyl]-2-sulfonamide

231D (640 mg, 1.2 mmol) was subjected to sodium borohydride reduction according to General Method 14. The crude residue was chromatographed on silica gel using 1:1 hexanes/ethyl acetate as eluant to
15 provide **231E** (550 mg) as a brown oil.

F. 2'-(1,1-Difluoroethyl)-N-(3,4-dimethylisoxazol-5-yl)-4'-(methanesulfonyloxy)methyl-N-[(2-trimethylsilyl)ethoxymethyl] [1,1'-biphenyl]-2-sulfonamide

231E (360 mg, 0.7 mmol) was converted to the corresponding methanesulfonate ester according to General Method 3. The entire crude product **231F** was used directly in the next step.

G. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-(1,1-difluoroethyl)-N-(3,4-dimethylisoxazol-5-yl)-N-[(2-trimethylsilyl)ethoxymethyl] [1,1'-biphenyl]-2-sulfonamide

231F was used to alkylate 2-butyl-1,3-diazaspiro[4.4]non-1-en-4-one according to General Method 4. The crude residue was chromatographed on silica gel using 1:1 hexanes/ethyl acetate as eluant to provide **231G**
30 (170 mg) as a yellow oil.


H. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-(1,1-difluoroethyl)-N-(3,4-dimethylisoxazol-5-yl)-[1,1'-biphenyl]-2-sulfonamide

231G (150 mg, 0.2 mmol) was deprotected according to General Method 8 (EtOH). The crude product was chromatographed on silica gel using 1:1 hexanes/ethyl acetate as eluant to provide the title compound (36 mg) as a white foam; LRMS m/z 599 (ESI+ mode); HPLC retention time 3.58 min (Method A); HPLC purity >98%.

10

Example 232

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-(2,2,2-trifluoroethyl)-N-(3,4-dimethylisoxazol-5-yl) [1,1'-biphenyl]-2-sulfonamide



15

A. Methyl 4-bromo-3-(bromomethyl)benzoate

Methyl 4-bromo-3-methylbenzoate (20 g, 87 mmol) was brominated according to General Method 13. The crude orange solid was triturated with 4:1 hexanes/ethyl acetate to provide **232A** (14.7 g) as a white solid.

B. Methyl 4-bromo-3-(2,2,2-trifluoroethyl)benzoate

Methyl 2,2-difluoro-2-(fluorosulfonyl)acetate (3.1 ml, 25 mmol) was added to a mixture of **232A** (7.0 g, 22 mmol) and copper(I) iodide (420 mg, 2.2 mmol) in DMF (45 ml) under a nitrogen atmosphere. The mixture was heated to 85 °C for 16 h. The mixture was cooled to RT and the solvent was evaporated. Hexanes and brine were added, and the organic layer

was dried over sodium sulfate and concentrated to give a yellow solid (6.0 g), which contained **232A** and **232B**.

The solid was dissolved in DMF (15 ml) and stirred at RT with potassium acetate (1.0 g, 10 mmol) for 3 h. The solvent was evaporated and the residue partitioned between ethyl acetate and water. The organic layer was dried over sodium sulfate and concentrated, and the residue was chromatographed on silica gel using 4:1 hexanes/ethyl acetate as eluant to provide **232B** (3.4 g) as a white solid.

10 C. N-(3,4-Dimethylisoxazol-5-yl)-4'-(methoxycarbonyl)-2'-(2,2,2-trifluoroethyl)-N-[(2-trimethylsilyl)ethoxymethyl] [1,1'-biphenyl]-2-sulfonamide

Suzuki coupling of **232B** (1.8 g, 5.9 mmol) according to the procedure of Example 229, step F, provided **232C** (1.3 g) as a yellow solid following silica gel chromatography using 3:1 hexanes/ethyl acetate as eluant.

20 D. N-(3,4-Dimethylisoxazol-5-yl)-4'-hydroxymethyl-2'-(2,2,2-trifluoroethyl)-N-[(2-trimethylsilyl)ethoxymethyl] [1,1'-biphenyl]-2-sulfonamide

232C (1.3 g, 2.2 mmol) was treated with DIBAL-H according to the procedure of Example 230, step E, to provide **232D** (650 mg) as an oil following silica gel chromatography using 3:1 hexanes/ethyl acetate as eluant.

25 E. N-(3,4-Dimethylisoxazol-5-yl)-4'-(methanesulfonyloxy)methyl-2'-(2,2,2-trifluoroethyl)-N-[(2-trimethylsilyl)ethoxymethyl] [1,1'-biphenyl]-2-sulfonamide

232D (650 mg, 1.1 mmol) was converted to the corresponding methanesulfonate ester according to General Method 3. The entire crude product **232E** was used directly in the next step.

F. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethylisoxazol-5-yl)-2'-(2,2,2-trifluoroethyl)-N-[(2-trimethylsilyl)ethoxymethyl] [1,1'-biphenyl]-2-sulfonamide


5 **232E** was used to alkylate 2-butyl-1,3-diazaspiro[4.4]non-1-en-4-one according to General Method 4. The crude residue was chromatographed on silica gel using 2:1 hexanes/ethyl acetate as eluant to provide **232F** (440 mg) as a yellow oil.

10 G. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethylisoxazol-5-yl)-2'-(2,2,2-trifluoroethyl) [1,1'-biphenyl]-2-sulfonamide

15 **232F** was deprotected according to General Method 8 (EtOH). The crude product was purified by reverse-phase preparative HPLC, followed by an extraction with ethyl acetate from pH 5 sodium phosphate buffer to provide the title compound (78 mg) as a white foam; LRMS m/z 617 (ESI+ mode); HPLC retention time 1.69 min (Method H); HPLC purity >98%.

Example 233

20 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethylisoxazol-5-yl)-2'-[(2-methyl)propoxy] [1,1'-biphenyl]-2-sulfonamide



25

A. 2-Bromo-5-methylphenol

A solution of sodium nitrite (2.8 g, 41 mmol) in 5 ml water was added rapidly with stirring to an ice-cooled mixture of 6-amino-m-cresol

(5.0 g, 41 mmol) and 48% hydrobromic acid (17 ml, 100 mmol). The temperature was kept below 10 °C by addition of ice chips. The diazonium salt solution was then added in portions over a period of 30 min to a boiling mixture of copper(I) bromide (6.4 g, 22 mmol) and 48%
5 hydrobromic acid (5 ml). The resulting mixture was refluxed for an additional 30 min, then was cooled and extracted with ether (2x100ml). The combined organic extracts were washed with water, dried over magnesium sulfate, and evaporated. The residue was chromatographed on silica gel using 98:2 hexanes/ethyl acetate to afford **233A** (1.6 g, 20%) as
10 an oil.

B. 4-Bromo-3-(2-methylpropoxy)toluene

Isobutyl bromide (0.70 ml, 6.4 mmol) was added to a mixture of **233A** (800 mg, 4.3 mmol), potassium carbonate (1.2 g, 8.5 mmol), and
15 DMF (2 ml). The mixture was stirred for 36 h at 45 °C and was then cooled and concentrated under vacuum. The residue was added to water and extracted with ethyl acetate (3 X 50ml). The combined organic extracts were washed with water and dried over sodium sulfate to give **233B** (960 mg, 93%) as an oil.

20

C. 4-Bromomethyl-2-(2-methylpropoxy)-1-bromobenzene

233B (960 mg, 3.9 mmol) was subjected to NBS bromination according to General Method 13. The crude **233C** (1.1 g, 86%) was used without further purification.

25

D. 4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2-(2-methylpropoxy)-1-bromobenzene

233C (1.1 g, 3.4 mmol) was used to alkylate 2-butyl-1,3-diazaspiro[4.4]non-1-en-4-one according to General Method 4. The crude
30 residue was chromatographed on silica gel using 1:1 hexanes/ethyl acetate as eluant to provide **233D** (550 mg, 40%) as an oil.

E. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethylisoxazol-5-yl)-2'-[(2-methyl)propoxy]-N-[(2-trimethylsilyl)ethoxymethyl] [1,1'-biphenyl]-2-sulfonamide

5 Suzuki coupling of **233D** (550 mg, 1.3 mmol) according to General Method 1 provided **233E** (660 mg, 60%) as an oil following silica gel chromatography using 2:3 hexanes/ethyl acetate as eluant.

10 F. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethylisoxazol-5-yl)-2'-[(2-methyl)propoxy][1,1'-biphenyl]-2-sulfonamide


233E (660 mg) was deprotected according to General Method 8 (EtOH). Water was added to the crude residue and the mixture was extracted with ethyl acetate (3 X 50ml). The combined organic extracts
15 were washed with water and dried and evaporated. The residue was chromatographed on silica gel using 1:1 hexanes/ethyl acetate to provide the title compound (270 mg, 60%) as a white solid, mp 58-61°C ; LRMS m/e 607 (ESI+ mode); HPLC retention time 26.32 min (Method B); HPLC purity >98%.

20

Example 234

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethylisoxazol-5-yl)-2'-(2-methoxyethoxy) [1,1'-biphenyl]-2-sulfonamide

25



A. 4-Bromo-3-(2-methoxyethoxy)toluene

233A (800 mg, 4.3 mmol) was alkylated with 1-bromo-2-methoxyethane (0.89 g, 6.4 mmol) according to the procedure of Example 233, step B, to give **234A** (840 mg, 81%) as an oil.

5

B. 4-Bromomethyl-2-(2-methoxyethoxy)-1-bromobenzene

234A (840 mg, 3.4 mmol) was subjected to NBS bromination according to General Method 13. The crude **234B** (510 mg, 46%) was used without further purification.

10

C. 4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2-(2-methoxyethoxy)-1-bromobenzene

234B (510 mg, 1.6 mmol) was used to alkylate 2-butyl-1,3-diazaspiro[4.4]non-1-en-4-one according to General Method 4. The crude residue was chromatographed on silica gel using 4:1 hexanes/ethyl acetate as eluant to provide **234C** (600 mg, 88%) as an oil.

15

D. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethylisoxazol-5-yl)-2'-(2-methoxyethoxy)-N-[(2-trimethylsilyl)ethoxymethyl] [1,1'-biphenyl]-2-sulfonamide

20

Suzuki coupling of **234C** (600 mg, 1.4 mmol) according to General Method 1 provided **234D** (550 mg, 54%) as an oil following silica gel chromatography using 2:3 hexanes/ethyl acetate as eluant.

25

E. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethylisoxazol-5-yl)-2'-(2-methoxyethoxy)[1,1'-biphenyl]-2-sulfonamide

30


234D (550 mg, 0.8 mmol) was deprotected according to General Method 8 (EtOH). Water was added to the crude residue and the mixture was extracted with ethyl acetate (3 X 50ml). The combined organic extracts were washed with water and dried and evaporated. The residue

was purified by reverse-phase preparative HPLC to provide the title compound (125 mg, 30%) as a white solid, mp 55-58 °C ; MS m/e 609 (ESI+ mode); HPLC retention time 22.75 min (Method B); HPLC purity >98%.

5

Example 235

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-butyl-N-(3,4-dimethylisoxazol-5-yl)[1,1'-biphenyl]-2-sulfonamide



10

A. 4-Bromo-3-(1-butenyl)benzonitrile

2A (4.0 g, 19 mmol) was reacted with propyltriphenylphosphonium bromide following the procedure of Example 27, step A. The crude residue was chromatographed on silica gel using 9:1 hexanes/ethyl acetate to afford **235A** (2.2 g, 49%) as an E/Z mixture.

B. 4-Bromo-3-butylbenzonitrile

A mixture of **235A** (2.2 g, 9.3 mmol) and 210 mg of PtO₂ in 40 ml EtOH was hydrogenated at 35 PSI for 40 min. Filtration and concentration gave 1.4 g of **235B** (62%).

C. 4-Bromo-3-butylbenzaldehyde

235B (1.4 g, 5.8 mmol) was treated with DIBAL-H according to General Method 14 to provide crude **235C** (1.2 g, 90%) as an oil.

D. N-(3,4-Dimethyl-5-isoxazolyl)-2'-butyl-4'-formyl-N-[(2-trimethylsilyl)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

235C (1.2 g, 5.1 mmol) was subjected to Suzuki coupling according to General Method 1 to provide **235D** as a crude oil.

E. N-(3,4-Dimethyl-5-isoxazolyl)-2'-butyl-4'-hydroxymethyl-N-[(2-trimethylsilyl)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

235D (entire sample) was reduced with sodium borohydride in methanol according to General Method 11 to provide **235E** (1.4 g, 50% from **235C**) as an oil.

F. N-(3,4-Dimethyl-5-isoxazolyl)-4'-(methanesulfonyl)oxymethyl-2'-butyl-N-[(2-trimethylsilyl)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

235E (1.4 g, 2.5 mmol) was converted to the corresponding methanesulfonate ester according to General Method 3 to provide **235F** (1.4 g, 90%) as an oil.

G. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-butyl-N-[(2-trimethylsilyl)ethoxymethyl]-N-(3,4-dimethyl-5-isoxazolyl)[1,1'-biphenyl]-2-sulfonamide


235F (1.3 g, 2.1 mmol) was used to alkylate 2-butyl-1,3-diazaspiro[4.4]non-1-en-4-one according to General Method 4. **235G** (1.3 g, 85%) was produced as an oil.

H. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-butyl-N-(3,4-dimethyl-5-isoxazolyl)[1,1'-biphenyl]-2-sulfonamide

235G (1.2 g, 1.7 mmol) was deprotected according to General Method 7. The crude product was purified by reverse-phase preparative HPLC to provide the title compound (620 mg, 62%) as a solid, mp 58-61 °C; MS m/e 591 (ESI+ mode); HPLC retention time 26.67 min (Method B); HPLC purity >98%.

Example 236

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3-methylisoxazol-5-yl)-2'-trifluoromethyl[1,1'-biphenyl]-2-sulfonamide



5

A. [2-[[[(3-methyl-5-isoxazolyl)][(2-trimethylsilyl)ethoxy]methyl]amino]sulfonylphenyl]boronic acid

10 **25B** (14 g, 31 mmol) was converted to the corresponding boronic acid according to the procedure of Example 45, step B. The crude product **236A** (16.5 g, estimated purity 60%) was produced as an amber oil, and was used without further purification.

15 B. 4'-Formyl-N-(3-methyl-5-isoxazolyl)-2'-trifluoromethyl-N-[(2-trimethylsilyl)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

20 **236A** (8.0 g, 20 mmol) was subjected to Suzuki coupling with **P6** according to General Method 1. Silica gel chromatography using 2:1 hexanes/ethyl acetate as eluant provided 1.8 g of a mixture of **236B** and a highly crystalline impurity. Trituration with 2:1 hexanes/ethyl acetate to remove this impurity provided **236B** (1.0 g) as an orange oil.

C. 4'-Hydroxymethyl-N-(3-methyl-5-isoxazolyl)-2'-trifluoromethyl-N-[(2-trimethylsilyl)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

25 **236B** (880 mg, 1.6 mmol) was reduced with sodium borohydride (0.3 eq.) in ethanol according to General Method 11. The crude residue was chromatographed on silica gel using 2:1 hexanes/ethyl acetate as eluant to provide **236C** (450 mg) as a yellow oil.

D. 4'-(Methanesulfonyloxy)methyl-N-(3-methyl-5-isoxazolyl)-2'-trifluoromethyl-N-[(2-trimethylsilyl)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

5 **236C** (450 mg) was converted to the corresponding methanesulfonate ester according to General Method 3. The entire crude product **236D** was used in the next reaction step.

10 **E.** 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl] N-(3-methyl-5-isoxazolyl)-2'-trifluoromethyl-N-[(2-trimethylsilyl)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

236D was used to alkylate 2-butyl-1,3-diazaspiro[4.4]non-1-en-4-one according to General Method 4. The crude residue was chromatographed on silica gel using 2:1 hexanes/acetone as eluant to
15 provide **236E** (170 mg) as a yellow oil.


F. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl] N-(3-methyl-5-isoxazolyl)-2'-trifluoromethyl [1,1'-biphenyl]-2-sulfonamide

20 **236E** (100 mg) was deprotected according to General Method 8 (ethanol) to provide the title compound as the hydrochloride salt (77 mg), which required no additional purification: MS m/e 589 (ESI+ mode); HPLC retention time 3.72 min (Method C); HPLC purity 97%.

25

Example 237

N-(4-Bromo-3-methyl-5-isoxazolyl)-4'-[(2-butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-trifluoromethyl [1,1'-biphenyl]-2-sulfonamide




5 Bromine (33 mg/ml solution in acetic acid, 18 mg, 0.11 mmol) was
 added in portions to a solution of **236** (53 mg, 0.09 mmol) and sodium
 acetate (35 mg, 0.42 mmol) in acetic acid (4 ml) at RT. The solvent was
 evaporated, aqueous potassium phosphate solution was added, and the pH
 was adjusted to 8. The mixture was extracted with ethyl acetate, and the
 combined organic layers were dried over sodium sulfate. The residue was
 10 chromatographed on silica gel using 1:3 hexanes/acetone as eluant to
 provide the title compound (31 mg) as a white powder following
 lyophilization: MS m/e 667, 669 (ESI+ mode); HPLC retention time 3.83
 min (Method C); HPLC purity 94%.

15

Example 238

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4-chloro-3-
methyl-5-isoxazolyl)-2'-trifluoromethyl [1,1'-biphenyl]-2-sulfonamide



20

Clorox bleach (5.25% sodium hypochlorite solution, 225 µl) was
 added in portions to a solution of **236** (32 mg, 0.054 mmol) in THF (2 ml)
 at RT. The solvent was evaporated and the residue was purified by
 25 reverse-phase preparative HPLC, followed by preparative thin-layer silica


gel chromatography using 10% methanol in dichloromethane as eluant, providing the title compound (1.0 mg) as a white powder following lyophilization: MS m/e 624 (ESI+ mode); HPLC retention time 3.53 min (Method A); HPLC purity 95%.

5

Example 239

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-(N-methoxy-N-methylaminomethyl) [1,1'-biphenyl]-2-sulfonamide

10



A. N-(3,4-Dimethyl-5-isoxazolyl)-4'-hydroxymethyl-2'-(N-methoxy-N-methylaminomethyl)-N-[(2-methoxyethoxy)methyl] [1,1'-biphenyl]-2-sulfonamide

P21 (230 mg, 0.48 mmol) was subjected to reductive amination with N-methoxy-N-methylamine according to General Method 5, yielding **239A** (169 mg, 69%) as an oil.

20

B. 4'-Bromomethyl-N-(3,4-dimethyl-5-isoxazolyl)-2'-(N-methoxy-N-methylaminomethyl)-N-[(2-methoxyethoxy)methyl] [1,1'-biphenyl]-2-sulfonamide

239A (165 mg, 0.33 mmol) was converted to the corresponding bromide according to General Method 2, yielding **239B** (174 mg, 92%) as an oil following silica gel chromatography using hexanes/ethyl acetate as eluant.

25

C. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-(N-methoxy-N-methylaminomethyl)-N-[(2-methoxyethoxy)methyl][1,1'-biphenyl]-2-sulfonamide


239B (170 mg, 0,30 mmol) was used to alkylate 2-butyl-1,3-diazaspiro[4.4]non-1-en-4-one according to General Method 4. The crude residue was chromatographed on silica gel using hexanes/ethyl acetate as eluant to provide **239C** (155 mg, 72%) as a yellow oil.

D. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-(N-methoxy-N-methylaminomethyl)[1,1'-biphenyl]-2-sulfonamide

239C (150 mg) was deprotected according to General Method 7. The crude product was purified by reverse-phase preparative HPLC to provide the title compound (117 mg, 89%) as a white solid: MS m/e 608 (ESI+ mode); HPLC retention time 17.75 min (Method E); HPLC purity >97%.

Example 240

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-(2,2-difluoroethoxymethyl)-N-(3,4-dimethylisoxazol-5-yl)[1,1'-biphenyl]-2-sulfonamide



A. Methyl 4-bromo-3-(hydroxymethyl)benzoate

A mixture of methyl 4-bromo-3-methylbenzoate (168 g, 735 mmol), N-bromosuccinimide (141 g, 792 mmol), benzoyl peroxide (3.5 g, 15 mmol), and carbon tetrachloride (900 ml) was refluxed for 17 h. The mixture was

cooled and filtered, and the filtrate was washed once with water and once with brine. The filtrate was then dried over sodium sulfate and concentrated to provide an orange oil (261 g), which was judged (^1H NMR) be about 70 mol% methyl 4-bromo-3-(bromomethyl)benzoate.

- 5 The crude orange oil (261 g) was dissolved in 400 ml DMF and treated with potassium acetate (74 g, 750 mmol) at 0 °C. The mixture was allowed to warm to RT and was stirred for 54 h. The solvent was evaporated and the residue was taken up in 1:1 hexanes ethyl acetate and washed twice with half-saturated brine, then once with saturated brine.
- 10 The organic layer was dried over sodium sulfate and concentrated to provide a new orange oil (208 g), which was judged to contain about 70 mol% methyl 4-bromo-3-(acetoxymethyl)benzoate.

- The crude acetate mixture (208 g) was dissolved in methanol (1 l) and treated with potassium carbonate (12 g, 87 mmol) at 0 °C. The
- 15 mixture was allowed to warm to RT and was stirred for 18 h. The solvent was evaporated and the residue was treated with 130 ml 1N hydrochloric acid at 0 °C. The mixture was extracted once with 1:3 hexanes/ethyl acetate, once with 1:1 hexanes/ethyl acetate, and once with ethyl acetate. The combined organic layers were dried over sodium sulfate and
- 20 concentrated, and the residue was chromatographed on silica gel using 2:1 hexanes/ethyl acetate. The product fractions were combined, evaporated, and triturated with 1:1 hexanes/ethyl acetate to provide **240A** (102 g) as a white solid.

- 25 B. Methyl 4-bromo-3-[(tetrahydro-2H-pyran-2-yl)oxymethyl]benzoate
 p-Toluenesulfonic acid hydrate (50 mg) was added to a solution of **240A** (10 g, 41 mmol) and 3,4-dihydro-2H-pyran (10 g, 120 mmol) in dichloromethane (100 ml) at 0 °C. After 1h aqueous sodium bicarbonate solution was added, the layers were separated, and the organic layer was
- 30 dried over sodium sulfate and concentrated to provide crude **240B** (16 g) as a slightly yellow oil.

C. N-(3,4-Dimethyl-5-isoxazolyl)-4'-methoxycarbonyl-2'-[(tetrahydro-2H-pyran-2-yl)oxymethyl]-N-[(2-trimethylsilyl)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

5 Crude **240B** (10 g, approximately 25 mmol) was subjected to Suzuki coupling according to General Method 1. Silica gel chromatography using 3:1 hexanes/ethyl acetate as eluant provided **240C** (14 g) as a yellow oil.

D. N-(3,4-Dimethyl-5-isoxazolyl)-4'-hydroxymethyl-2'-[(tetrahydro-2H-pyran-2-yl)oxymethyl]-N-(2-trimethylsilyl)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

10 A solution of **240C** (10.5 g, 17 mmol) in THF (200 ml) was treated with DIBAL-H (23.4 ml of a 1.5 M solution in toluene, 35 mmol) at -78 °C. The temperature was allowed to rise to -25 °C and the mixture was stirred for 2 h. Water (10 ml) and ether (100 ml) were added and the mixture was stirred at RT for 2 h. Ethyl acetate and hexanes were added, and the mixture was washed three times with aqueous sodium bicarbonate solution. The organic layer was dried over sodium sulfate and concentrated to provide crude **240D** (9 g) as a deep maroon-colored oil.

20

E. N-(3,4-Dimethyl-5-isoxazolyl)-4'-(methanesulfonyloxy)methyl-2'-[(tetrahydro-2H-pyran-2-yl)oxymethyl]-N-[(2-trimethylsilyl)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

25 **240D** (4.5 g, 7.5 mmol) was converted to the corresponding methanesulfonate ester according to General Method 3. The entire crude product **240E** was used in the next reaction step.

F. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-[(tetrahydro-2H-pyran-2-yl)oxymethyl]-N-[(2-trimethylsilyl)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

30

240E was used to alkylate 2-butyl-1,3-diazaspiro[4.4]non-1-en-4-one according to General Method 4. The crude residue was chromatographed on silica gel using 3:2 hexanes/ethyl acetate as eluant to provide **240F** (6.0 g) as a slightly yellow oil.

5

G. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-hydroxymethyl-N-[(2-trimethylsilyl)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

A solution of **240F** (6.0 g, 7.7 mmol) and 2N hydrochloric acid (6 ml, 12 mmol) in methanol (150 ml) was stirred for 16 h at RT. The mixture was neutralized with aqueous sodium bicarbonate solution and the methanol was evaporated. Aqueous sodium bicarbonate solution was added and the mixture was extracted twice with ethyl acetate. The combined organic extracts were dried over sodium sulfate and concentrated to provide **240G** (5.0 g) as a crude orange oil.

H. 2'-Bromomethyl-4'-[(2-butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-N-[(2-trimethylsilyl)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

240G (590 mg, 0.85 mmol) was converted to the corresponding bromide according to General Method 2, providing **240H** (373 mg, 58%) as a yellow oil following silica gel chromatography using hexanes/ethyl acetate as eluant.

I. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-(2,2-difluoroethoxymethyl)-N-(3,4-dimethyl-5-isoxazolyl)-N-[(2-trimethylsilyl)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

240H (370 mg, 0.49 mmol) was used to alkylate 2,2-difluoroethanol according to General Method 4. **240I** (209 mg, 56%) was produced as a yellow oil following silica gel chromatography using hexanes/ethyl acetate as eluant.


J. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-(2,2-difluoroethoxymethyl)-N-(3,4-dimethylisoxazol-5-yl)[1,1'-biphenyl]-2-sulfonamide

5 **240I** (205 mg, 0.27 mmol) was deprotected according to General Method 7. The crude product was purified by reverse-phase preparative HPLC to provide the title compound (104 mg, 60%) as a white solid: MS m/e 629 (ESI+ mode); HPLC retention time 17.18 min (Method E); HPLC purity >97%.

10

Example 241

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethylisoxazol-5-yl)-2'-(2-fluoroethyl)[1,1'-biphenyl]-2-sulfonamide



15

A. Methyl 3-(2-hydroxyethyl)-4-(trifluoromethanesulfonyloxy)benzoate

An ozone/oxygen gas mixture was bubbled through a solution of
 20 **230A** (8.3 g, 26 mmol) in methanol (100 ml) at -78 °C until a light blue color persisted. The solution was sparged with nitrogen to remove excess ozone, then triphenylphosphine (10 g, 38 mmol) was added in portions. The cooling bath was removed and the mixture was allowed to warm RT, then was concentrated. Ethanol (100 ml) was added to the residue and
 25 the resulting mixture was cooled to 0 °C. Sodium borohydride (1.9 g, 51 mmol) was added. After 1h, the mixture was allowed to warm to RT and the solvent was evaporated. The residue was taken up in 10% aqueous potassium dihydrogen phosphate solution and was extracted with ethyl

acetate. The combined organic extracts were dried over sodium sulfate and concentrated, and the residue chromatographed on silica gel using 1:1 hexanes/ethyl acetate as eluant to yield **241A** (4.5 g) as a colorless oil.

- 5 B. N-(3,4-Dimethyl-5-isoxazolyl)-2'-(2-hydroxyethyl)-4'-(methoxycarbonyl)-N-[(2-trimethylsilyl)ethoxymethyl] [1,1'-biphenyl]-2-sulfonamide

 Suzuki coupling of **241A** (4.5 g, 14 mmol) according to the procedure of Example 229, step F, provided **241B** (4.1 g) as a yellow solid following
10 silica gel chromatography using 1:1 hexanes/ethyl acetate as eluant.

- C. N-(3,4-Dimethyl-5-isoxazolyl)-2'-(2-fluoroethyl)-4'-(methoxycarbonyl)-N-[(2-trimethylsilyl)ethoxymethyl] [1,1'-biphenyl]-2-sulfonamide

15 A mixture of **241B** (1.0 g, 1.8 mmol) and (diethylamino)sulfur trifluoride (0.71 ml, 5.4 mmol) was stirred at RT for 20 h. The mixture was poured onto ice. Aqueous sodium bicarbonate solution was added and the mixture was extracted with two portions of dichloromethane. The organic extracts were dried over sodium sulfate, concentrated, and the
20 residue was chromatographed on silica gel using 1:1 hexanes/ethyl acetate as eluant to provide **241C** (230 mg) as a yellow oil.

- D. N-(3,4-Dimethyl-5-isoxazolyl)-2'-(2-fluoroethyl)-4'-(hydroxymethyl)-N-[(2-trimethylsilyl)ethoxymethyl] [1,1'-biphenyl]-2-sulfonamide

25 **241C** (230 mg, 0.41 mmol) was treated with DIBAL-H according to the procedure of Example 230, step E, to provide crude **241D** (320 mg) as a yellow oil

- E. N-(3,4-Dimethyl-5-isoxazolyl)-2'-(2-fluoroethyl)-4'-(methanesulfonyloxy)methyl-N-[(2-trimethylsilyl)ethoxymethyl] [1,1'-biphenyl]-2-sulfonamide
- 30

241D (320 mg) was converted to the corresponding methanesulfonate ester according to General Method 3. The entire crude product **241E** was used directly in the next step.

- 5 F. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-(2-fluoroethyl)-N-[(2-trimethylsilyl)ethoxymethyl] [1,1'-biphenyl]-2-sulfonamide

241E was used to alkylate 2-butyl-1,3-diazaspiro[4.4]non-1-en-4-one according to General Method 4, to provide **241F** (300 mg) as a yellow oil,
10 which was used without further purification.


- G. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethylisoxazol-5-yl)-2'-(2-fluoroethyl) [1,1'-biphenyl]-2-sulfonamide

15 Crude **241F** (300 mg) was deprotected according to General Method 8 (ethanol). The crude product was purified by silica gel chromatography using hexanes/ethyl acetate as eluant provided the title compound (16 mg) as a tan solid; LRMS m/z 581 (ESI+ mode); HPLC retention time 2.05 min (Method H); HPLC purity 97%.

20

Example 242

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethylisoxazol-5-yl)-2'-(2-hydroxyethyl)[1,1'-biphenyl]-2-sulfonamide



25

A. N-(3,4-Dimethyl-5-isoxazolyl)-4'-methoxycarbonyl-2'-[2-[(tetrahydro-2H-pyran-2-yl)oxy]ethyl]-N-[(2-trimethylsilyl)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

Pyridinium p-toluenesulfonate (2 mg) was added to a solution of
5 **241B** (1.0 g, 1.8 mmol) and 3,4-dihydro-2H-pyran (0.48 ml, 5.4 mmol) in dichloromethane (4 ml) at 0 °C. After 48h aqueous sodium bicarbonate solution was added, the layers were separated, and the organic layer was dried over sodium sulfate and concentrated. The residue was
10 **242A** (320 mg) as a slightly yellow oil.

B. N-(3,4-Dimethyl-5-isoxazolyl)-4'-(hydroxymethyl)-2'-[2-[(tetrahydro-2H-pyran-2-yl)oxy]ethyl]-N-[(2-trimethylsilyl)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

15 **242A** (320 mg, 0.50 mmol) was treated with DIBAL-H according to the procedure of Example 230, step E, to provide crude **242B** (250 mg) as a yellow oil.

C. N-(3,4-Dimethyl-5-isoxazolyl)-4'-(methanesulfonyloxy)methyl-2'-[2-[(tetrahydro-2H-pyran-2-yl)oxy]ethyl]-N-[(2-trimethylsilyl)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

20 **242B** (250 mg) was converted to the corresponding methanesulfonate ester according to General Method 3. The entire crude product **242C** was used in the next reaction step.

25

D. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-[2-[(tetrahydro-2H-pyran-2-yl)oxy]ethyl]-N-[(2-trimethylsilyl)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

242C was used to alkylate 2-butyl-1,3-diazaspiro[4.4]non-1-en-4-one
30 according to General Method 4. The crude **242D** was used without further purification.


E. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-(2-hydroxyethyl)[1,1'-biphenyl]-2-sulfonamide

5 **242D** was deprotected according to General Method 8 (ethanol). The crude product was purified by silica gel chromatography using 3% methanol in dichloromethane as eluant to provide the title compound (45 mg) as a white solid following lyophilization: MS m/e 579 (ESI+ mode); HPLC retention time 1.42 min (Method H); HPLC purity >98%.

10

Example 243

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethylisoxazol-5-yl)-2'-(3-methylbutyl)[1,1'-biphenyl]-2-sulfonamide



15

A. 4-Bromo-3-(3-methyl-1-butenyl)benzonitrile

20 **2A** (2.0 g, 9.5 mmol) was reacted with isobutyltriphenylphosphonium bromide following the procedure of Example 27, step A. The crude residue was chromatographed on silica gel using 9:1 hexanes/ethyl acetate to afford **243A** (2.2 g, 91%) as an E/Z mixture.

25 B. 4-Bromo-3-(3-methylbutyl)benzonitrile

A mixture of **243A** (2.2 g) and 220 mg of PtO₂ in 30 ml EtOH was hydrogenated at 35 PSI for 20 min. Filtration and concentration gave 1.9 g of **243B** (86%).

C. 4-Bromo-3-(3-methylbutyl)benzaldehyde

243B (1.9 g) was treated with DIBAL-H according to General Method 14 to provide crude **243C** (810 mg, 43%) as an oil.

5

D. N-(3,4-Dimethyl-5-isoxazolyl)-4'-formyl-2'-(3-methylbutyl)-N-[(2-methoxyethoxy)methyl][1,1'-biphenyl]-2-sulfonamide

243C (810 mg) was subjected to Suzuki coupling according to General Method 1 to provide **243D** as a crude oil.

10

E. N-(3,4-Dimethyl-5-isoxazolyl)-4'-hydroxymethyl-2'-(3-methylbutyl)-N-[(2-methoxyethoxy)methyl][1,1'-biphenyl]-2-sulfonamide

243D (entire sample) was reduced with sodium borohydride in methanol according to General Method 11 to provide **243E** (490 mg, 30% from **243C**) as an oil following silica gel chromatography using hexanes/ethyl acetate as eluant.

15

F. 4'-Bromomethyl-N-(3,4-dimethyl-5-isoxazolyl)-2'-(3-methylbutyl)-N-[(2-methoxyethoxy)methyl][1,1'-biphenyl]-2-sulfonamide

243E (490 mg) was converted to the corresponding bromide according to General Method 2 to provide **243F** (430 mg, 78%) as an oil following silica gel chromatography using hexanes/ethyl acetate as eluant.

20

G. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-(3-methylbutyl)-N-[(2-methoxyethoxy)methyl]-N-(3,4-dimethyl-5-isoxazolyl)[1,1'-biphenyl]-2-sulfonamide

243F (430 mg) was used to alkylate 2-butyl-1,3-diazaspiro[4.4]non-1-en-4-one according to General Method 4. **243G** (300 mg, 58%) was produced as an oil following silica gel chromatography using hexanes/ethyl acetate as eluant.

25

30


H. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethylisoxazol-5-yl)-2'-(3-methylbutyl)[1,1'-biphenyl]-2-sulfonamide

243G was deprotected according to General Method 7. The crude product was purified by reverse-phase preparative HPLC to provide the title compound (165 mg, 63%) as a white solid: mp 50-53 °C; MS m/e 605 (ESI+ mode); HPLC retention time 27.42 min (Method B); HPLC purity >97%.

10

Example 244

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethylisoxazol-5-yl)-2'-(2-methylpropyl)[1,1'-biphenyl]-2-sulfonamide



15

A. 4-[(2-Methyl-2-propenyl)oxy]benzonitrile

A mixture of 4-cyanophenol (9.0 g, 75 mmol), potassium carbonate (21 g, 150 mmol), and DMF (50 ml) was treated with methallyl bromide (7.8 ml, 77 mmol) at 0 °C. The mixture was stirred at RT for 16 h, then was added to water and extracted with ethyl acetate (3 X 100 ml). The combined organic extracts were washed with water and dried and evaporated to give **244A** (13 g, 99%) as an oil.

25 B. 4-Cyano-2-(2-methyl-2-propenyl)phenol

A solution of **244A** (13 g, 75 mmol) and BHT (165 mg) in 1,2,4-trichlorobenzene (40 ml) was heated at 200 °C for 5 days. The mixture was cooled and diluted with ethyl acetate, then the organic layer was

extracted with 10% NaOH solution (3 X 200ml). The aqueous phase was made acidic with hydrochloric acid and extracted with ether (3 X 100ml). The combined ether extracts were washed with water and dried and evaporated. The residue was chromatographed on silica gel using 9:1
5 hexanes/ethyl acetate to afford **244C** (5.8 g, 45%) as a solid.

C. 4-Cyano-2-(2-methylpropyl)phenol

A mixture of **244B** (1.1 g) and 110 mg of PtO₂ in 20 ml EtOH was hydrogenated at 35 PSI for 15 min. Filtration and concentration gave
10 0.73 g of **244C** (66%).

D. 4-Hydroxy-3-(2-methylpropyl)benzaldehyde

244C (0.73 g) was treated with DIBAL-H according to General Method 14 to provide crude **244D** (530 mg, 72%) as an oil.

15

E. 3-(2-Methylpropyl)-4-(trifluoromethanesulfonyloxy)benzaldehyde

244D (530 mg) was converted to the corresponding trifluoromethanesulfonate ester according to the procedure of Example 229, step E. The residue was chromatographed on silica gel using 4:1
20 hexanes/ethyl acetate to give **244E** (295 mg, 33%) as an oil.

F. N-(3,4-Dimethyl-5-isoxazolyl)-4'-formyl-2'-(2-methylpropyl)-N-[(2-methoxyethoxy)methyl][1,1'-biphenyl]-2-sulfonamide

Suzuki coupling of **244E** (295 mg) according to the procedure of
25 Example 229, step F, provided **244F** (170 mg, 36%) as a yellow solid following silica gel chromatography using hexanes/ethyl acetate as eluant.

G. N-(3,4-Dimethyl-5-isoxazolyl)-4'-hydroxymethyl-2'-(2-methylpropyl)-N-[(2-methoxyethoxy)methyl][1,1'-biphenyl]-2-sulfonamide

244F (170 mg) was reduced with sodium borohydride in methanol according to General Method 11 to provide **244G** (67 mg, 39%) as an oil.

30

H. 4'-Bromo-methyl-N-(3,4-dimethyl-5-isoxazolyl)-2'-(2-methylpropyl)-N-[(2-methoxyethoxy)methyl][1,1'-biphenyl]-2-sulfonamide

244G (67 mg) was converted to the corresponding bromide

5 according to General Method 2 to provide **244H** (42 mg, 56%) as an oil.

I. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-(2-methylpropyl)-N-[(2-methoxyethoxy)methyl]-N-(3,4-dimethyl-5-isoxazolyl)[1,1'-biphenyl]-2-sulfonamide


10 **244H** (42 mg) was used to alkylate 2-butyl-1,3-diazaspiro[4.4]non-1-en-4-one according to General Method 4. **244I** (36 mg, 45%) was produced as an oil following silica gel chromatography using hexanes/ethyl acetate as eluant.

15 J. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-(2-methylpropyl)-N-(3,4-dimethyl-5-isoxazolyl)[1,1'-biphenyl]-2-sulfonamide

244I (36 mg) was deprotected according to General Method 7. The crude product was purified by reverse-phase preparative HPLC to provide
20 the title compound (25 mg, 86%) as a white solid: mp 58-61 °C; MS m/e 591 (ESI+ mode); HPLC retention time 28.21 min (Method B); HPLC purity >98%.

Example 245

25 4'-[[2-(3,3-Difluorobutyl)-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl]methyl]-N-(3,4-dimethylisoxazol-5-yl)-2'-(ethoxymethyl)[1,1'-biphenyl]-2-sulfonamide



A. Methyl 1-[(3,3-difluorobutanoyl)aminocyclopentane-1-carboxylate

A solution of 4,4-Difluoropentanoic acid (600 mg, 4.4 mmol,
5 prepared according to Larsson, U.; Carlson, R.; Leroy, J. *Acta Chem. Scand.* **1993**, 47, 380-90) in dichloromethane (10 ml) was treated at RT with oxalyl chloride (4.4 ml of a 2.0 M solution in dichloromethane, 8.8 mmol) and DMF (10 µl). After 20 min the mixture was evaporated and 10 ml fresh dichloromethane was added. The mixture was cooled to 0 °C and
10 methyl 1-aminocyclopentane-1-carboxylate hydrochloride (1.6 g, 8.8 mmol) was added, followed by triethylamine (3.6 ml, 26 mmol) and DMAP (10 mg). The mixture was stirred at RT for 3 hours. Aqueous sodium bicarbonate solution was added and the mixture was extracted three times with dichloromethane. The combined organic extracts were dried over
15 sodium sulfate and concentrated. The residue was chromatographed on silica gel using 1:2 hexanes/ethyl acetate as eluant to give **245A** (520 mg, 46%) as an orange oil.

B. 2-(3,3-Difluorobutyl)-1,3-diazaspiro[4.4]non-1-en-4-one

20 **245A** (520 mg, 2.0 mmol) was treated according to the procedure of Example 22, step B. The crude residue was chromatographed on silica gel using 1:3 hexanes/ethyl acetate as eluant to yield **245B** (150 mg, 33%) as a yellow oil.

25 C. 4'-[[2-(3,3-Difluorobutyl)-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl]methyl]-N-(3,4-dimethylisoxazol-5-yl)-2'-(ethoxymethyl)-N-(2-methoxyethoxy)methyl[1,1'-biphenyl]-2-sulfonamide

245B (75 mg, 0.42 mmol) was alkylated with 4'-bromomethyl-N-(3,4-dimethyl-5-isoxazolyl)-2'-ethoxymethyl-N-(2-
30 methoxyethoxy)methyl[1,1'-biphenyl]-2-sulfonamide (prepared as


described in Example 226) according to General Method 4. Crude **245C** (220 mg) was used without further purification.

D. 4'-[[2-(3,3-Difluorobutyl)-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl]methyl]-N-(3,4-dimethylisoxazol-5-yl)-2'-(ethoxymethyl)[1,1'-biphenyl]-2-sulfonamide

245C (220 mg) was deprotected according to General Method 8 (EtOH). The crude product was purified by preparative thin-layer silica gel chromatography using 1:1 hexanes/acetone to provide the title compound (57 mg) as a white lyophilized powder; MS m/e 629 (ESI+ mode); HPLC retention time 3.79 min (Method A); HPLC purity 96%.

Example 246

N-(3,4-Dimethyl-5-isoxazolyl)-4'-[[[(3-methoxy-2,6-dimethyl-4-pyridinyl)oxy]methyl]-2'-(3,3,3-trifluoropropyl)[1,1'-biphenyl]-2-sulfonamide



A. N-(3,4-Dimethyl-5-isoxazolyl)-4'-[[[(3-methoxy-2,6-dimethyl-4-pyridinyl)oxy]methyl]-2'-(3,3,3-trifluoropropyl)-N-[(2-trimethylsilyl)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide


229H (400 mg, 0.60 mmol) was used to alkylate 3-methoxy-2,6-dimethyl-4-(4H)-pyridinone according to General Method 22. The crude product was purified by silica gel chromatography using 1:2 hexanes/ethyl acetate as eluant to provide **246A** (130 mg) as a slightly yellow oil.

B. N-(3,4-Dimethyl-5-isoxazolyl)-4'-[[[(3-methoxy-2,6-dimethyl-4-pyridinyl)oxy]methyl]-2'-(3,3,3-trifluoropropyl)[1,1'-biphenyl]-2-sulfonamide

5 **246A** (130 mg, 0.18 mmol) was deprotected according to General Method 8 (EtOH). The crude product was purified by silica gel chromatography using 5% methanol in chloroform as eluant to provide the title compound (82 mg) as a pale orange powder after lyophilization; MS m/e 590 (ESI+ mode); HPLC retention time 3.35 min (Method A); HPLC
10 purity 97%.

Example 247

15 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethylisoxazol-5-yl)-2'-[(1,1-dimethylethoxy)methyl][1,1'-biphenyl]-2-sulfonamide



20 A. Methyl 4-bromo-3-[(1,1-dimethylethoxy)methyl]benzoate

To a solution of **240A** (4.90 g, 20 mmol) in 40 ml cyclohexane and 20 ml dichloromethane was added t-butyl 2,2,2-trichloroacetimidate (4.81 g, 22 mmol) followed by 0.4 ml boron trifluoride diethyl etherate. The mixture was stirred at RT for 4 hours. 1 g solid sodium bicarbonate was
25 added and the mixture was stirred for 10 min. The mixture was chromatographed directly on silica gel eluting with 20:1 hexanes/ethyl acetate to give compound **247A** as an oil (5.33 g, 88%).

B. N-(3,4-Dimethyl-5-isoxazolyl)-2'-[(1,1-dimethylethoxy)methyl]-4'-methoxycarbonyl-N-[(2-trimethylsilyl)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

5 **247A** (5.3 g, 17.5 mmol) was subjected to Suzuki coupling according to General Method 1. Silica gel chromatography using hexanes/ethyl acetate as eluant provided **247B** (6.4 g, 88%) as an oil.

C. N-(3,4-Dimethyl-5-isoxazolyl)-2'-[(1,1-dimethylethoxy)methyl]-4'-hydroxymethyl-N-(2-trimethylsilyl)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

10

247B (6.4 g, 10.7 mmol) was reduced with DIBAL-H according to the procedure of Example 230, step E, to provide crude **247C** (5.5 g, 89%) as an orange oil.

15 D. 4'-Bromomethyl-N-(3,4-dimethyl-5-isoxazolyl)-2'-[(1,1-dimethylethoxy)methyl]-N-[(2-trimethylsilyl)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

247C (5.5 g, 9.6 mmol) was converted to the corresponding bromide according to General Method 2, providing **247D** (5.6 g, 92%) as a yellow oil


20 following silica gel chromatography using hexanes/ethyl acetate as eluant.

E. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-[(1,1-dimethylethoxy)methyl]-N-[(2-trimethylsilyl)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

25 **247D** (380 mg, 0.60 mmol) was used to alkylate 2-butyl-1,3-diazaspiro[4.4]non-1-en-4-one according to General Method 4. The crude residue was chromatographed on silica gel using hexanes/ethyl acetate as eluant to provide **247E** (410 mg, 92%) as a slightly yellow oil.

E. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-[(1,1-dimethylethoxy)methyl][1,1'-biphenyl]-2-sulfonamide


- 247E (410 mg, 0.55 mmol) was deprotected with TBAF according to General Method 10. The crude product was purified by reverse-phase preparative HPLC to provide the title compound (230 mg, 66%) as a white solid: MS m/e 621 (ESI+ mode); HPLC retention time 20.97 min (Method E); HPLC purity 96%.

E X A M P L E	S T R U C T U R E	N A M E	S T A R T I N G M A T E R I A L	General Methods Applied (yield, %)	M/z (MH) ⁺	HPLC % Puri- ty	HPLC ret time, min (meth- od)
248		1-[[2'-[[[(3,4-dimethyl-5-isoxazolyl)-amino]sulfonyl-2-methoxymethyl][1,1'-biphenyl]-4-yl)methyl]-4-ethyl-2-propyl-1H-imidazole-5-carboxamide	P2	22 (39); 8, EtOH, 15 (96); 12 (38)	566	95	2.79 (C)

249		1-[[2'-[[[(3,4-dimethyl-5-isoxazolyl)-amino]sulfonyl-2-methoxymethyl][(1,1'-biphenyl)-4-yl]methyl]-4-ethyl-N-methyl-2-propyl-1H-imidazole-5-carboxamide	P2	22 (39); 8, EtOH, 15 (96); 12 (34)	580	96	2.90 (C)
250		1-[[2'-[[[(3,4-dimethyl-5-isoxazolyl)-amino]sulfonyl-2-methyl][(1,1'-biphenyl)-4-yl]methyl]-4-ethyl-2-propyl-1H-imidazole-5-carboxamide	P13	22 (56); 8, EtOH, 15 (94); 12 (38)	536	95	2.90 (C)
251		1-[[2'-[[[(3,4-dimethyl-5-isoxazolyl)-amino]sulfonyl-2-methyl][(1,1'-biphenyl)-4-yl]methyl]-4-ethyl-N-methyl-2-propyl-1H-imidazole-5-carboxamide	P13	22 (56); 8, EtOH, 15 (94); 12 (37)	550	97	2.99 (C)

Example 252

N-(4,5-Dimethyl-3-isoxazolyl)-2'-ethoxymethyl-4'-[[[(3-methoxy-2,6-dimethyl-4-pyridinyl)oxy]methyl] [1,1'-biphenyl]-2-sulfonamide



A. 4-Bromo-3-(ethoxymethyl)benzonitrile

P2A (8.7 g, 32 mmol) was treated with sodium hydride and ethanol according to General Method 4, producing crude **252A** (5.8 g, 77%) as an oil.

B. 4-Bromo-3-(ethoxymethyl)benzaldehyde

Crude **252A** (5.8 g, 24 mmol) was reduced with DIBAL-H according to General Method 14, yielding **252B** (4.7 g, 80%) as an oil following silica gel chromatography with hexanes/ethyl acetate as eluant.

C. N-(4,5-Dimethyl-3-isoxazolyl)-2'-ethoxymethyl-4'-formyl-N-[(2-methoxyethoxy)methyl] [1,1'-biphenyl]-2-sulfonamide

252B (800 mg, 3.3 mmol) was subjected to Suzuki coupling with [2-[[[(4,5-dimethyl-3-isoxazolyl)](2-methoxyethoxy)methyl]amino]-sulfonyl]phenyl]boronic acid according to General Method 1. **252C** (1.2 g, 70%) was obtained as an orange oil following silica gel chromatography using hexanes/ethyl acetate as eluant.

D. N-(4,5-Dimethyl-3-isoxazolyl)-2'-ethoxymethyl-4'-hydroxymethyl-N-[(2-methoxyethoxy)methyl] [1,1'-biphenyl]-2-sulfonamide

252C (1.2 g, 2.3 mmol) was reduced with sodium borohydride in methanol according to General Method 11 to provide crude **252D**. This material was used without further purification.

E. 4'-Bromomethyl-N-(4,5-dimethyl-3-isoxazolyl)-2'-ethoxymethyl-N-[(2-methoxyethoxy)methyl][1,1'-biphenyl]-2-sulfonamide

252D (entire sample) was converted to the corresponding bromide according to General Method 2. **252E** (1.1 g, 84% over two steps) was obtained as an oil following silica gel chromatography using hexanes/ethyl acetate as eluant.

F. N-(4,5-Dimethyl-3-isoxazolyl)-2'-ethoxymethyl-4'-[[[(3-methoxy-2,6-dimethyl-4-pyridinyl)oxy]methyl]- N-[(2-methoxyethoxy)methyl] [1,1'-biphenyl]-2-sulfonamide


5 **252E** (360 mg, 0.63 mmol) was used to alkylate 3-methoxy-2,6-dimethyl-4-(1H)-pyridinone according to General Method 22. **252F** (310 mg, 76%) was produced as an oil following silica gel chromatography using hexanes/ethyl acetate as eluant.

10 G. N-(4,5-Dimethyl-3-isoxazolyl)-2'-ethoxymethyl-4'-[[[(3-methoxy-2,6-dimethyl-4-pyridinyl)oxy]methyl] [1,1'-biphenyl]-2-sulfonamide
 252F (310 mg, 0.49 mmol) was deprotected according to General Method 7. The crude product was purified by silica gel column chromatography using methanol/dichloromethane as eluant to provide the
 15 title compound (240 mg, 91%) as a light yellow solid: mp 78-82 °C; MS m/e 552 (ESI+ mode); HPLC retention time 12.42 min (Method E); HPLC purity 97%.

20

Example 253

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-(2-fluoroethoxymethyl) [1,1'-biphenyl]-2-sulfonamide



25

A. 4-Bromo-3-[(2-fluoroethoxy)methyl]benzonitrile

P2A (4.1 g, 15 mmol) was treated with sodium hydride and 2-fluoroethanol according to General Method 4. Water was added to the reaction mixture to precipitate **253A** (2.9 g, 75%) as a brown solid.

5

B. 4-Bromo-3-[(2-fluoroethoxy)methyl]benzaldehyde

253A (2.9 g, 11 mmol) was reduced with DIBAL-H according to General Method 14, yielding **253B** (2.5 g) as an oil, which was used without further purification.

10

C. N-(4,5-Dimethyl-3-isoxazolyl)-2'-[(2-fluoroethoxy)methyl]-4'-formyl-N-[(2-methoxyethoxy)methyl] [1,1'-biphenyl]-2-sulfonamide

253B (2.5 g) was subjected to Suzuki coupling with [2-[(4,5-dimethyl-3-isoxazolyl)(2-methoxyethoxy)methyl]amino]-

15 sulfonylphenyl]boronic acid according to General Method 1. **253C** (1.4 g, 23% over two steps) was obtained as an oil following silica gel chromatography using hexanes/ethyl acetate as eluant.

D. N-(4,5-Dimethyl-3-isoxazolyl)-2'-[(2-fluoroethoxy)methyl]-4'-hydroxymethyl-N-[(2-methoxyethoxy)methyl] [1,1'-biphenyl]-2-sulfonamide

20

253C (1.4 g, 2.6 mmol) was reduced with sodium borohydride in methanol according to General Method 11 to provide **253D** as an oil. This material was used without further purification.

25

E. 4'-Bromomethyl-N-(4,5-dimethyl-3-isoxazolyl)-2'-[(2-fluoroethoxy)methyl]-N-[(2-methoxyethoxy)methyl][1,1'-biphenyl]-2-sulfonamide

253D (entire sample) was converted to the corresponding bromide according to General Method 2. **253E** (1.3 g, 80% over two steps) was

30

obtained as an oil following silica gel chromatography using hexanes/ethyl acetate as eluant.

F. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-[(2-fluoroethoxy)methyl]-N-[(2-methoxyethoxy)methyl] [1,1'-biphenyl]-2-sulfonamide


253E (430 mg) was used to alkylate 2-butyl-1,3-diazaspiro[4.4]non-1-en-4-one according to General Method 4. **253F** (400 mg, 78%) was produced as an oil following silica gel chromatography using hexanes/ethyl acetate as eluant.

G. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-[(2-fluoroethoxy)methyl] [1,1'-biphenyl]-2-sulfonamide

253F (400 mg, 0.57 mmol) was deprotected according to General Method 7. The crude product was purified by silica gel column chromatography using methanol/dichloromethane as eluant to provide the title compound (310 mg, 89%) as a white solid: mp 68-73 °C; MS m/e 611 (ESI+ mode); HPLC retention time 16.07 min (Method E); HPLC purity >97%.

Example 254

N-(4,5-Dimethyl-3-isoxazolyl)-2'-[(2-fluoroethoxy)methyl]-4'-[(3-methoxy-2,6-dimethyl-4-pyridinyl)oxy]methyl] [1,1'-biphenyl]-2-sulfonamide



B. N-(4,5-Dimethyl-3-isoxazolyl)-2'-[(2-fluoroethoxy)methyl]-4'-[[[(3-methoxy-2,6-dimethyl-4-pyridinyl)oxy]methyl]-N-[(2-methoxyethoxy)methyl] [1,1'-biphenyl]-2-sulfonamide


5 **253E** (450 mg, 0.77 mmol) was used to alkylate 3-methoxy-2,6-dimethyl-4-(1H)-pyridinone according to General Method 22. The crude product was purified by silica gel chromatography using hexanes/ethyl acetate as eluant to provide **254A** (380 mg, 76%) as a slightly yellow oil.

10 B. N-(4,5-Dimethyl-3-isoxazolyl)-2'-[(2-fluoroethoxy)methyl]-4'-[[[(3-methoxy-2,6-dimethyl-4-pyridinyl)oxy]methyl] [1,1'-biphenyl]-2-sulfonamide

15 **254A** (380 mg, 0.58 mmol) was deprotected according to General Method 7. The crude product was purified by silica gel chromatography using methanol/chloroform as eluant to provide the title compound (310 mg, 93%) as a light yellow solid; mp 81-86 °C; MS m/e 570 (ESI+ mode); HPLC retention time 10.54 min (Method E); HPLC purity >97%.

Example 255

20 N-(3,4-Dimethyl-5-isoxazolyl)-2'-[(2-fluoroethoxy)methyl]-4'-[[[(3-methoxy-2,6-dimethyl-4-pyridinyl)oxy]methyl] [1,1'-biphenyl]-2-sulfonamide



25

A. N-(3,4-Dimethyl-5-isoxazolyl)-2'-[(2-fluoroethoxy)methyl]-4'-formyl-N-[[[(2-trimethylsilyl)ethoxy]methyl] [1,1'-biphenyl]-2-sulfonamide

253B (1.3 g, 4.9 mmol) was subjected to Suzuki coupling with [2-
[[[3,4-dimethyl-5-isoxazolyl]][(2-trimethylsilyl)ethoxy)methyl]amino]-
sulfonyl]phenyl]boronic acid according to General Method 1. **255A** (2.5 g,
90%) was obtained as an oil following silica gel chromatography using
5 hexanes/ethyl acetate as eluant.

B. N-(3,4-Dimethyl-5-isoxazolyl)-2'-[(2-fluoroethoxy)methyl]-4'-
hydroxymethyl-N-[[[(2-trimethylsilyl)ethoxy)methyl] [1,1'-biphenyl]-
2-sulfonamide

10 **255A** (2.5 g, 4.4 mmol) was reduced with sodium borohydride in
methanol according to General Method 11 to provide **255B** as an oil. This
material was used without further purification.

C. 4'-Bromomethyl-N-(3,4-dimethyl-5-isoxazolyl)-2'-[(2-
15 fluoroethoxy)methyl]-N-[[[(2-trimethylsilyl)ethoxy)methyl] [1,1'-
biphenyl]-2-sulfonamide

255B (entire sample) was converted to the corresponding bromide
according to General Method 2, furnishing **255C** (2.1 g, 76% over two
steps) as an oil following silica gel chromatography using hexanes/ethyl
20 acetate as eluant.

D. N-(3,4-dimethyl-5-isoxazolyl)-2'-[(2-fluoroethoxy)methyl]-4'-[[[3-
methoxy-2,6-dimethyl-4-pyridinyl]oxy)methyl]-N-[[[(2-
trimethylsilyl)ethoxy)methyl] [1,1'-biphenyl]-2-sulfonamide


25 **255C** (590 mg, 0.93 mmol) was used to alkylate 3-methoxy-2,6-
dimethyl-4-(1H)-pyridinone according to General Method 22. **255D** (290
mg, 44%) was produced as an oil following silica gel chromatography using
hexanes/ethyl acetate as eluant.

E. N-(3,4-Dimethyl-5-isoxazolyl)-2'-[(2-fluoroethoxy)methyl]-4'-[[[(3-methoxy-2,6-dimethyl-4-pyridinyl)oxy]methyl] [1,1'-biphenyl]-2-sulfonamide

255D was deprotected according to General Method 7. The crude product was purified by silica gel column chromatography using methanol/dichloromethane as eluant to provide the title compound (180 mg, 75%) as a white solid: mp 90-95 °C; MS m/e 570 (ESI+ mode); HPLC retention time 10.71 min (Method E); HPLC purity >97%.

Example 256

N-(3,4-Dimethyl-5-isoxazolyl)-2'-ethoxymethyl-4'-[[4-oxo-2-(3,3,3-trifluoropropyl)-1,3-diazaspiro[4.4]non-1-en-3-yl]methyl] [1,1'-biphenyl]-2-sulfonamide



A. Methyl 1-[(4,4,4-trifluorobutanoyl)amino]cyclopentane-1-carboxylate

A solution of 4,4,4-trifluorobutanoic acid (5.0 g, 35 mmol) in dichloromethane (90 ml) was treated at 0 °C with oxalyl chloride (26 ml of a 2.0 M solution in dichloromethane, 52 mmol) and DMF (10 µl). After 20 min the mixture was concentrated to a volume of about 10 ml and 90 ml fresh dichloromethane was added. The mixture was cooled to 0 °C and methyl 1-aminocyclopentane-1-carboxylate hydrochloride (12.6 g, 70 mmol) was added, followed by triethylamine (30 ml, 210 mmol) and DMAP (10 mg). The mixture was stirred at RT for 60 hours. Aqueous sodium

bicarbonate solution was added and the mixture was extracted three times with dichloromethane. The combined organic extracts were dried over sodium sulfate and concentrated. The residue was chromatographed on silica gel using 1:1 hexanes/ethyl acetate as eluant to give **256A** (3.1 g, 33%) as an orange oil.

B. 2-(3,3,3-Trifluoropropyl)-1,3-diazaspiro[4.4]non-1-en-4-one
256A (3.1 g, 12 mmol) was treated according to the procedure of Example 22, step B. Evaporation of the crude extract provided **256B** (2.0 g, 71%) as a white solid, which was used without further purification.

C. N-(3,4-Dimethyl-5-isoxazolyl)-2'-ethoxymethyl-4'-formyl-N-[[2-trimethylsilyl)ethoxy)methyl] [1,1'-biphenyl]-2-sulfonamide
252B (7.6 g, 31 mmol) was subjected to Suzuki coupling with [2-[[[(3,4-dimethyl-5-isoxazolyl)[[(2-trimethylsilyl)ethoxy)methyl]amino]-sulfonyl]phenyl]boronic acid according to General Method 1. **256C** (12.3 g, 72%) was obtained as an oil following silica gel chromatography using hexanes/ethyl acetate as eluant.

D. N-(3,4-Dimethyl-5-isoxazolyl)-2'-ethoxymethyl-4'-hydroxymethyl-N-[[2-trimethylsilyl)ethoxy)methyl] [1,1'-biphenyl]-2-sulfonamide
256C (1.0 g, 1.8 mmol) was reduced with sodium borohydride in ethanol according to General Method 11 to provide **256D** (0.97 g, 95%) as a crude brown oil, which was used without further purification.

E. N-(3,4-Dimethyl-5-isoxazolyl)-2'-ethoxymethyl-4'-(methanesulfonyloxy)methyl-N-[[2-trimethylsilyl)ethoxy)methyl] [1,1'-biphenyl]-2-sulfonamide
256D (490 mg) was converted to the corresponding mesylate according to General Method 3 to provide **256E** (560 mg) as crude orange oil.

F. N-(3,4-dimethyl-5-isoxazolyl)-2'-ethoxymethyl-4'-[[4-oxo-2-(3,3,3-trifluoropropyl)-1,3-diazaspiro[4.4]non-1-en-3-yl]methyl]-N-[(2-trimethylsilyl)ethoxy]methyl] [1,1'-biphenyl]-2-sulfonamide

5 **256E** (560 mg) was used to alkylate **256B** (270 mg) according to General Method 4. **256F** (570 mg) was produced as a crude brown oil, which was used without further purification.


G. N-(3,4-Dimethyl-5-isoxazolyl)-2'-ethoxymethyl-4'-[[4-oxo-2-(3,3,3-trifluoropropyl)-1,3-diazaspiro[4.4]non-1-en-3-yl]methyl] [1,1'-biphenyl]-2-sulfonamide

10 **256F** (570 mg) was deprotected according to General Method 8 (EtOH). The crude product was purified by silica gel column chromatography using 2:1 hexanes/ethyl acetate as eluant to provide the
15 title compound (92 mg, 8% from **256D**) as a white solid: MS m/e 633 (ESI+ mode); HPLC retention time 3.96 min (Method C); HPLC purity 97%

20

Example 257

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-ethyl [1,1'-biphenyl]-2-sulfonamide



25

A. Methyl 4-bromo-3-ethylbenzoate

Methylolithium (57 ml of a 1.4 M solution in ether, 80 mmol) was added dropwise to a suspension of copper(I) iodide (76 g, 40 mmol) in ether (20 ml) at 0 °C. The mixture was stirred for 20 min at 0 °C. Solid **232A** (12.3 g, 40 mmol) was added in portions over 45 min, after which THF (50 ml) was added. The heterogeneous mixture was stirred at 0 °C for 1 h, then aqueous ammonium chloride and aqueous ammonium hydroxide were added. The mixture was extracted with ethyl acetate, and the combined organic layers were concentrated. The residue was taken up in chloroform and partitioned against aqueous ammonium chloride. The chloroform layer was dried over sodium sulfate and concentrated, and the residue was chromatographed on silica gel, eluting with 9:1 hexanes/ethyl acetate, to provide **257A** (3.2 g, 32%) as a yellow oil.

B. N-(3,4-Dimethyl-5-isoxazolyl)-2'-ethyl-4'-methoxycarbonyl-N-(((2-trimethylsilyl)ethoxy)methyl) [1,1'-biphenyl]-2-sulfonamide

257A (1.4 g, 5.6 mmol) was subjected to Suzuki coupling with [2-(((3,4-dimethyl-5-isoxazolyl) (((2-trimethylsilyl)ethoxy)methyl)amino)sulfonyl]phenyl]boronic acid according to General Method 1. Silica gel chromatography provided **257B** (3.2 g) as an oil, contaminated with byproducts deriving from the boronic acid.

C. N-(3,4-Dimethyl-5-isoxazolyl)-2'-ethyl-4'-hydroxymethyl-N-(((2-trimethylsilyl)ethoxy)methyl) [1,1'-biphenyl]-2-sulfonamide

257B (3.2 g) was treated with DIBAL-H according to the procedure of Example 230, step E, to provide **257C** (4.0 g) as a crude oil.

D. N-(3,4-Dimethyl-5-isoxazolyl)-2'-ethyl-4'-(methanesulfonyloxy)methyl-N-(((2-trimethylsilyl)ethoxy)methyl) [1,1'-biphenyl]-2-sulfonamide

257C (4.0 g) was converted to the corresponding mesylate according to General Method 3 to provide **257D** (3.7 g) as crude orange oil.

E. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-ethyl- N-[(2-trimethylsilyl)ethoxy)methyl][1,1'-biphenyl]-2-sulfonamide


5 **257D** (0.92 g) was used to alkylate 2-butyl-1,3-diazaspiro[4.4]non-1-en-4-one according to General Method 4. **257E** (760 mg) was produced as a colorless oil following silica gel chromatography using 2:1 hexanes/ethyl acetate as eluant.

10 F. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-ethyl [1,1'-biphenyl]-2-sulfonamide

257D (760 mg) was deprotected according to General Method 8 (EtOH). The crude product was purified by reverse-phase preparative HPLC to provide the title compound (200 mg) as a white solid following
15 lyophilization: MS m/e 563 (ESI+ mode); HPLC retention time 3.69 min (Method C); HPLC purity >98%

Example 258

20 (+/-)-4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-(1-hydroxyethyl) [1,1'-biphenyl]-2-sulfonamide



25

A. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-formyl-N-[(2-trimethylsilyl)ethoxymethyl][1,1'-biphenyl]-2-sulfonamide

23A (600 mg, 1.0 mmol) was used to alkylate 2-butyl-1,3-diazaspiro[4.4]non-1-en-4-one according to General Method 4. **258A** (510 mg, 74%) was produced as a yellow oil following silica gel chromatography using 1:1 hexanes/ethyl acetate as eluant.

5

B. (+/-)-4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-(1-hydroxyethyl)-N-[(2-trimethylsilyl)ethoxymethyl] [1,1'-biphenyl]-2-sulfonamide

A solution of **258A** (260 mg, 0.37 mmol) in dry ether (1 ml) was treated dropwise at -78 °C with methylmagnesium bromide (3.0 M solution in ether, 0.25 ml, 0.74 mmol). The mixture was allowed to warm to 0 °C and was then quenched with aqueous ammonium chloride. The mixture was extracted with ether. The combined organic extracts were dried over magnesium sulfate and concentrated to provide **258B** (210 mg) as a colorless oil, which was used without further purification.

10

C. (+/-)-4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-(1-hydroxyethyl) [1,1'-biphenyl]-2-sulfonamide


258B (210 mg) was deprotected according to General Method 8 (EtOH). The crude product was purified by preparative thin-layer silica gel chromatography using 9:1 chloroform/methanol as eluant to provide the title compound (6 mg, 3% from **258A**) as a white solid: MS m/e 579 (ESI+ mode); HPLC retention time 3.07 and 3.40 min (Method A; product appears as two peaks); HPLC purity 92%.

15

Example 259
4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-propyl [1,1'-biphenyl]-2-sulfonamide

20

30



- 5 A. N-(4,5-Dimethyl-3-isoxazolyl)-4'-formyl-N-methoxymethyl-2'-propyl
[1,1'-biphenyl]-2-sulfonamide

27C (570 mg, 2.5 mmol) was subjected to Suzuki coupling with [2-
 [[(4,5-dimethyl-3-isoxazolyl)(methoxymethyl)amino]-
 sulfonyl]phenyl]boronic acid according to General Method 1. **259A** (1.1 g)
 10 was obtained as an impure yellow oil following silica gel chromatography
 using 4:1 hexanes/ethyl acetate.

- B. N-(4,5-Dimethyl-3-isoxazolyl)-4'-hydroxymethyl-N-methoxymethyl-2'-propyl [1,1'-biphenyl]-2-sulfonamide

15 **259A** (1.1 g) was reduced with sodium borohydride in methanol according to General Method 11. **259B** (520 mg, 47% over two steps) was obtained as an oil following silica gel chromatography with 1:1 hexanes/ethyl acetate as eluant.

- 20 C. 4'-Bromomethyl-N-(4,5-dimethyl-3-isoxazolyl)-N-methoxymethyl-2'-
propyl [1,1'-biphenyl]-2-sulfonamide

259B (520 mg, 1.2 mmol) was converted to the corresponding bromide according to General Method 2, providing **259C** (425 mg, 71%) as a yellow solid following silica gel chromatography using 9:1 hexanes/ethyl acetate as eluant.

D. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-N-methoxymethyl-2'-propyl [1,1'-biphenyl]-2-sulfonamide


259C (800 mg, 1.6 mmol) was used to alkylate 2-butyl-1,3-diazaspiro[4.4]non-1-en-4-one according to General Method 4. **259D** (720 mg) was obtained as a crude oil, which was used without further purification.

E. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-propyl [1,1'-biphenyl]-2-sulfonamide

259D (720 mg) was deprotected according to General Method 7. The crude product was purified by silica gel column chromatography using 1:1 hexanes/ethyl acetate as eluant to provide the title compound (390 mg, 42% over two steps) as a white solid: mp 64-66 °C; MS m/e 577 (ESI+ mode); HPLC retention time 30.78 min (Method B); HPLC purity >98%.

Example 260

N-(4,5-Dimethyl-3-isoxazolyl)-4'-[[[(3-methoxy-2,6-dimethyl-4-pyridinyl)oxy]methyl]-2'-propyl [1,1'-biphenyl]-2-sulfonamide



25

A. N-(4,5-Dimethyl-3-isoxazolyl)-4'-[[[(3-methoxy-2,6-dimethyl-4-pyridinyl)oxy]methyl]-N-methoxymethyl-2'-propyl [1,1'-biphenyl]-2-sulfonamide


259C (1.9 g, 3.7 mmol) was used to alkylate 3-methoxy-2,6-dimethyl-4-(1H)-pyridinone according to General Method 22. The crude product was chromatographed on silica gel using 1:1 hexanes/ethyl acetate as eluant to provide **260A** (1.3 g, 62%) as an oil.

B. N-(4,5-Dimethyl-3-isoxazolyl)-4'-[[3-methoxy-2,6-dimethyl-4-pyridinyl)oxy]methyl]-2'-propyl [1,1'-biphenyl]-2-sulfonamide

260A (1.3 g, 2.2 mmol) was deprotected according to General Method 7. The crude product was purified by silica gel column chromatography using 3% methanol in dichloromethane as eluant, providing the title compound (540 mg, 45%) as a white solid: mp 56-59 °C; MS m/e 536 (ESI+ mode); HPLC retention time 26.73 min (Method B); HPLC purity 97%.

Example 261

1-[[2'-[[3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl-2-(ethoxymethyl)]] [1,1'-biphenyl]-4-yl]methyl]-4-ethyl-2-propyl-1H-imidazole-5-carboxamide



A. N-(3,4-Dimethyl-5-isoxazolyl)-4'-[(5-ethoxycarbonyl-4-ethyl-2-propylimidazol-1-yl)methyl]-2'-ethoxymethyl-N-[(2-trimethylsilyl)ethoxymethyl] [1,1'-biphenyl]-2-sulfonamide

Ethyl 4-ethyl-2-propylimidazole-5-carboxylate (140 mg, 0.66 mmol) was alkylated with **256E** according to General Method 22. Silica gel


chromatography using 1:3 hexanes/ethyl acetate as eluant provided **261A** (120 mg, 25%) as an orange oil.

- 5 B. 4'-[(5-Carboxy-4-ethyl-2-propylimidazol-1-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-ethoxymethyl[1,1'-biphenyl]-2-sulfonamide
261A (120 mg, 0.16 mmol) was subjected to sulfonamide deprotection according to General Method 8 (EtOH). The crude residue was dissolved in methanol (2 ml) and was then treated with 45% aqueous potassium hydroxide (2 ml) and heated at 65 °C for 3 h. The mixture was cooled, adjusted to pH 4 by the addition of concentrated hydrochloric acid and aqueous sodium dihydrogen phosphate, and was then extracted with ethyl acetate. The combined organic extracts were dried over sodium sulfate and concentrated to provide **261B** (64 mg) as a crude orange oil.
- 10
- 15 C. 1-[[2'-[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl-2-(ethoxymethyl)][1,1'-biphenyl]-4-yl]methyl]-4-ethyl-2-propyl-1H-imidazole-5-carboxamide
261B (20 mg) was subjected to amide formation according to General Method 12 using aqueous ammonia as the amine component. The crude material was purified by preparative thin-layer silica gel chromatography using 1:1 hexanes/acetone as eluant to provide the title compound (13 mg) as a white solid: MS m/e 580 (ESI+ mode); HPLC retention time 2.97 min (Method C); HPLC purity 98%.
- 20

25

Example 262


1-[[2'-[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl-2-(ethoxymethyl)][1,1'-biphenyl]-4-yl]methyl]-4-ethyl-N-methyl-2-propyl-1H-imidazole-5-carboxamide



5 **261B** (44 mg) was subjected to amide formation according to
 General Method 12 using aqueous methylamine as the amine component.
 The crude material was purified by preparative thin-layer silica gel
 chromatography using 1:1 hexanes/acetone as eluant to provide the title
 compound (21 mg) as an off-white solid: MS m/e 594 (ESI+ mode); HPLC
 10 retention time 3.07 min (Method C); HPLC purity 97%.

Example 263

15 1-[[2'-[[[(4,5-Dimethyl-3-isoxazolyl)amino]sulfonyl-2-methyl][1,1'-biphenyl]-
4-yl]methyl]-4-ethyl-N-methyl-2-propyl-1H-imidazole-5-carboxamide



20

A. N-(4,5-Dimethyl-3-isoxazolyl)-4'-methoxycarbonyl-N-
methoxymethyl-2'-methyl[1,1'-biphenyl]-2-sulfonamide

Methyl 4-bromo-3-methylbenzoate (2.3 g, 10 mmol) was subjected to
 Suzuki coupling with [2-[[[(4,5-dimethyl-3-
 25 isoxazolyl)(methoxymethyl)amino]-sulfonyl]phenyl]boronic acid (3.2 g, 6.5

mmol) according to General Method 1. **263A** (3.1 g) was obtained as an impure yellow oil following silica gel chromatography using 3:1 hexanes/ethyl acetate as eluant.

5 B. N-(4,5-Dimethyl-3-isoxazolyl)-4'-hydroxymethyl-N-methoxymethyl-2'-methyl[1,1'-biphenyl]-2-sulfonamide

263A (3.1 g) was treated with DIBAL-H according to the procedure of Example 230, step E, to provide **263B** (2.7 g) as a crude yellow oil.

10 C. 4'-Bromomethyl-N-(4,5-dimethyl-3-isoxazolyl)- N-methoxymethyl-2'-methyl[1,1'-biphenyl]-2-sulfonamide

263B (2.7 g) was converted to the corresponding bromide according to General Method 2, providing **263C** (1.7 g, 55% over 3 steps) as a colorless oil following silica gel chromatography using 5:1 hexanes/ethyl acetate as eluant.

15 D. N-(4,5-Dimethyl-3-isoxazolyl)-4'-[(5-ethoxycarbonyl-4-ethyl-2-propylimidazol-1-yl)methyl]-N-methoxymethyl-2'-methyl [1,1'-biphenyl]-2-sulfonamide

20 **263C** (1.7 g) was used to alkylate ethyl 4-ethyl-2-propylimidazole-5-carboxylate according to General Method 22. Silica gel chromatography using 3:1 hexanes/ethyl acetate as eluant provided **263D** (880 mg, 41%) as an orange oil.

25 E. 4'-[(5-Carboxy-4-ethyl-2-propylimidazol-1-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-methyl[1,1'-biphenyl]-2-sulfonamide

263D (880 mg) was subjected to the procedure of Example 261, step B, to provide **263E** (870 mg) as a crude yellow solid.

F. 1-[[2'-[[[(4,5-Dimethyl-3-isoxazolyl)amino]sulfonyl-2-methyl][1,1'-biphenyl]-4-yl]methyl]-4-ethyl-N-methyl-2-propyl-1H-imidazole-5-carboxamide


263E (830 mg) was subjected to amide formation according to

5 General Method 12 using aqueous methylamine as the amine component. The crude material was purified by silica gel chromatography using 1:1 hexanes/acetone as eluant to provide the title compound (400 mg) as a white solid following lyophilization: MS m/e 550 (ESI+ mode); HPLC retention time 2.98 min (Method C); HPLC purity >98%.

10

Example 264

N,4-Diethyl-1-[[2'-[[[(4,5-dimethyl-3-isoxazolyl)amino]sulfonyl-2-methyl][1,1'-biphenyl]-4-yl]methyl]-2-propyl-1H-imidazole-5-carboxamide



15

263E (20 mg) was subjected to amide formation according to


General Method 12 using aqueous ethylamine as the amine component.

20 The crude material was purified by preparative thin-layer silica gel chromatography using 9:1 methanol/chloroform as eluant to provide the title compound (10 mg) as a white solid following lyophilization: MS m/e 564 (ESI+ mode); HPLC retention time 3.10 min (Method C); HPLC purity 97%.

25

Example 265

1-[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl-2-ethyl][1,1'-biphenyl]-4-yl]methyl]-4-ethyl-2-propyl-1H-imidazole-5-carboxamide



5

A. N-(3,4-Dimethyl-5-isoxazolyl)-4'-[(5-ethoxycarbonyl-4-ethyl-2-propyl-imidazol-1-yl)methyl]-2'-ethyl-N-[(2-trimethylsilyl)ethoxymethyl] [1,1'-biphenyl]-2-sulfonamide

Ethyl 4-ethyl-2-propylimidazole-5-carboxylate (600 mg, 2.9 mmol)
 10 was alkylated with **257D** (1.76 g) according to General Method 22. Silica gel chromatography using 1:1 hexanes/ethyl acetate as eluant provided impure **265A** (785 mg) as an orange oil.


B. 4'-[(5-Carboxy-4-ethyl-2-propylimidazol-1-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-ethyl [1,1'-biphenyl]-2-sulfonamide
 15 **265A** (785 mg) was subjected to the procedure of Example 261, step B. Purification of the crude material by preparative reverse-phase HPLC provided **265B** (240 mg) as yellow oil.

20 C. 1-[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl-2-ethyl][1,1'-biphenyl]-4-yl]methyl]-4-ethyl-2-propyl-1H-imidazole-5-carboxamide

265B (240 mg) was subjected to amide formation according to General Method 12 using aqueous ammonia as the amine component. The
 25 crude material was purified by silica gel chromatography using 1:2 hexanes/acetone as eluant to provide the title compound (110 mg) as a white solid following lyophilization: MS m/e 550 (ESI+ mode); HPLC retention time 3.05 min (Method C); HPLC purity >98%.

Example 266

1-[[2'-[[[(4,5-Dimethyl-3-isoxazolyl)amino]sulfonyl-2-methyl][1,1'-biphenyl]-
4-yl)methyl]-4-ethyl-N-(1-methylethyl)-2-propyl-1H-imidazole-5-
carboxamide




10

263E (150 mg) was subjected to amide formation according to General Method 12 using isopropylamine as the amine component. The crude material was chromatographed on silica gel using 1:1
 15 hexanes/acetone as eluant to provide the title compound (27 mg) as a white solid: MS m/e 578 (ESI+ mode); HPLC retention time 3.21 min (Method C); HPLC purity 96%.

Example 267

1-[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl-2-(2-
fluoroethoxymethyl)][1,1'-biphenyl]-4-yl)methyl]-4-ethyl-2-propyl-1H-
imidazole-5-carboxamide



25

A. N-(3,4-Dimethyl-5-isoxazolyl)-4'-[(5-ethoxycarbonyl-4-ethyl-2-propyl-imidazol-1-yl)methyl]-2'-(2-fluoroethoxymethyl)-N-[(2-trimethylsilyl)ethoxymethyl] [1,1'-biphenyl]-2-sulfonamide

5 **255C** (660 mg, 1.1 mmol) was used to alkylate ethyl 4-ethyl-2-propylimidazole-5-carboxylate according to General Method 22. Silica gel chromatography using 7:3 hexanes/ethyl acetate as eluant provided **267A** (460 mg) as a yellow oil.

10 B. 4'-[(5-Carboxy-4-ethyl-2-propylimidazol-1-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-(2-fluoroethoxymethyl) [1,1'-biphenyl]-2-sulfonamide

267A (460 mg) was subjected to the procedure of Example 261, step

15 B. **267B** (360 mg) was obtained as a crude yellow oil, which was used without further purification.


C. 1-[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl-2-(2-fluoroethoxymethyl)]] [1,1'-biphenyl]-4-yl)methyl]-4-ethyl-2-propyl-1H-imidazole-5-carboxamide

20 **267B** (360 mg) was subjected to amide formation according to General Method 12 using aqueous ammonia as the amine component. The crude material was purified by silica gel chromatography using 1:2 hexanes/acetone as eluant to provide the title compound (180 mg) as a white solid following lyophilization: MS m/e 598 (ESI+ mode); HPLC
25 retention time 2.83 min (Method C); HPLC purity >98%.

Example 268

N-(4,5-Dimethyl-3-isoxazolyl)-2'-ethoxymethyl-4'-[[[(6-ethyl-3-methoxy-2-methyl-4-pyridinyl)oxy]methyl] [1,1'-biphenyl]-2-sulfonamide

30



- 5 A. N-(4,5-Dimethyl-3-isoxazolyl)-2'-ethoxymethyl-4'-[[[(6-ethyl-3-methoxy-2-methyl-4-pyridinyl)oxy]methyl]-N-[(2-methoxyethoxy)methyl]] [1,1'-biphenyl]-2-sulfonamide


252E (200 mg, 0.38 mmol) was used to alkylate 6-ethyl-3-methoxy-2-methyl-4-(1H)-pyridinone (prepared according to Katano, K.; et. al. *Meiji Seika Kenkyu Nenpo*, **1996**, 35, 62-65) following General Method 22. **268A** (155 mg, 70%) was produced as an oil following silica gel chromatography using 50:50:0.2 hexanes/ethyl acetate/triethylamine as eluant.

- 15 B. N-(4,5-Dimethyl-3-isoxazolyl)-2'-ethoxymethyl-4'-[[[(6-ethyl-3-methoxy-2-methyl-4-pyridinyl)oxy]methyl]] [1,1'-biphenyl]-2-sulfonamide

268A was deprotected according to General Method 7. The crude product was purified by silica gel column chromatography using 1:30 methanol/dichloromethane as eluant, providing the title compound (108 mg, 74%) as a light yellow solid: mp 64-72 °C; MS m/e 566 (ESI+ mode); HPLC retention time 13.87 min (Method E); HPLC purity 95%.

Example 269

- 25 N2-[[2'-[[[(4,5-Dimethyl-3-isoxazolyl)amino]sulfonyl]] [1,1'-biphenyl]-4-yl]methyl]-N-methyl-N2-(1-oxobutyl)-L-valinamide



5 A. N-(4,5-Dimethyl-3-isoxazolyl)-N-[[[(2-trimethylsilyl)ethoxy]methyl]-2-bromobenzenesulfonamide

SEM-Cl (6.4 ml, 36 mmol) was added at 0 °C to a mixture of N-(4,5-dimethyl-3-isoxazolyl)-2-bromobenzenesulfonamide (11 g, 34 mmol), potassium carbonate (9.4 g, 68 mmol), and DMF (100 ml). The mixture
10 was allowed to warm to rt and was stirred for 18 h. The solvent was evaporated, water (200 mL) was added, and the mixture was extracted with ethyl acetate. The crude extract was dried over sodium sulfate and concentrated, and the residue was chromatographed on silica gel using 4:1 hexanes/ethyl acetate as eluant to provide **269A** (11 g, 70%) as an oil.

15

B. N-(4,5-Dimethyl-3-isoxazolyl)-4'-formyl-N-[[[(2-trimethylsilyl)ethoxy]methyl] [1,1'-biphenyl]-2-sulfonamide

269A (2.3 g, 5.0 mmol) was subjected to Suzuki coupling with 4-formylphenylboronic acid according to General Method 1. **269B** (2.0 g, 83
20 %) was obtained as yellow crystalline solid following silica gel chromatography using 4:1 hexanes /ethyl acetate as eluant.

C. N2-[[[2'-[[N-(4,5-Dimethyl-3-isoxazolyl)-N-[[[(2-trimethylsilyl)ethoxy]methyl]amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-methyl-L-valinamide

25

269B (490 mg, 1.0 mmol) and L-valine N-methylamide were subjected to reductive amination according to General Method 5. Crude **269C** (780 mg) was produced as an oil.

D. N2-[[2'-[[N-(4,5-Dimethyl-3-isoxazolyl)-N-[(2-trimethylsilyl)ethoxy]methyl]amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-methyl-N2-(1-oxobutyl)-L-valinamide

5 **269C** (750 mg) was acylated with butanoyl chloride according to General Method 6 to provide crude **269D** (680 mg) as an oil.


E. N2-[[2'-[[[(4,5-Dimethyl-3-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-methyl-N2-(1-oxobutyl)-L-valinamide

10 **269D** (680 mg) was deprotected according to General Method 10. Silica gel chromatography of the crude residue using dichloromethane/methanol as eluant provided the title compound (360 mg) as a white amorphous solid: MS m/e 541 (ESI+ mode); HPLC retention time 3.61 min (Method C); HPLC purity 95%.

15

Example 270

N2-[[2'-[[[(4,5-Dimethyl-3-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N,N-dimethyl-N2-(1-oxopentyl)-L-valinamide



20

A. N2-[[2'-[[N-(4,5-Dimethyl-3-isoxazolyl)-N-[(2-trimethylsilyl)ethoxy]methyl]amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N,N-dimethyl-L-valinamide

25

269B (490 mg, 1.0 mmol) and L-valine N,N-dimethylamide were subjected to reductive amination according to General Method 5. Crude **270A** (780 mg) was produced as an oil.


- 5 B. N2-[[2'-[[N-(4,5-Dimethyl-3-isoxazolyl)-N-[[[(2-trimethylsilyl)ethoxy)methyl]amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N,N-dimethyl-N2-(1-oxopentyl)-L-valinamide
270A (750 mg) was acylated with pentanoyl chloride according to General Method 6 to provide crude **270B** (710 mg) as an oil.

10

- C. N2-[[2'-[[[(4,5-Dimethyl-3-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N,N-dimethyl-N2-(1-oxopentyl)-L-valinamide
270B (710 mg) was deprotected according to General Method 10. Silica gel chromatography of the crude residue using dichloromethane/
 15 methanol as eluant provided the title compound (400 mg) as a white amorphous solid: MS m/e 569 (ESI+ mode); HPLC retention time 3.82 min (Method C); HPLC purity 95%.

Example 271

- 20 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-ethyl [1,1'-biphenyl]-2-sulfonamide



25

- A. N-(4,5-Dimethyl-3-isoxazolyl)-2'-ethyl-4'-methoxycarbonyl-N-(methoxymethyl) [1,1'-biphenyl]-2-sulfonamide

257A (5.9 g, 24 mmol) was subjected to Suzuki coupling with 2-[[N-(4,5-dimethyl-3-isoxazolyl)-N-(methoxymethyl)amino]sulfonyl]phenylboronic acid according to General Method 1. Silica gel chromatography provided **271A** (6.0 g) as an oil, contaminated with byproducts deriving from the boronic acid. **257A** (4.5 g) was also recovered. A single recycling of the recovered **257A** yielded 6.6 g of the **271A** product mix (12.6 g total combined yield).

B. N-(4,5-Dimethyl-3-isoxazolyl)-2'-ethyl-4'-hydroxymethyl-N-(methoxymethyl) [1,1'-biphenyl]-2-sulfonamide

271A (11.1 g) was treated with DIBAL-H according to the procedure of Example 230, step E, to provide **271B** (6.5 g) as a crude oil.

C. 4'-Bromomethyl-N-(4,5-dimethyl-3-isoxazolyl)-2'-ethyl-N-(methoxymethyl) [1,1'-biphenyl]-2-sulfonamide

271B (6.5 g) was converted to the corresponding bromide according to General Method 2 to provide **271C** (4.2 g) as an orange oil following silica gel chromatography using 3:1 hexanes/ethyl acetate as eluant.

D. 4'-[[2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-ethyl- N-(methoxymethyl) [1,1'-biphenyl]-2-sulfonamide

271C (2.0 g) was used to alkylate 2-butyl-1,3-diazaspiro[4.4]non-1-en-4-one according to General Method 4. **271D** (1.6 g) was produced as a colorless oil following silica gel chromatography using 1:1 hexanes/ethyl acetate as eluant.

E. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-ethyl [1,1'-biphenyl]-2-sulfonamide


271D (1.6 g) was deprotected according to General Method 8 (EtOH). The crude product was purified by silica gel chromatography

using 1:1 hexanes/ethyl acetate as eluant to provide the title compound (815 mg) as a white amorphous solid: MS m/e 563 (ESI+ mode); HPLC retention time 2.18 min (Method H); HPLC purity >98%.

5

Example 272

4'-[(5-Acetyl-4-ethyl-2-propylimidazol-1-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-methyl[1,1'-biphenyl]-2-sulfonamide



10

A. 5-Acetyl-4-ethyl-2-propylimidazole

4-Ethyl-5-formyl-2-propylimidazole (4 g, 24 mmol) was subjected to the procedure of Example 38, steps A and B. Silica gel chromatography using 1:3 hexanes/ethyl acetate as eluant provided **272A** (2.4 g, 55% over two steps) as a brown oil.

B. 4'-[(5-Acetyl-4-ethyl-2-propylimidazol-1-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-N-methoxymethyl-2'-methyl[1,1'-biphenyl]-2-sulfonamide

272A (800 mg, 4.4 mmol) was alkylated with **263C** according to General Method 22. Silica gel chromatography using 1:3 hexanes/ethyl acetate as eluant provided **272B** (990 mg, 38%) as an orange oil.


25

C. 4'-[(5-Acetyl-4-ethyl-2-propylimidazol-1-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-methyl[1,1'-biphenyl]-2-sulfonamide

272B (940 mg, 1.6mmol) was subjected to sulfonamide deprotection according to General Method 8 (water). The crude material was purified by reverse-phase preparative HPLC followed by preparative thin-layer silica gel chromatography (chloroform/methanol eluant) to provide the title compound (9 mg) as a white solid following lyophilization: MS m/e 535 (ESI+ mode); HPLC retention time 1.98 min (Method H); HPLC purity 98%.

Example 273

10 N-(4,5-Dimethyl-3-isoxazolyl)-2'-ethyl-4'-[[[(3-methoxy-2,6-dimethyl-4-pyridinyl)oxy]methyl] [1,1'-biphenyl]-2-sulfonamide



15

A. N-(4,5-Dimethyl-3-isoxazolyl)-2'-ethyl-4'-[[[(3-methoxy-2,6-dimethyl-4-pyridinyl)oxy]methyl]-N-(methoxymethyl) [1,1'-biphenyl]-2-sulfonamide

20 **271C** (2.0 g, 4.0 mmol) was used to alkylate 3-methoxy-2,6-dimethyl-4-(1H)-pyridinone according to General Method 22. The crude product was purified by silica gel chromatography using 1:1 hexanes/ethyl acetate as eluant to provide **273A** (1.2 g, 52%) as an orange oil.

25 **B.** N-(4,5-Dimethyl-3-isoxazolyl)-2'-ethyl-4'-[[[(3-methoxy-2,6-dimethyl-4-pyridinyl)oxy]methyl] [1,1'-biphenyl]-2-sulfonamide


273A (1.2 g, 2.2 mmol) was deprotected according to General Method 8 (EtOH). The crude product was crystallized from

methanol/dichloromethane to provide the title compound (172 mg) as a white solid: MS m/e 522 (ESI+ mode); HPLC retention time 1.53 min (Method H); HPLC purity 95%.

5

Example 274

N-(4,5-Dimethyl-3-isoxazolyl)-4'-[[[(6-ethyl-3-methoxy-2-methyl-4-pyridinyl)oxy]methyl]-2'-propyl [1,1'-biphenyl]-2-sulfonamide



10

A. N-(4,5-Dimethyl-3-isoxazolyl)-4'-[[[(6-ethyl-3-methoxy-2-methyl-4-pyridinyl)oxy]methyl]-N-methoxymethyl-2'-propyl [1,1'-biphenyl]-2-sulfonamide

15

259C (1.97 g) was used to alkylate 6-ethyl-3-methoxy-2-methyl-4-(1H)-pyridinone (500 mg, prepared according to Katano, K.; et. al. *Meiji Seika Kenkyu Nenpo*, **1996**, 35, 62-65) following General Method 22. **274A** (1.23 g, 76%) was produced as an oil following silica gel chromatography using hexanes/ethyl acetate/triethylamine as eluant.

20


B. N-(4,5-Dimethyl-3-isoxazolyl)-4'-[[[(6-ethyl-3-methoxy-2-methyl-4-pyridinyl)oxy]methyl]-2'-propyl [1,1'-biphenyl]-2-sulfonamide

25

274A (1.23 g) was deprotected according to General Method 7. The crude product was purified by silica gel column chromatography using methanol/dichloromethane as eluant, providing the title compound (820 mg, 72%) as a white solid: mp 63-65 °C; MS m/e 550 (ESI+ mode); HPLC retention time 3.04 min (Method A); HPLC purity >98%.

Example 275

5 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-(ethoxymethyl) [1,1'-biphenyl]-2-sulfonamide [crystalline]



(alternative synthesis of Example 227)

10

A. Ethyl 4-bromo-3-(bromomethyl)benzoate

Ethyl 4-bromo-3-methylbenzoate (110 g, 450 mmol) was treated with NBS according to the procedure of **P2A**. Silica gel chromatography with hexanes/ethyl acetate as eluant provided **275A** (91 g, 62%) as a white solid.

15

B. Ethyl 4-bromo-3-(ethoxymethyl)benzoate

A solution of **275A** (89 g, 280 mmol) in a mixture of ethanol (300 ml) and DMF (50 ml) was treated at 0 °C with sodium ethoxide (135 ml of a 21% solution in ethanol). The mixture was allowed to warm to rt and was stirred for 16 h. The ethanol was evaporated under reduced pressure. Ethyl acetate was added to the residue and the mixture was washed with water and brine. The organic layer was dried over sodium sulfate and concentrated, and the residue was chromatographed on silica gel using hexanes/ethyl acetate as eluant to provide **275B** (67 g, 84%) as a slightly yellow oil.

25

C. N-(4,5-Dimethyl-3-isoxazolyl)-4'-(ethoxycarbonyl)-2'-(ethoxymethyl)-N-(methoxymethyl) [1,1'-biphenyl]-2-sulfonamide

275B (32 g, 100 mmol) was subjected to Suzuki coupling with 2-[[N-(4,5-dimethyl-3-isoxazolyl)-N-(methoxymethyl)amino]sulfonyl]-

phenylboronic acid according to General Method 1. Silica gel chromatography using hexanes/ethyl acetate as eluant provided **275C** (52 g) as a yellow oil, contaminated with byproducts deriving from the boronic acid.

- 5 D. N-(4,5-Dimethyl-3-isoxazolyl)-2'-(ethoxymethyl)-4'-(hydroxymethyl)-N-(methoxymethyl) [1,1'-biphenyl]-2-sulfonamide

275C (entire sample) was treated with DIBAL-H according to the procedure of Example 230, step E, with the following difference: Workup was by addition of saturated aqueous ammonium chloride to the cooled reaction mixture, followed by
10 extraction with ethyl acetate. The combined organic layers were dried over sodium sulfate and concentrated to provide **275D** as a crude yellow oil.

- E. 4'-(Bromomethyl)-N-(4,5-dimethyl-3-isoxazolyl)-2'-(ethoxymethyl)-N-(methoxymethyl) [1,1'-biphenyl]-2-sulfonamide

15 **275D** (entire sample) was converted to the corresponding bromide according to General Method 2. Silica gel chromatography using hexanes/ethyl acetate as eluant provided **275E** (38 g, purity estimated to be 83% by ¹H NMR) as a light yellow oil.

- F. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-(ethoxymethyl)-N-(methoxymethyl) [1,1'-biphenyl]-2-sulfonamide

275E (entire sample) was used to alkylate 2-butyl-1,3-diazaspiro[4.4]non-1-en-4-one according to General Method 4. The crude residue was chromatographed on silica gel using hexanes/ethyl acetate/triethylamine as eluant to provide **275F** (32 g,
25 53% from **275B**) as a slightly yellow oil.

- G. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-(ethoxymethyl) [1,1'-biphenyl]-2-sulfonamide

275F (32 g, 53 mmol) was deprotected according to General Method 7. The
30 crude product was purified by silica gel chromatography using hexanes/ethyl acetate/acetic acid as eluant to provide the title compound (26 g, 88%) as an

amorphous foam: MS m/e 593 (ESI+ mode); HPLC retention time 18.75 min (Method E); HPLC purity >96%.


H. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-(ethoxymethyl) [1,1'-biphenyl]-2-sulfonamide

(crystallization)

The amorphous **275F** (1 g) was dissolved in 5 mL of isopropanol and 5 mL of water was added to the mixture dropwise and the mixture was warmed up to 40°C to provide a clear solution. The solution was let stand at room temperature and the white crystals thus obtained was filtered and washed with a small amount of 2:1 mixture of isopropanol / water and dried to give 0.87 g of a white crystalline solid. mp. 148°C.

EXAMPLE 276

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-3'-chloro-N-(4,5-dimethyl-3-isoxazolyl)-5-methoxy[1,1'-biphenyl]-2-sulfonamide



A. (2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl-2-chloro-4-bromobenzene.

α -Bromomethyl-4-bromo-2-chlorobenzene (1.6gm) was used to alkylate 2-butyl-1,3-diazaspiro[4.4]non-1-en-4-one (1.2gm) according to General Method 4 to yield **276A** (2.2gm, 98%) as a solid. MS m/e 399.13.

B. 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-3'-chloro-N-[(2-methoxyethoxy)methyl]-N-(4,5-dimethyl-3-isoxazolyl)[1,1'-biphenyl]-2-sulfonamide.

276A (218 mg) was subjected to Suzuki coupling with (2-[[4,5-dimethyl-3-isoxazolyl]{2-methoxyethoxy)methyl}amino]-sulfonyl]-5-methoxyphenyl]boronic acid (270 mg) according to General Method 1 to provide **276B** (280 mg, 67%) as an oil.

C. Title Compound

276B (280 mg) was deprotected according to General Method 8 to provide the title compound (178 mg, 98%) as a white amorphous solid: MS m/e 583.

Examples 277-297

The compounds of Examples 277-297 were prepared in a manner similar to that described in Example 276. The reported HPLC retention times were obtained under the following conditions.

Column: YMC S5 ODS 4.6X50 mm (4min); Wavelength: 220nm;






Solvent: Gradient elution 10%MeOH--90% MeOH in water





(0.2% H₃PO₄).

E X A M P L E	S T R U C T U R E	N A M E	M/z (MH) ⁺	HPLC ret time, min

277		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-3'-fluoro-4-methoxy[1,1'-biphenyl]-2-sulfonamide	583.3	3.2min
278		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-5'-chloro-N-(4,5-dimethyl-3-isoxazolyl)-2'-fluoro-4-methoxy[1,1'-biphenyl]-2-sulfonamide	617.3	3.3min
279		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-3'-chloro-N-(4,5-dimethyl-3-isoxazolyl)-4-methoxy[1,1'-biphenyl]-2-sulfonamide	599.2	3.4min
280		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-4-methoxy[1,1'-biphenyl]-2-sulfonamide	565.3	3.0min
281		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-(ethoxymethyl)-4-methoxy[1,1'-biphenyl]-2-sulfonamide	623.51	3.11min
282		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-3'-fluoro-5-methoxy[1,1'-biphenyl]-2-sulfonamide	583	3.1min

283		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-5'-chloro-N-(4,5-dimethyl-3-isoxazolyl)-2'-fluoro-5-methoxy[1,1'-biphenyl]-2-sulfonamide	617.5	3.3min
284		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-5-methoxy[1,1'-biphenyl]-2-sulfonamide	565.3	3.0min
285		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-(ethoxymethyl)-5-methoxy[1,1'-biphenyl]-2-sulfonamide	623.6	3.03min
286		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-3'-fluoro-4,5-dimethoxy[1,1'-biphenyl]-2-sulfonamide	613.4	3.04min
287		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-5'-chloro-N-(4,5-dimethyl-3-isoxazolyl)-2'-fluoro-4,5-dimethoxy[1,1'-biphenyl]-2-sulfonamide	647.2	3.2min

288		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-3'-chloro-N-(4,5-dimethyl-3-isoxazolyl)-4,5-dimethoxy[1,1'-biphenyl]-2-sulfonamide	629.2	3.2min
289		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-4,5-dimethoxy[1,1'-biphenyl]-2-sulfonamide	595.3	2.8min
290		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-(ethoxymethyl)-4,5-dimethoxy[1,1'-biphenyl]-2-sulfonamide	653.48	2.89min
291		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-5'-chloro-N-(4,5-dimethyl-3-isoxazolyl)-2'-fluoro[1,1'-biphenyl]-2-sulfonamide	587.2	3.3min
292		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-3'-chloro-N-(4,5-dimethyl-3-isoxazolyl)[1,1'-biphenyl]-2-sulfonamide	569.2	3.2min

293		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-4-methoxy-3'-methyl[1,1'-biphenyl]-2-sulfonamide	579.4	3.2min
294		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-3'-fluoro-4-methoxy[1,1'-biphenyl]-2-sulfonamide	553.1	3.0min
295		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-3'-methyl[1,1'-biphenyl]-2-sulfonamide	549.2	3.01min
296		4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-5-fluoro[1,1'-biphenyl]-2-sulfonamide	553.23	2.90min





Examples 297-309






5 The compounds of Examples 297-309 were prepared in a manner similar to that described in Example 269. The reported HPLC retention times were obtained under the following conditions.





Column: YMC S5 ODS 4.6X50 mm (4min); Wavelength: 220nm;

Solvent: Gradient elution 10%MeOH--90% MeOH in water

10 (0.2% H3PO4).

E X A M P L E	S T R U C T U R E	N A M E	M/z (MH) ⁺	HPLC ret time, min
297		N ² -(Cyclopropylcarbonyl)-N ² -[[2'-[[[(4,5-dimethyl-3-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-methyl-L-valinamide	539.17	3.08min
298		N ² -[[2'-[[[(4,5-Dimethyl-3-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N,3-dimethyl-N ² -(1-oxobutyl)-L-valinamide	555.24	3.40min
299		N ² -[[2'-[[[(4,5-Dimethyl-3-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-methyl-N ² -(2-methyl-1-oxopropyl)-L-valinamide	541.29	3.17min
300		N ² -(Cyclopentylcarbonyl)-N ² -[[2'-[[[(4,5-dimethyl-3-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-methyl-L-valinamide	567.28	3.40min

301		N ² -[[2'-[[[(4,5-Dimethyl-3-isoxazolyl)amino]sulfonyl]-3-fluoro[1,1'-biphenyl]-4-yl]methyl]-N-methyl-N ² -(1-oxobutyl)-L-valinamide	559.2	3.40min
302		N ² -[[2'-[[[(4,5-Dimethyl-3-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-(1-methylethyl)-N ² -(1-oxobutyl)-L-valinamide	569.3	3.40min
303		N ² -[[2'-[[[(4,5-Dimethyl-3-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-(2-methoxyethyl)-N ² -(1-oxobutyl)-L-valinamide	585.3	3.27min
304		N-(Cyclopropylmethyl)-N ² -[[2'-[[[(4,5-dimethyl-3-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N ² -(1-oxobutyl)-L-valinamide	581.3	3.44min
305		N ² -[[2'-[[[(4,5-Dimethyl-3-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N ² -(1-oxobutyl)-N-(3-pyridinyl)-L-valinamide	604.3	2.87min

306		N²-[[2'-[[[(4,5-Dimethyl-3-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-methyl-N²-(1-oxopentyl)-L-valinamide	555.17	3.36min
307		N-Methyl-N²-[[2'-[[[(5-methyl-3-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N²-(1-oxopentyl)-L-valinamide	541.2	3.40min
308		N²-[[2'-[[[(4,5-Dimethyl-3-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-ethyl-N²-(1-oxobutyl)-L-valinamide	555.3	3.17min
309		N²-[[2'-[[[(4,5-Dimethyl-3-isoxazolyl)amino]sulfonyl]-5'-fluoro[1,1'-biphenyl]-4-yl]methyl]-N-methyl-N²-(1-oxobutyl)-L-valinamide	558.76	3.28min



Examples 310-311

5 The compounds of Examples 310-311 were prepared in a manner similar to that described in Example 252. The reported HPLC retention times were obtained under the following conditions.

Column: YMC S5 ODS 4.6X50 mm (4min); Wavelength: 220nm;

Solvent: Gradient elution 10%MeOH--90% MeOH in water

(0.2% H₃PO₄).

E X A M P L E	S T R U C T U R E	N A M E	M/z (MH) ⁺	HPLC ret time, min
310		N-(4,5-Dimethyl-3-isoxazolyl)-4'-[[[(2,6-dimethyl-3-methoxy-4-pyridinyl)oxy)methyl][1,1'-biphenyl]-2-sulfonamide	494	2.28min
311		N-(4,5-Dimethyl-3-isoxazolyl)-4'-[[[(2,6-dimethyl-3-methoxy-4-pyridinyl)oxy)methyl]-3'-fluoro[1,1'-biphenyl]-2-sulfonamide	512	2.41min




5


Examples 312-314

The compounds of Examples 312-314 were prepared in a manner similar to that described in Example 228. The reported HPLC retention times were obtained under the following conditions.


Column: YMC S5 ODS 4.6X50 mm (4min); Wavelength: 220nm;

10 Solvent: Gradient elution 10%MeOH--90% MeOH in water
(0.2% H₃PO₄).


E X A M P L E	S T R U C T U R E	N A M E	M/z (MH) ⁺	HPLC ret time, min
312		N-(4,5-Dimethyl-3-isoxazolyl)-4'-[(1,4,5,6,7,8-hexahydro-8-oxo-2-propyl-1-cycloheptimidazolyl)methyl][1,1'-biphenyl]-2-sulfonamide	533.3	2.35min
313		N-(4,5-Dimethyl-3-isoxazolyl)-3'-fluoro-4'-[(1,4,5,6,7,8-hexahydro-8-oxo-2-propyl-1-cycloheptimidazolyl)methyl][1,1'-biphenyl]-2-sulfonamide	551.3	2.35min
314		N-(4,5-Dimethyl-3-isoxazolyl)-5-fluoro-4'-[(1,4,5,6,7,8-hexahydro-8-oxo-2-propyl-1-cycloheptimidazolyl)methyl][1,1'-biphenyl]-2-sulfonamide	551.17	2.32min

Example 315

5 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-(hydroxymethyl)[1,1'-biphenyl]-2-sulfonamide

A.

5E (4.49 g, 11.48 mmol) was subjected to Suzuki coupling with [2-[(4,5-dimethyl-3-isoxazolyl)-N-methoxymethyl]amino]-sulfonylphenylboronic acid according to General Method 1. **316A** (5.0 g, 72%) was obtained as an orange oil following silica gel chromatography using hexanes/ethyl acetate as eluant.


10 **B.**

315A (0.7g) was dissolved in 10 mL of methylene chloride and 2 mL of triethylsilane and 2 mL of trifluoroacetic acid were added to the mixture. The solution was stirred at room temperature for 1h and concentrated.

15 **315B** (0.6 g, 85%) was obtained as an orange oil following silica gel chromatography using hexanes/ethyl acetate as eluant.

C.

315B (0.6 g) was deprotected according to General Method 8 (EtOH). The crude product was purified by silica gel chromatography using 1:1 hexanes/ethyl acetate as eluant to provide the title compound (0.44 g, 79%) as a white amorphous solid. mp. 90-98°C. **M+H: 565.2**

Example 316

- 5 N-[[2'-[[4,5-Dimethyl-3-isoxazolyl]amino]sulfonyl]-2-(ethoxymethyl)[1,1'-biphenyl]-4-yl]methyl]-1-[(1-oxopentyl)amino]cyclopentanamide
(A metabolite of Example 227)

227 (30 mg) was dissolved in 1mL DMF and 0.6 mL of 20% aqueous NaOH
10 was added and the mixture was stirred at 60°C for 24h. The mixture was purified by HPLC to provide the target compound as a white solid. HPLC RT: 3.21 min. M+H: 611.3

15 **Example 317**

(+)-4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-(ethoxymethyl) [1,1'-biphenyl]-2-sulfonamide
and

20 (-)-4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-(ethoxymethyl) [1,1'-biphenyl]-2-sulfonamide


10µl of a solution of **275F** (2mg/mL) in isopropanol was injected onto an HPLC column (Chiral AD, 250x4.6 (10µ)) eluting with a mobil phase of
25 80% hexane, 20% IPA, 0.1%TFA, 0.1%TEA at 0.75mL/min. The HPLC column temperature was maintained at 15°C. Detection was carried out

at 210nm using a UV detector. The enantiomers of **275** were resolved into two peaks, with one enantiomer eluting at 11.6 min and the other at 15.1 min. yielding the two title compounds as pure enantiomers.


What is claimed is:


1. A compound of the following formula I, or an enantiomer, diastereomer, salt or metabolite thereof:

5



wherein:







R_2 is hydrogen, halogen, -CHO, alkyl, haloalkyl, (cycloalkyl)alkyl, alkenyl,
 5 alkynyl, alkoxyalkyl, haloalkoxyalkyl, alkoxy, aryloxy alkoxyalkoxy,
 cyano, hydroxy, hydroxyalkyl, nitro, $-\text{CH}(\text{OR}_{13})(\text{OR}_{14})$, $-(\text{CH}_2)_w\text{Y}$;
 with the proviso that when R_1 is **B**, R_2 is not hydrogen, halogen,
 alkyl, haloalkyl, alkoxy, hydroxyalkyl, nitro, $-(\text{CH}_2)_w\text{NR}_{19}\text{R}_{20}$ or
 $-\text{NHSO}_2\text{R}_{22}$;


10 R_3 is heteroaryl;

R_4 and R_5 are each independently alkyl, hydroxyalkyl, cycloalkyl, hydroxy
 substituted cycloalkyl, alkoxyalkyl, or hydroxy substituted
 alkoxyalkyl, or R_4 and R_5 together form a cyclobutyl, cyclopentyl,
 cyclohexyl, tetrahydrofuranyl or tetrahydropyranyl ring which may
 15 be optionally substituted with one or more hydroxy groups;

R_6 is alkyl, hydroxyalkyl, haloalkyl, hydroxy substituted haloalkyl,
 cycloalkyl, hydroxy substituted cycloalkyl, (cycloalkyl)alkyl,
 hydroxy substituted (cycloalkyl)(alkyl), aralkyl, alkoxy, hydroxy
 substituted alkoxy, alkoxyalkyl, hydroxy substituted alkoxyalkyl, or
 20 $-\text{NR}_{16}\text{R}_{17}$;


- R_7 is $-(CH_2)_w-CO_2R_{15}$, $-(CH_2)_w-(C=O)NR_{16}R_{17}$, $-(CH_2)_w-NR_{15}(C=O)NR_{16}R_{17}$, $-(CH_2)_w-CH_2OH$, $-(CH_2)_w-(C=O)R_{15}$, tetrazolyl, oxadiazolyl or triazolyl wherein said tetrazolyl, oxadiazolyl or triazolyl may optionally be substituted with hydrogen, alkyl, hydroxy or halogen;
- 5 R_8 , R_9 , R_{9a} , R_{10} and R_{12} are each independently hydrogen, halogen, alkyl, hydroxyalkyl, cycloalkyl, (cycloalkyl)alkyl, aryl, heteroaryl, arylalkyl, alkylthioalkyl, alkoxy or alkoxyalkyl, or R_9 and R_{9a} together with the carbon atom to which they are bonded form a cycloalkyl ring;
- 10 R_{11} and R_{11a} are each independently hydrogen, alkoxy, or together form a carbonyl;
- R_{13} and R_{14} are alkyl or together form a five to six-membered ring;
- R_{15} , R_{16} and R_{17} are independently hydrogen, alkyl, hydroxyalkyl, cycloalkyl, (cycloalkyl)alkyl, alkoxyalkyl, aralkyl, heterocycloalkyl,
- 15 aryl, heteroaryl or $-(CH_2)_wQ$, or R_{16} and R_{17} may together form a four to six-membered heterocyclic ring;
- n is 1 or 2;
- w is 0, 1, or 2;
- Y is heteroaryl, $-COOH$, $-COOR_{18}$, $-CONR_{19}R_{20}$, $-NR_{19}R_{20}$, $-NR_{19}-OR_{20}$,
- 20 $-NR_{21}(C=O)R_{22}$, $-NR_{21}(C=O)NR_{19}R_{20}$, $-N(R_{19})-(alk)-NR_{21}(C=O)R_{22}$, $-NR_{21}(C=O)OR_{18}$, $-NR_{21}SO_2R_{22}$, $-SO_2R_{22}$, Q, R or S;





R_{18} , R_{19} , R_{20} , R_{21} and R_{22} are each independently hydrogen, alkyl, haloalkyl, alkoxyalkyl, cycloalkyl, alkenyl, alkynyl, aryl, aralkyl, heteroaryl, or R_{19} and R_{20} may together form a four to seven-membered heterocyclic ring;

R_{23} and R_{24} are each independently hydrogen, alkyl or cycloalkyl, or may together form a three to seven membered cycloalkyl ring;




x is 2, 3 or 4;

R_{25} , R_{26} and R_{27} are each independently hydrogen, alkyl or cycloalkyl, or R_{26} and R_{27} may together form a three to seven-membered cycloalkyl ring;


R_{101} , R_{102} , R_{103} , and R_{104} are each independently hydrogen, halogen, -CHO, alkyl, haloalkyl, (cycloalkyl)alkyl, alkenyl, alkynyl, alkoxyalkyl, haloalkoxyalkyl, alkoxy, alkoxyalkoxy, cyano, hydroxy, hydroxyalkyl, nitro, -CH(OR₁₃)(OR₁₄), or -(CH₂)_wY; wherein said rings; aryl alone or as part of another group; or heteroaryl alone or as part of another group may each optionally be substituted by one or more hydrogen, halogen, cyano, alkyl, hydroxyalkyl, alkoxy, nitro or trifluoromethyl groups;

provided that when R₁ is A said compound is other than



2. A compound of claim 1, wherein R_3 is isoxazol-5-yl or isoxazol-3-yl independently substituted with two substituents independently selected from alkyl or halogen.

5




3. A compound of claim 2, wherein R_1 is A.

4. A compound of claim 3, wherein R_2 is alkyl, haloalkyl, alkoxyalkyl or haloalkoxyalkyl and R_{101} , R_{102} , R_{103} , R_{104} are each
10 independently hydrogen, halogen, or alkyl.

5. A compound of claim 3, wherein R_2 is $-CH_2Y$.

6. A compound of claim 5, wherein Y is Q.



15


7. A compound of claim 2, wherein R_1 is E.

8. A compound of claim 7, wherein R_2 is alkyl, haloalkyl, alkoxyalkyl or haloalkoxyalkyl and R_{101} , R_{102} , R_{103} , R_{104} are each
independently hydrogen, halogen, or alkyl.

20

9. A compound of claim 7, wherein R_2 is $-CH_2Y$.

10. A compound of claim 9, wherein Y is Q.



11. A compound of claim 2, wherein R_1 is

12. A compound of claim 11, wherein R_2 is alkyl, haloalkyl, alkoxyalkyl or haloalkoxyalkyl and R_{101} , R_{102} , R_{103} , R_{104} are each

5 independently hydrogen, halogen, or alkyl.

13. A compound of claim 11, wherein R_2 is $-\text{CH}_2\text{Y}$.

14. A compound of claim 13, wherein Y is $\underline{\text{Q}}$.

10

15. A compound of claim 1, wherein R_2 is alkoxyalkyl alkyl, haloalkyl or haloalkoxyalkyl

16. A compound of claim 15, wherein R_3 is isoxazol-5-yl or isoxazol-3-yl independently substituted with two substituents selected from alkyl or halogen.

15

17. A compound of claim 1, wherein R_2 is $-\text{CH}_2\text{Y}$.

20

18. A compound of claim 17, wherein R_3 is isoxazol-5-yl or isoxazol-3-yl independently substituted with two substituents selected from alkyl or halogen.

19. A compound of claim 17, wherein Y is $\underline{\text{Q}}$.

25

20. A compound of claim 19, wherein R_3 is isoxazol-5-yl or isoxazol-3-yl independently substituted with two substituents selected from alkyl or halogen.

21. A compound of claim 1, wherein said compound is selected from
 N,4-Diethyl-1-[[2'-[[[(4,5-dimethyl-3-isoxazolyl)amino]sulfonyl-2-
 methyl][1,1'-biphenyl]-4-yl]methyl]-2-propyl-1H-imidazole-5-carboxamide;
 5 1-[[2'-[[[(4,5-dimethyl-3-isoxazolyl)amino]sulfonyl-2-methyl][1,1'-biphenyl]-4-
 yl]methyl]-4-ethyl-N-(1-methylethyl)-2-propyl-1H-imidazole-5-carboxamide;
 N-(4,5-Dimethyl-3-isoxazolyl)-2'-ethoxymethyl-4'-[[[(6-ethyl-3-methoxy-2-
 methyl-4-pyridinyl)oxy]methyl] [1,1'-biphenyl]-2-sulfonamide;
 4'-[(5-Acetyl-4-ethyl-2-propylimidazol-1-yl)methyl]-N-(4,5-dimethyl-
 10 3-isoxazolyl)-2'-methyl[1,1'-biphenyl]-2-sulfonamide;
 N-(4,5-Dimethyl-3-isoxazolyl)-4'-[[[(6-ethyl-3-methoxy-2-methyl-4-
 pyridinyl)oxy]methyl]-2'-propyl [1,1'-biphenyl]-2-sulfonamide;
 and salts, enantiomers, diastereomers and metabolites thereof.
22. A compound of claim 1, wherein said compound is selected from
 N,4-Diethyl-1-[[2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl-2-
 methyl][1,1'-biphenyl]-4-yl]methyl]-2-propyl-1H-imidazole-5-carboxamide;
 1-[[2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl-2-methyl][1,1'-
 biphenyl]-4-yl]methyl]-4-ethyl-N-(1-methylethyl)-2-propyl-1H-imidazole-5-
 20 carboxamide;
 N-(3,4-Dimethyl-5-isoxazolyl)-2'-ethoxymethyl-4'-[[[(6-ethyl-3-
 methoxy-2-methyl-4-pyridinyl)oxy]methyl] [1,1'-biphenyl]-2-sulfonamide;
 4'-[(5-Acetyl-4-ethyl-2-propylimidazol-1-yl)methyl]-N-(3,4-dimethyl-
 5-isoxazolyl)-2'-methyl[1,1'-biphenyl]-2-sulfonamide;
 25 N-(3,4-Dimethyl-5-isoxazolyl)-4'-[[[(6-ethyl-3-methoxy-2-methyl-4-
 pyridinyl)oxy]methyl]-2'-propyl [1,1'-biphenyl]-2-sulfonamide;
 (S)-2-[N-[2'-[[N-(5-Methyl-3-isoxazolyl)amino]sulfonyl][1,1'-
 biphenyl]-4-yl]methyl]-N-(1-oxopentyl)amino]-3,N-dimethylbutanamide;
 (S)-2-[N-[2'-[[N-(4-Bromo-5-methyl-3-isoxazolyl)amino]sulfonyl][1,1'-
 30 biphenyl]-4-yl]methyl]-N-(1-oxopentyl)amino]-3,N-dimethylbutanamide;
 4'-[(2-Ethylquinolin-4-yl)oxymethyl]-N-(5-methylisoxazol-3-yl)

- [1,1'-biphenyl]-2-sulfonamide;
4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4-bromo-5-methyl-3-isoxazolyl)-2'-[(3,3-dimethyl-2-oxo-1-pyrrolidiny)methyl][1,1'-biphenyl]-2-sulfonamide;
- 5 N-[[4-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-[[4,5-dimethyl-3-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-2-yl)methyl]-N-methyl-2-pyrazinecarboxamide;
- N-[[2'-[(4,5-Dimethyl-3-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N-(1-oxopentyl)-L-valine;
- 10 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-2'-cyano-N-(4,5-dimethyl-3-isoxazolyl)[1,1'-biphenyl]-2-sulfonamide;
- (S)-N-[[2'-[[4,5-Dimethyl-3-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N-(1-piperidiny)carbonylpropyl]pentanamide;
- and salts, enantiomers, diastereomers and metabolites thereof.

15

23. A method for the treatment of an endothelin-dependent or angiotensin II-dependent disorder (such as hypertension, pulmonary hypertension, primary pulmonary hypertension, low renin hypertension, male erectile dysfunction, male or female sexual dysfunction, heart
- 20 failure, atherosclerosis, restenosis, endotoxemia, inhibition of cell growth, cancer, migraine, asthma, ischemia, subarachnoid hemorrhage, benign prostatic hypertrophy, renal glomerular or mesangial cell disorders, acute or chronic renal failure, chronic obstructive pulmonary disease, pain associated with prostate cancer, organ damage associated with the cell
- 25 proliferative effects of endothelin, general morbidity and mortality associated with endothelin-dependent or angiotensin II-depedendent disorders, diabetic nephropathy, and dementia), comprising the step of administering to a subject in need thereof an amount effective therefor of at least one compound of claim 1.

30

24. A pharmaceutical composition for the treatment of an endothelin-dependent or angiotensin II-dependent disorder, comprising a pharmaceutically acceptable vehicle or diluent and at least one compound of claim 1 in an amount effective therefor.

5

25. A compound of claim 1, wherein said compound is selected from the group consisting of:

N-(4,5-Dimethyl-3-isoxazolyl)-4'-[[[6-ethyl-3-methoxy-2-methyl-4-pyridinyl)oxy]methyl]-2'-ethyl [1,1'-biphenyl]-2-sulfonamide;

10 N-(3,4-Dimethyl-5-isoxazolyl)-4'-[[[6-ethyl-3-methoxy-2-methyl-4-pyridinyl)oxy]methyl]-2'-ethyl [1,1'-biphenyl]-2-sulfonamide;

N-(4,5-Dimethyl-3-isoxazolyl)-4'-[[[3-methoxy-2-methyl-6-propyl-4-pyridinyl)oxy]methyl]-2'-ethyl [1,1'-biphenyl]-2-sulfonamide;

15 N-(3,4-Dimethyl-5-isoxazolyl)-4'-[[[3-methoxy-2-methyl-6-propyl-4-pyridinyl)oxy]methyl]-2'-ethyl [1,1'-biphenyl]-2-sulfonamide;

N-(4,5-Dimethyl-3-isoxazolyl)-4'-[[[3-methoxy-2-methyl-6-propyl-4-pyridinyl)oxy]methyl]-2'-(ethoxymethyl) [1,1'-biphenyl]-2-sulfonamide;

N-(3,4-Dimethyl-5-isoxazolyl)-4'-[[[3-methoxy-2-methyl-6-propyl-4-pyridinyl)oxy]methyl]-2'-(ethoxymethyl) [1,1'-biphenyl]-2-sulfonamide;

20 N-(4,5-Dimethyl-3-isoxazolyl)-4'-[[[3-methoxy-2-methyl-6-propyl-4-pyridinyl)oxy]methyl]-2'-propyl [1,1'-biphenyl]-2-sulfonamide;

N-(3,4-Dimethyl-5-isoxazolyl)-4'-[[[3-methoxy-2-methyl-6-propyl-4-pyridinyl)oxy]methyl]-2'-propyl [1,1'-biphenyl]-2-sulfonamide;

25 N-(4,5-Dimethyl-3-isoxazolyl)-4'-[[[3-ethoxy-6-ethyl-2-methyl-4-pyridinyl)oxy]methyl]-2'-ethyl [1,1'-biphenyl]-2-sulfonamide;

N-(3,4-Dimethyl-5-isoxazolyl)-4'-[[[3-ethoxy-6-ethyl-2-methyl-4-pyridinyl)oxy]methyl]-2'-ethyl [1,1'-biphenyl]-2-sulfonamide;

N-(4,5-Dimethyl-3-isoxazolyl)-4'-[[[3-ethoxy-6-ethyl-2-methyl-4-pyridinyl)oxy]methyl]-2'-ethoxymethyl [1,1'-biphenyl]-2-sulfonamide;

30 N-(3,4-Dimethyl-5-isoxazolyl)-4'-[[[3-ethoxy-6-ethyl-2-methyl-4-pyridinyl)oxy]methyl]-2'-ethoxymethyl [1,1'-biphenyl]-2-sulfonamide;

N-(4,5-Dimethyl-3-isoxazolyl)-4'-[[[(3-ethoxy-6-ethyl-2-methyl-4-pyridinyl)oxy]methyl]-2'-propyl [1,1'-biphenyl]-2-sulfonamide;

N-(3,4-Dimethyl-5-isoxazolyl)-4'-[[[(3-ethoxy-6-ethyl-2-methyl-4-pyridinyl)oxy]methyl]-2'-propyl [1,1'-biphenyl]-2-sulfonamide;

5 N-(4,5-Dimethyl-3-isoxazolyl)-4'-[[[(3-ethoxy-2-methyl-6-propyl-4-pyridinyl)oxy]methyl]-2'-ethyl [1,1'-biphenyl]-2-sulfonamide;

N-(3,4-Dimethyl-5-isoxazolyl)-4'-[[[(3-ethoxy-2-methyl-6-propyl-4-pyridinyl)oxy]methyl]-2'-ethyl [1,1'-biphenyl]-2-sulfonamide;

10 N-(4,5-Dimethyl-3-isoxazolyl)-4'-[[[(3-ethoxy-2-methyl-6-propyl-4-pyridinyl)oxy]methyl]-2'-ethoxymethyl [1,1'-biphenyl]-2-sulfonamide;

N-(3,4-Dimethyl-5-isoxazolyl)-4'-[[[(3-ethoxy-2-methyl-6-propyl-4-pyridinyl)oxy]methyl]-2'-ethoxymethyl [1,1'-biphenyl]-2-sulfonamide;

N-(4,5-Dimethyl-3-isoxazolyl)-4'-[[[(3-ethoxy-2-methyl-6-propyl-4-pyridinyl)oxy]methyl]-2'-propyl [1,1'-biphenyl]-2-sulfonamide;

15 N-(3,4-Dimethyl-5-isoxazolyl)-4'-[[[(3-ethoxy-2-methyl-6-propyl-4-pyridinyl)oxy]methyl]-2'-propyl [1,1'-biphenyl]-2-sulfonamide;

N-(4,5-Dimethyl-3-isoxazolyl)-4'-[[[(2,6-dimethyl-3-ethoxy-4-pyridinyl)oxy]methyl]-2'-ethyl [1,1'-biphenyl]-2-sulfonamide;

20 N-(3,4-Dimethyl-5-isoxazolyl)-4'-[[[(2,6-dimethyl-3-ethoxy-4-pyridinyl)oxy]methyl]-2'-ethyl [1,1'-biphenyl]-2-sulfonamide;

N-(4,5-Dimethyl-3-isoxazolyl)-4'-[[[(2,6-dimethyl-3-ethoxy-4-pyridinyl)oxy]methyl]-2'-ethoxymethyl [1,1'-biphenyl]-2-sulfonamide;


N-(3,4-Dimethyl-5-isoxazolyl)-4'-[[[(2,6-dimethyl-3-ethoxy-4-pyridinyl)oxy]methyl]-2'-ethoxymethyl [1,1'-biphenyl]-2-sulfonamide;

25 N-(4,5-Dimethyl-3-isoxazolyl)-4'-[[[(2,6-dimethyl-3-ethoxy-4-pyridinyl)oxy]methyl]-2'-propyl [1,1'-biphenyl]-2-sulfonamide;


N-(3,4-Dimethyl-5-isoxazolyl)-4'-[[[(2,6-dimethyl-3-ethoxy-4-pyridinyl)oxy]methyl]-2'-propyl [1,1'-biphenyl]-2-sulfonamide;

and salts, enantiomers, diastereomers and metabolites thereof.

26. A compound of formula LXX, or an enantiomer, diastereomer, salt or metabolite thereof



- 5 wherein R_2 , R_{101} , R_{102} , R_{103} , R_{104} are as defined in claim 1;
 R_3 is isoxazol-5-yl or isoxazol-3-yl independently substituted with two
substituents selected from alkyl or halogen; and
 R_1 is any group such that the resulting compound demonstrates affinity
(IC₅₀) for both the AT₁ receptor and ET_A receptor of less than 5
10 micromolar at both receptors.



27. A compound of claim 2, wherein R_1 is M.
28. A compound of claim 27, wherein R_2 is alkyl, haloalkyl,
alkoxyalkyl or haloalkoxyalkyl and R_{101} , R_{102} , R_{103} , R_{104} are each
15 independently hydrogen, halogen, or alkyl.

29. A compound of claim 27, wherein R_2 is -CH₂Y.

30. A compound of claim 29, wherein Y is Q.

20

31. A crystalline form of 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-(ethoxymethyl) [1,1'-biphenyl]-2-sulfonamide having a melting point of about 148°C.

32. A compound of claim 1 selected from

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-3'-chloro-
N-(4,5-dimethyl-3-isoxazolyl)-5-methoxy[1,1'-biphenyl]-2-sulfonamide;

5 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-
dimethyl-3-isoxazolyl)-3'-fluoro-4-methoxy[1,1'-biphenyl]-2-sulfonamide;

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-5'-chloro-
N-(4,5-dimethyl-3-isoxazolyl)-2'-fluoro-4-methoxy[1,1'-biphenyl]-2-
sulfonamide;

10 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-3'-chloro-
N-(4,5-dimethyl-3-isoxazolyl)-4-methoxy[1,1'-biphenyl]-2-sulfonamide;

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-
dimethyl-3-isoxazolyl)-4-methoxy[1,1'-biphenyl]-2-sulfonamide;

15 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-
dimethyl-3-isoxazolyl)-2'-(ethoxymethyl)-4-methoxy[1,1'-biphenyl]-2-
sulfonamide;

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-
dimethyl-3-isoxazolyl)-3'-fluoro-5-methoxy[1,1'-biphenyl]-2-sulfonamide;

20 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-5'-chloro-
N-(4,5-dimethyl-3-isoxazolyl)-2'-fluoro-5-methoxy[1,1'-biphenyl]-2-
sulfonamide;

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-
dimethyl-3-isoxazolyl)-5-methoxy[1,1'-biphenyl]-2-sulfonamide;

25 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-
dimethyl-3-isoxazolyl)-2'-(ethoxymethyl)-5-methoxy[1,1'-biphenyl]-2-
sulfonamide;

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-
dimethyl-3-isoxazolyl)-3'-fluoro-4,5-dimethoxy[1,1'-biphenyl]-2-
sulfonamide;

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-5'-chloro-
N-(4,5-dimethyl-3-isoxazolyl)-2'-fluoro-4,5-dimethoxy[1,1'-biphenyl]-2-
sulfonamide;

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-3'-chloro-
5 N-(4,5-dimethyl-3-isoxazolyl)-4,5-dimethoxy[1,1'-biphenyl]-2-sulfonamide;

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-
dimethyl-3-isoxazolyl)-4,5-dimethoxy[1,1'-biphenyl]-2-sulfonamide;

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-
dimethyl-3-isoxazolyl)-2'-(ethoxymethyl)-4,5-dimethoxy[1,1'-biphenyl]-2-
10 sulfonamide;

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-5'-chloro-
N-(4,5-dimethyl-3-isoxazolyl)-2'-fluoro[1,1'-biphenyl]-2-sulfonamide;

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-3'-chloro-
N-(4,5-dimethyl-3-isoxazolyl)[1,1'-biphenyl]-2-sulfonamide;

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-
dimethyl-3-isoxazolyl)-4-methoxy-3'-methyl[1,1'-biphenyl]-2-sulfonamide;

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-
dimethyl-3-isoxazolyl)-3'-fluoro-4-methoxy[1,1'-biphenyl]-2-sulfonamide;

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-
20 dimethyl-3-isoxazolyl)-3'-methyl[1,1'-biphenyl]-2-sulfonamide;

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-
dimethyl-3-isoxazolyl)-5-fluoro[1,1'-biphenyl]-2-sulfonamide;

N²-(Cyclopropylcarbonyl)-N²-[[2'-[(4,5-dimethyl-3-
isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N-methyl-L-
25 valinamide;

N²-[[2'-[(4,5-Dimethyl-3-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-
yl)methyl]-N,3-dimethyl-N²-(1-oxobutyl)-L-valinamide;

N²-[[2'-[(4,5-Dimethyl-3-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-
yl)methyl]-N-methyl-N²-(2-methyl-1-oxopropyl)-L-valinamide;

N²-(Cyclopentylcarbonyl)-N²-[[2'-[[[(4,5-dimethyl-3-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-methyl-L-valinamide;

5 N²-[[2'-[[[(4,5-Dimethyl-3-isoxazolyl)amino]sulfonyl]-3-fluoro[1,1'-biphenyl]-4-yl]methyl]-N-methyl-N²-(1-oxobutyl)-L-valinamide;

N²-[[2'-[[[(4,5-Dimethyl-3-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-(1-methylethyl)-N²-(1-oxobutyl)-L-valinamide;

N²-[[2'-[[[(4,5-Dimethyl-3-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-(2-methoxyethyl)-N²-(1-oxobutyl)-L-valinamide;

10 N-(Cyclopropylmethyl)-N²-[[2'-[[[(4,5-dimethyl-3-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N²-(1-oxobutyl)-L-valinamide;

N²-[[2'-[[[(4,5-Dimethyl-3-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N²-(1-oxobutyl)-N-(3-pyridinyl)-L-valinamide;

15 N²-[[2'-[[[(4,5-Dimethyl-3-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-methyl-N²-(1-oxopentyl)-L-valinamide;

N-Methyl-N²-[[2'-[[[(5-methyl-3-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N²-(1-oxopentyl)-L-valinamide;

20 N²-[[2'-[[[(4,5-Dimethyl-3-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-ethyl-N²-(1-oxobutyl)-L-valinamide;

N²-[[2'-[[[(4,5-Dimethyl-3-isoxazolyl)amino]sulfonyl]-5'-fluoro[1,1'-biphenyl]-4-yl]methyl]-N-methyl-N²-(1-oxobutyl)-L-valinamide;

N-(4,5-Dimethyl-3-isoxazolyl)-4'-[[[(2,6-dimethyl-3-methoxy-4-pyridinyl)oxy]methyl][1,1'-biphenyl]-2-sulfonamide;

25 N-(4,5-Dimethyl-3-isoxazolyl)-4'-[[[(2,6-dimethyl-3-methoxy-4-pyridinyl)oxy]methyl]-3'-fluoro[1,1'-biphenyl]-2-sulfonamide;

N-(4,5-Dimethyl-3-isoxazolyl)-4'-[(1,4,5,6,7,8-hexahydro-8-oxo-2-propyl-1-cycloheptimidazolyl)methyl][1,1'-biphenyl]-2-sulfonamide;

30 N-(4,5-Dimethyl-3-isoxazolyl)-3'-fluoro-4'-[(1,4,5,6,7,8-hexahydro-8-oxo-2-propyl-1-cycloheptimidazolyl)methyl][1,1'-biphenyl]-2-sulfonamide;

N-(4,5-Dimethyl-3-isoxazolyl)-5-fluoro-4'-[(1,4,5,6,7,8-hexahydro-8-oxo-2-propyl-1-cycloheptimidazolyl)methyl][1,1'-biphenyl]-2-sulfonamide;

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-(hydroxymethyl)[1,1'-biphenyl]-2-sulfonamide;

5 N-[[2'-[(4,5-Dimethyl-3-isoxazolyl)amino]sulfonyl]-2-(ethoxymethyl)[1,1'-biphenyl]-4-yl)methyl]-1-[(1-oxopentyl)amino]cyclopentanamide;

and salts, enantiomers, diastereomers and metabolites thereof.

10 33. A compound of claim 1 selected from

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-3'-chloro-N-(3,4-dimethyl-5-isoxazolyl)-5-methoxy[1,1'-biphenyl]-2-sulfonamide;

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-3'-fluoro-4-methoxy[1,1'-biphenyl]-2-sulfonamide;

15 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-5'-chloro-N-(3,4-dimethyl-5-isoxazolyl)-2'-fluoro-4-methoxy[1,1'-biphenyl]-2-sulfonamide;

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-3'-chloro-N-(3,4-dimethyl-5-isoxazolyl)-4-methoxy[1,1'-biphenyl]-2-sulfonamide;

20 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-4-methoxy[1,1'-biphenyl]-2-sulfonamide;

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-(ethoxymethyl)-4-methoxy[1,1'-biphenyl]-2-sulfonamide;

25 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-3'-fluoro-5-methoxy[1,1'-biphenyl]-2-sulfonamide;

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-5'-chloro-N-(3,4-dimethyl-5-isoxazolyl)-2'-fluoro-5-methoxy[1,1'-biphenyl]-2-sulfonamide;

30 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-5-methoxy[1,1'-biphenyl]-2-sulfonamide;

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-(ethoxymethyl)-5-methoxy[1,1'-biphenyl]-2-sulfonamide;

5 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-3'-fluoro-4,5-dimethoxy[1,1'-biphenyl]-2-sulfonamide;

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-5'-chloro-N-(3,4-dimethyl-5-isoxazolyl)-2'-fluoro-4,5-dimethoxy[1,1'-biphenyl]-2-sulfonamide;

10 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-3'-chloro-N-(3,4-dimethyl-5-isoxazolyl)-4,5-dimethoxy[1,1'-biphenyl]-2-sulfonamide;

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-4,5-dimethoxy[1,1'-biphenyl]-2-sulfonamide;

15 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-(ethoxymethyl)-4,5-dimethoxy[1,1'-biphenyl]-2-sulfonamide;

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-5'-chloro-N-(3,4-dimethyl-5-isoxazolyl)-2'-fluoro[1,1'-biphenyl]-2-sulfonamide;

20 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-3'-chloro-N-(3,4-dimethyl-5-isoxazolyl)[1,1'-biphenyl]-2-sulfonamide;

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-4-methoxy-3'-methyl[1,1'-biphenyl]-2-sulfonamide;

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-3'-fluoro-4-methoxy[1,1'-biphenyl]-2-sulfonamide;

25 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-3'-methyl[1,1'-biphenyl]-2-sulfonamide;

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-5-fluoro[1,1'-biphenyl]-2-sulfonamide;

30 N²-(Cyclopropylcarbonyl)-N²-[[2'-[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl)methyl]-N-methyl-L-valinamide;

N^2 -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N,3-dimethyl- N^2 -(1-oxobutyl)-L-valinamide;

N^2 -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-methyl- N^2 -(2-methyl-1-oxopropyl)-L-valinamide;

5 N^2 -(Cyclopentylcarbonyl)- N^2 -[[2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-methyl-L-valinamide;

N^2 -[[2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl]-3-fluoro[1,1'-biphenyl]-4-yl]methyl]-N-methyl- N^2 -(1-oxobutyl)-L-valinamide;

10 N^2 -[[2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-(1-methylethyl)- N^2 -(1-oxobutyl)-L-valinamide;

N^2 -[[2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-(2-methoxyethyl)- N^2 -(1-oxobutyl)-L-valinamide;

15 N-(Cyclopropylmethyl)- N^2 -[[2'-[[[(3,4-dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]- N^2 -(1-oxobutyl)-L-valinamide;

N^2 -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]- N^2 -(1-oxobutyl)-N-(3-pyridinyl)-L-valinamide;

20 N^2 -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-methyl- N^2 -(1-oxopentyl)-L-valinamide;

N-Methyl- N^2 -[[2'-[[[(3-methyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]- N^2 -(1-oxopentyl)-L-valinamide;

N^2 -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]methyl]-N-ethyl- N^2 -(1-oxobutyl)-L-valinamide;

25 N^2 -[[2'-[[[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl]-5'-fluoro[1,1'-biphenyl]-4-yl]methyl]-N-methyl- N^2 -(1-oxobutyl)-L-valinamide;

N-(3,4-Dimethyl-5-isoxazolyl)-4'-[[[(2,6-dimethyl-3-methoxy-4-pyridinyl)oxy]methyl][1,1'-biphenyl]-2-sulfonamide;

30 N-(3,4-Dimethyl-5-isoxazolyl)-4'-[[[(2,6-dimethyl-3-methoxy-4-pyridinyl)oxy]methyl]-3'-fluoro[1,1'-biphenyl]-2-sulfonamide;

N-(3,4-Dimethyl-5-isoxazolyl)-4'-[(1,4,5,6,7,8-hexahydro-8-oxo-2-propyl-1-cycloheptimidazolyl)methyl][1,1'-biphenyl]-2-sulfonamide;

N-(3,4-Dimethyl-5-isoxazolyl)-3'-fluoro-4'-[(1,4,5,6,7,8-hexahydro-8-oxo-2-propyl-1-cycloheptimidazolyl)methyl][1,1'-biphenyl]-2-sulfonamide;

5 N-(3,4-Dimethyl-5-isoxazolyl)-5-fluoro-4'-[(1,4,5,6,7,8-hexahydro-8-oxo-2-propyl-1-cycloheptimidazolyl)methyl][1,1'-biphenyl]-2-sulfonamide;

4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(3,4-dimethyl-5-isoxazolyl)-2'-(hydroxymethyl)[1,1'-biphenyl]-2-sulfonamide;

10 N-[[2'-[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl]-2-(ethoxymethyl)[1,1'-biphenyl]-4-yl)methyl]-1-[(1-oxopentyl)amino]cyclopentanamide;

and salts, enantiomers, diastereomers and metabolites thereof.

34. The compound 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-(2-fluoroethoxymethyl) [1,1'-biphenyl]-2-sulfonamide or a salt, enantiomer, diastereomer or metabolite thereof.


35. The compound 4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-propyl [1,1'-biphenyl]-2-sulfonamide or a salt, enantiomer, diastereomer or metabolite thereof.

36. The compound N-(4,5-Dimethyl-3-isoxazolyl)-2'-ethoxymethyl-4'-[[3-methoxy-2,6-dimethyl-4-pyridinyl]oxy]methyl [1,1'-biphenyl]-2-sulfonamide or a salt, enantiomer, diastereomer or metabolite thereof.

37. The compound N-(4,5-Dimethyl-3-isoxazolyl)-2'-[(2-fluoroethoxy)methyl]-4'-[[3-methoxy-2,6-dimethyl-4-pyridinyl]oxy]methyl [1,1'-biphenyl]-2-sulfonamide or a salt, enantiomer, diastereomer or metabolite thereof.

30

38. The compound N-(4,5-Dimethyl-3-isoxazolyl)-4'-[[[(3-methoxy-2,6-dimethyl-4-pyridinyl)oxy]methyl]-2'-propyl [1,1'-biphenyl]-2-sulfonamide or a salt, enantiomer, diastereomer or metabolite thereof.



5 39. A compound of claim 2, where in R_1 is D .

40. A compound of claim 39, wherein R_2 is hydrogen, alkyl, haloalkyl, alkoxyalkyl or haloalkoxyalkyl and R_{101} , R_{102} , R_{103} , R_{104} are each independently hydrogen, halogen, or alkyl.

10

41. A compound of claim 39 wherein R_2 is $-\text{CH}_2\text{Y}$.

42. A compound of claim 41, wherein Y is Q.

15

43. A compound of claim 1 wherein R_3 is other than pyridyl when R_1 is A.

44. The pharmaceutical composition of claim 46 further
 20 comprising at least one ACE inhibitor (such as captopril, zofenopril, fosinopril, ceranapril, alacepril, enalapril, delapril, pentopril, quinapril, ramipril, or lisinopril), vasopepsidase inhibitor (such as omapatrilat or gemopatrilat), HMG CoA reductase inhibitor (such as pravastatin, lovastatin, atorvastatin, simvastatin, NK-104 or ZD-
 25 4522), anti-platelet agent (such as clopidigrel, ticlopidine, CS-747 or aspirin), anti-diabetic agent (such as biguanides or biguanide/glyburide combinations), beta-adrenergic agent (such as

carvedilol or metoprolol), or mineralocorticoid receptor antagonist (such as spironolactone or eplerenone).

45. The compound (+)-4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-(ethoxymethyl) [1,1'-biphenyl]-2-sulfonamide or metabolites and salts thereof.

46. The compound (-)-4'-[(2-Butyl-4-oxo-1,3-diazaspiro[4.4]non-1-en-3-yl)methyl]-N-(4,5-dimethyl-3-isoxazolyl)-2'-(ethoxymethyl) [1,1'-biphenyl]-2-sulfonamide or metabolites and salts thereof.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
21 June 2001 (21.06.2001)

PCT

(10) International Publication Number
WO 01/44239 A3

(51) International Patent Classification⁷: **C07D 413/12**,
413/14, 403/14, 487/04, 261/16, 401/12, 417/12, A61K
31/42, A61P 9/12

08822 (US). GU, Zhengxiang [CN/US]; 92 York Road,
Princeton, NJ 08540 (US).

(21) International Application Number: PCT/US00/33730

(74) Agents: **ALGIERI, Aldo, A.** et al.; Bristol-Myers Squibb
Co., P.O. Box 4000, Lawrenceville-Princeton Road,
Princeton, NJ 08543 (US).

(22) International Filing Date:
13 December 2000 (13.12.2000)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE,
DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID,
IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT,
LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL,
PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ,
UA, UG, US, UZ, VN, YU, ZA, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/464,037 15 December 1999 (15.12.1999) US
09/481,197 11 January 2000 (11.01.2000) US
09/513,779 25 February 2000 (25.02.2000) US
09/604,322 26 June 2000 (26.06.2000) US
09/643,640 22 August 2000 (22.08.2000) US

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **BRIS-
TOL-MYERS SQUIBB CO.** [US/US]; P.O. Box 4000,
Lawrenceville-Princeton Road, Princeton, NJ 08543-4000
(US).

Published:
— with international search report

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **MURUGESAN,**
Natesan [US/US]; 42 Millbrook Drive, Princeton Junc-
tion, NJ 08550 (US). **TELLEW, John, E.** [US/US]; 28
Avalon Road, Pennington, NJ 08534 (US). **MACOR,**
John, E. [US/US]; 258 Old Farm Road, Flemington, NJ

(88) Date of publication of the international search report:
1 November 2001

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: BIPHENYL SULFONAMIDES AS DUAL ANGIOTENSIN ENDOTHELIN RECEPTOR ANTAGONISTS

(57) Abstract: Novel biphenyl sulfonamide compounds which are combined angiotensin and endothelin receptor antagonists are claimed along with methods of using such compounds in the treatment of conditions such as hypertension and other diseases, as well as pharmaceutical compositions containing such compounds.

WO 01/44239 A3

INTERNATIONAL SEARCH REPORT

International Application No

PC US 00/33730

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D413/12 C07D413/14 C07D403/14 C07D487/04 C07D261/16
 C07D401/12 C07D417/12 A61K31/42 A61P9/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5 612 359 A (MURUGESAN) 18 March 1997 (1997-03-18) column 4, line 58 -column 6, line 2; claims; example 15 ---	1-46
X	US 5 846 990 A (MURUGESAN) 8 December 1998 (1998-12-08) claims; examples ---	26
Y	US 5 780 473 A (MURUGESAN ET. AL.) 14 July 1998 (1998-07-14) column 6, line 33 -column 7, line 45; claims; examples ---	1-46
Y	US 5 514 696 A (MURUGESAN ET. AL.) 7 May 1996 (1996-05-07) column 5, line 13 -column 6, line 2; claims; examples --- -/--	1-46

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

15 May 2001

Date of mailing of the international search report

29.05.01

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Helps, I

INTERNATIONAL SEARCH REPORT

International Application No

PC IS 00/33730

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	GB 2 264 710 A (MERCK & CO) 8 September 1993 (1993-09-08) claims; examples ---	1-46
X	WO 98 33780 A (BRISTOL MYERS SQUIBB) 6 August 1998 (1998-08-06) claims; examples ---	1,26,39
X	WO 98 04260 A (BRISTOL MYERS SQUIBB) 5 February 1998 (1998-02-05) claims; examples ---	26
Y	US 5 554 625 A (RIVERO ET. AL.) 10 September 1996 (1996-09-10) claims; examples ---	1
Y	US 5 399 578 A (BÜHLMAYER ET. AL.) 21 March 1998 (1998-03-21) claims; examples ---	1,39
Y	US 5 260 328 A (MARKWALDER ET. AL.) 9 November 1993 (1993-11-09) column 2, line 1 - line 38; claims; examples ---	1,39
Y	US 5 760 038 A (MURUGESAN ET. AL.) 2 June 1998 (1998-06-02) claims; examples -----	1

INTERNATIONAL SEARCH REPORT

international application No.
PCT/US 00/33730

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claim 23 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☒ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1(part),2-25,27-31,32(part),33-37,39-46

Compounds of formula I in which R1 is A, B, E, G - N

2. Claims: 1(part),32(part),38

Compounds of formula I in which R1 is D

3. Claim : 1(part)

Compounds of formula I in which R1 is F

4. Claim : 1(part)

Compounds of formula I in which R1 is O

5. Claim : 26

Compounds of formula LXX.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PC US 00/33730

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5612359 A	18-03-1997	US 5827869 A	27-10-1998
		AU 699138 B	26-11-1998
		AU 3026195 A	07-03-1996
		CA 2155447 A	27-02-1996
		CN 1128262 A,B	07-08-1996
		CZ 9502163 A	13-03-1996
		EP 0702012 A	20-03-1996
		FI 954008 A	27-02-1996
		HU 74097 A	28-11-1996
		IL 114829 A	30-11-1999
		JP 8183786 A	16-07-1996
		NO 953361 A	27-02-1996
		NZ 272812 A	24-11-1997
		PL 310191 A	04-03-1996
		SG 46142 A	20-02-1998
		BR 1100659 A	13-06-2000
US 5846990 A	08-12-1998	AU 720458 B	01-06-2000
		AU 2209897 A	02-09-1997
		CA 2240043 A	21-08-1997
		EP 0921800 A	16-06-1999
		WO 9729748 A	21-08-1997
		ZA 9701423 A	19-08-1998
US 5780473 A	14-07-1998	US 5760038 A	02-06-1998
		AU 3661397 A	20-02-1998
		WO 9804260 A	05-02-1998
		AU 708926 B	19-08-1999
		AU 4331496 A	15-08-1996
		CA 2168154 A	07-08-1996
		CN 1138581 A,B	25-12-1996
		CZ 9600292 A	11-09-1996
		EP 0725067 A	07-08-1996
		FI 960516 A	07-08-1996
		HU 9600256 A	28-10-1997
		JP 8245600 A	24-09-1996
		NO 960473 A	07-08-1996
		NZ 280879 A	24-11-1997
		PL 312650 A	19-08-1996
		SG 46218 A	20-02-1998
US 5514696 A	07-05-1996	US 6107320 A	22-08-2000
		AT 149155 T	15-03-1997
		AU 3838293 A	11-11-1993
		CA 2095174 A	07-11-1993
		CN 1084515 A	30-03-1994
		CY 2016 A	20-02-1998
		DE 69308229 D	03-04-1997
		DE 69308229 T	17-07-1997
		DK 569193 T	17-03-1997
		EP 0569193 A	10-11-1993
		ES 2100459 T	16-06-1997
		FI 932052 A	07-11-1993
		GR 3023421 T	29-08-1997
		HK 98197 A	08-08-1997
		HU 65751 A,B	28-07-1994
		IL 105481 A	05-12-1996
		JP 6049046 A	22-02-1994

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PC JS 00/33730

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5514696 A		MX 9302618 A NO 303065 B NZ 247440 A PL 298828 A RU 2133742 C ZA 9302914 A	31-05-1994 25-05-1998 27-04-1995 27-12-1993 27-07-1999 12-11-1993
GB 2264710 A	08-09-1993	NONE	
WO 9833780 A	06-08-1998	AU 730111 B AU 6134998 A BR 9807098 A CN 1244862 T CZ 9902362 A EP 0996618 A NO 993691 A PL 334773 A US 6043265 A	22-02-2001 25-08-1998 18-04-2000 16-02-2000 17-11-1999 03-05-2000 29-07-1999 13-03-2000 28-03-2000
WO 9804260 A	05-02-1998	US 5780473 A AU 3661397 A	14-07-1998 20-02-1998
US 5554625 A	10-09-1996	AU 4770893 A EP 0668864 A JP 7509464 T WO 9402142 A	14-02-1994 30-08-1995 19-10-1995 03-02-1994
US 5399578 A	21-03-1995	US 5965592 A AT 134624 T AU 644844 B AU 7115191 A CA 2036427 A CA 2232775 A CY 1978 A DE 59107440 D DK 443983 T EP 0443983 A ES 2084801 T FI 910747 A FI 980787 A GR 3019155 T HK 219996 A HU 61271 A IE 910548 A IL 97219 A JP 2749458 B JP 4235149 A KR 171409 B LU 90100 A LU 90362 A LV 5773 A MX 24598 A NO 304023 B NZ 237126 A PT 96799 A,B ZA 9101179 A	12-10-1999 15-03-1996 23-12-1993 22-08-1991 20-08-1991 20-08-1991 05-09-1997 04-04-1996 18-03-1996 28-08-1991 16-05-1996 20-08-1991 06-04-1998 31-05-1996 03-01-1997 28-12-1992 28-08-1991 08-12-1995 13-05-1998 24-08-1992 01-02-1999 25-09-1997 10-05-1999 20-12-1996 28-02-1994 12-10-1998 25-11-1994 31-10-1991 27-11-1991
US 5260328 A	09-11-1993	US 5352676 A	04-10-1994

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/33730

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5260328 A		AT 99934 T	15-01-1994
		AU 5416990 A	05-11-1990
		CA 2030530 A	07-10-1990
		DD 298478 A	27-02-1992
		DE 69005974 D	24-02-1994
		DE 69005974 T	05-05-1994
		WO 9011760 A	18-10-1990
		EP 0423269 A	24-04-1991
		HU 58518 A	30-03-1992
		JP 3505332 T	21-11-1991
		PT 93680 A	20-11-1990
		US 5196445 A	23-03-1993
		ZA 9002649 A	30-01-1991

US 5760038 A	02-06-1998	AU 708926 B	19-08-1999
		AU 4331496 A	15-08-1996
		CA 2168154 A	07-08-1996
		CN 1138581 A,B	25-12-1996
		CZ 9600292 A	11-09-1996
		EP 0725067 A	07-08-1996
		FI 960516 A	07-08-1996
		HU 9600256 A	28-10-1997
		JP 8245600 A	24-09-1996
		NO 960473 A	07-08-1996
		NZ 280879 A	24-11-1997
		PL 312650 A	19-08-1996
		SG 46218 A	20-02-1998
		US 5780473 A	14-07-1998
